


NEET | **JEE**
CLASS XI-XII ESSENTIALS



ACE YOUR WAY **CBSE**
Class XI-XII

CHEMISTRY

India's #1
CHEMISTRY MONTHLY FOR
JEE (Main & Advanced) & NEET

t  **oday**

**CHEMISTRY
MUSING**

**MONTHLY
PRACTICE
PROBLEMS
(XI & XII)**

**CONCEPT
MAP**

mtg

Trust of more than
1 Crore Readers
Since 1982

**CONCEPT
BOOSTER**

**EXAMINER'S
MIND**



2017100009372



ALERT!

The **IIT Joint Admission Board (JAB)** has announced that the **Joint Entrance Examination (Advanced)** will go online from 2018.



Don't panic! ALLEN Career Institute has always helped students for their right exam preparation with best resources prepared by ALLEN's Outstanding Academic and R&D team. JEE Advanced is going Online and to equip you with Exact Simulation for the Online Exam Temperament, ALLEN brings to you the Most Trusted and Advanced Online Test Series Platform www.onlinetestseries.in. Don't left behind; thousands are already Online ! Benchmark your Preparation Today with ALLEN !



Compete with Largest Pool of Talented Students (Inclusive our Classroom Students)



Full syllabus 12 Major Online Tests @ ₹600



Practice on exact Simulation of Online Exam

To know more logon to www.onlinetestseries.in

CORPORATE OFFICE : "SANKALP", CP-6, Indra Vihar, Kota-324005 (Raj.)

Email: support@onlinetestseries.in | **Website**: www.allen.ac.in | dlp.allen.ac.in

"At LPU, we had industry-sponsored labs from Google, Oracle, SAP where we worked on real life projects. This industry experience made cracking job interviews much easier."

At LPU, I learnt to
Think BIG

SONIA (LPU Alumni)
Software Engineer

amazon USA



UNBEATABLE PLACEMENT RECORD

- Thousands of LPU students have got placed in top MNCs in USA, UK, Canada, Australia etc.
- 600+ Offers alone made by Ten Fortune 500 companies including Amazon, Xerox and HP to 2017 batch students.
- 51 Companies that recruit from IITs also recruit at LPU.
- 500+ Multinational & National Companies recruit LPU students.
- Record 610 Students hired by Cognizant in a single day, highest in North India till date.
- 100+ Selections each by Top Companies like Capgemini, Tech Mahindra, Amazon from 2017 batch.



LOVELY
PROFESSIONAL
UNIVERSITY

Transforming Education Transforming India

Toll Free : 1800-102-4430

FOLLOW US@    

Jalandhar-Delhi G.T. Road, Phagwara, Punjab | admissions@lpu.co.in | SMS LPU to 53030

Application Open for

L P U N E S T 2018
NATIONAL ENTRANCE AND SCHOLARSHIP TEST
FOR LPU ENGINEERING AND OTHER PROGRAMMES

B.Tech.
B.Tech. (Hons.)
Integrated B.Tech.-M.Tech.
Integrated B.Tech.-MBA

Streams available:

ECE[†], CSE, IT[†], ME, ME - Mechatronics[†], Aerospace Engineering[†], CE[†], EE[†], EEE[†], Electronics and Computer Engg.[†], Biotech. (BT)[†], Food Technology[†], Automobile Engg. (AE)[†], Chemical Engg. (CHE)[†], CHE - Petroleum[†], Biomedical Engineering (BE)[†].

^{*}streams/majors are not offered in Honors programmes.

[†]streams/majors are not offered in Integrated B.Tech.- M.Tech. programme.

[^]streams/majors are not offered in Integrated B.Tech.- MBA programme.

Online Registration Open

Date of Examination : **10th - 30th April 2018**

For last date and other details visit : www.lpu.in



**Scholarships up to ₹ 7.5 Lakh
Per Student**

EXCEL ACADAMICS®

Bengaluru

A Premier Institute in KARNATAKA

Integrated PUC | PUC+CET | PUC+IIT-JEE | PUC+NEET

NEET

1 ALL INDIA RANK

18 44 97

SHIVANANDA S
H. T. No. : 61718963
All India Quota Cat.

PRAJWAL K
H.T. No. : 61717312
All India Quota Cat.

CHETHAN. M
H. T. No. : 61718306
All India Quota Cat.

SHAMANTH M
H. T. No. : 903604299

CET Ranks

18 36 66

V. CHETAN

NAVYA N
H. T. No. : SH213

SACHIN V A
H. T. No. : WB101

JEE(Main) Score

222 192 187

SACHIN VA

MADHU J N

SANJAY GOWDA M

227 181 172

DHEERAJ R

V CHETAN

NAVEEN RAJ

Helpline : 7676-91-7777, 7676-41-6666

CET, IIT-JEE (Main & Advanced), NEET, AAIMS, JIPMER, COMED-K, BITS, VIT, MAHE

REGISTRATION OPEN

<p>EXCLUSIVE NEET-2018</p> <p>New Batch</p> <p>PCB</p> <p>Classes with Daily test</p>	<p>K-CET / NEET / JEE CRASH COURSE 2018</p> <p>2nd PUC/12th/+2 Studying Students</p>	<p>46 Days Residential</p> <p>SUMMER COURSE</p> <p>1st PUC/11th, +1, Studying Students</p>	<p>1 Month Residential</p> <p>BRIDGE COURSE</p> <p>9th & 10th Studying Students (Maths & Science)</p>	<p>Fully Solved</p> <p>OFF-LINE TEST SERIES</p> <p>CET, IIT-JEE (Main & Advanced), NEET - 2018 & 2019</p> <p>₹ 3,540/- ONLY</p>	<p>Fully Solved</p> <p>ON-LINE TEST SERIES</p> <p>CET, IIT-JEE (Main & Advanced), NEET - 2018 & 2019</p> <p>₹ 7,080/- ONLY</p>
--	--	--	--	--	---

EXCEL ACADAMICS : #326, Opp. People Tree Hospital, Sheshadripuram College Road, Yelahanka New Town - 560064, Bangalore, KARNATAKA

Contact: 9535656277 / 9880286824 / 9900836461 / 9036357499

★ ★ ★ To get free Mock CET/ NEET/JEE SMS your complete postal address to 7676917777 ★ ★ ★

★ ★ ★ ★  **Separate Deluxe Hostel for Boys and Girls**  ★ ★ ★ ★

RESULT 2017

FELICITATION



Achievements

Academic Year		2014	2015	2016	2017
Selections	Medical	80	98	115	139
	Engineering	60	76	94	109

COURSES OFFERED

Two Years Integrated Programme

- ◆ NEET (Medical Entrance Exam.)
- ◆ IIT-JEE (Engineering Entrance Exam.)

One Year Integrated Programme

- ◆ NEET Repeater Batch

Test Series NEET/JEE (Online /Offline)

- ◆ **Special Online Distance Learning / Test Series Programme.**

- ◆ Tab with study material available at Lalit Tutorials.
- ◆ For Free Test Series - Download Lalit Eduapp from Google Play Store



- ◆ Well Equiped Digital Class Room
- ◆ Well Furnished Library & Laboratory



The Best Institute in Maharashtra for Medical & Engineering Entrance Exams ...

Kalpande Sir's



Toshniwal Lay-out, Akola (Maharashtra) Phone : 0724-2456 654, 7720 857 857

Opportunity For Faculties

Faculties	Min. Exp.	Min. Package / Annum
Senior	6 yr.	12 Lac
Junior	3 yr.	6 Lac

- ★ Special Incentives & Increments According to Performance
- ★ Special Reward to Faculty on Producing Best Ranks.

For Recruitment of Senior & Junior Faculties send your Resume to hrltmaharashtra@gmail.com



ACADEMIC BENEFITS*

- More than 800 Academic Hours & 500 Classes
- More than 15000 Academic Questions
- More than 100 Testing Hours

FINANCIAL BENEFITS*

- Upto Rs. 35000+ Saving on 1 Year Course Fee
- 50% Concession on Admission Form Fee
- Upto 90% Scholarship on Course Fee

* T&C Apply

ANNOUNCING SPECIAL SCHOLARSHIP PROGRAM FOR MERITORIOUS STUDENTS MOVING TO

CLASS 8th, 9th & 10th



- | | | |
|---|--|---|
| STEP 1
Complete ResoFAST Application form | STEP 2
Qualify ResoFAST followed by personal interview round | STEP 3
Pay once through Vardaan Scholarship & study till Class 12 th |
|---|--|---|

T&C Apply

ADMISSION ANNOUNCEMENT

Enroll Now for Academic Session 2018-19 @ Coaching Fee of 2017-18

CLASSROOM CONTACT PROGRAM FOR CLASSES



Target: JEE (Main+Advanced) | JEE (Main) AIMS/ NEET | Pre-foundation

Test Dates

26th November 2017
03th & 10th December 2017

Resonance Eduventures Limited

Registered & Corporate Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Rajasthan) - 324005
Tel. No.: 0744-3012100, 3012222, 6635555 | CIN: U80302RJ2007PLC024029

To know more: sms RESO at 56677 | website: contact@resonance.ac.in | e-mail: www.resonance.ac.in

Toll Free: 1800 258 5555 | [facebook.com/ResonanceEdu](https://www.facebook.com/ResonanceEdu) | twitter.com/ResonanceEdu | www.youtube.com/resowatch | blog.resonance.ac.in

CHEMISTRY today

Volume 26 No. 12 December 2017

Managing Editor

Mahabir Singh

Editor

Anil Ahlawat

Corporate Office:

Plot 99, Sector 44 Institutional area, Gurgaon -122 003 (HR).

Tel : 0124-6601200 e-mail : info@mtg.in website : www.mtg.in

Regd. Office:

406, Taj Apartment, Near Safdarjung Hospital, New Delhi - 110029.

CONTENTS

Class 11

NEET JEE Essentials	8
Examiner's Mind	23
Ace Your Way CBSE	31
MPP-8	39
Concept Map	42

Class 12

Concept Map	43
NEET JEE Essentials	44
Examiner's Mind	62
Ace Your Way CBSE	71
MPP-8	78

Competition Edge

Chemistry Musing Problem Set 53	82
Concept Booster	83
JEE Advanced Practice Problems	85
Chemistry Musing Solution Set 52	88

Subscribe online at www.mtg.in

	Individual Subscription Rates			Combined Subscription Rates			
	1 yr.	2 yrs.	3 yrs.		1 yr.	2 yrs.	3 yrs.
Mathematics Today	330	600	775	PCM	900	1500	1900
Chemistry Today	330	600	775	PCB	900	1500	1900
Physics For You	330	600	775	PCMB	1000	1800	2300
Biology Today	330	600	775				

Send D.D/M.O in favour of MTG Learning Media (P) Ltd.

Payments should be made directly to : MTG Learning Media (P) Ltd,

Plot No. 99, Sector 44, Gurgaon - 122003 (Haryana)

We have not appointed any subscription agent.

Printed and Published by Mahabir Singh on behalf of MTG Learning Media Pvt. Ltd. Printed at HT Media Ltd., B-2, Sector-63, Noida, UP-201307 and published at 406, Taj Apartment, Ring Road, Near Safdarjung Hospital, New Delhi - 110029.

Editor : Anil Ahlawat

Readers are advised to make appropriate thorough enquiries before acting upon any advertisements published in this magazine. Focus/Infocus features are marketing incentives. MTG does not vouch or subscribe to the claims and representations made by advertisers. All disputes are subject to Delhi jurisdiction only. Copyright © MTG Learning Media (P) Ltd.

All rights reserved. Reproduction in any form is prohibited.

NEET | JEE

ESSENTIALS

Class
XI

Maximise your chance of success, and high rank in NEET, JEE (Main and Advanced) by reading this column. This specially designed column is updated year after year by a panel of highly qualified teaching experts well-tuned to the requirements of these Entrance Tests.

Unit 6

The *p*-Block Elements (Group 13 and 14)

GROUP-13 ELEMENTS (BORON FAMILY)

INTRODUCTION

Group-13 of the periodic table consist of the following elements with general configuration ns^2np^1 . A new element is added to group-13 *i.e.*, nihonium (Nh), which is radioactive.

Element	Electronic configuration
B (5)	[He] $2s^2 2p^1$
Al (13)	[Ne] $3s^2 3p^1$
Ga (31)	[Ar] $3d^{10} 4s^2 4p^1$
In (49)	[Kr] $4d^{10} 5s^2 5p^1$
Tl (81)	[Xe] $4f^{14} 5d^{10} 6s^2 6p^1$
Nh (113)	[Rn] $5f^{14} 6d^{10} 7s^2 7p^1$

Physical Properties

Atomic and ionic radii

The atomic/ionic radius increases from B to Tl (exception $Al > Ga$).

Ionisation energy

I.E. decreases down the group. Successive *I.E.* increases in the order :

$$\Delta_i H_1 < \Delta_i H_2 < \Delta_i H_3$$

Electronegativity

Boron has the maximum electronegativity among the group-13 elements. Electronegativity first decreases from B to Al and then increases down the group.

Metallic character

B is a non-metal while Al is a metal and all other elements are almost metallic in character.

Density

As we move down the group, density increases.

Boiling point

Boiling points of group-13 elements follow a regular trend and decreases down the group.


Melting point

The melting points decreases from B to Ga and then increases from Ga to Tl.

The melting point of boron is very high because it has giant covalent polymer structure in both solid and liquid states.

mtg

Last-minute check on your NEET readiness



₹ 550



MTG's NEET Explorer helps students self-assess their readiness for success in NEET. Attempting the tests put together by MTG's experienced team of editors and experts strictly on the NEET pattern and matching difficulty levels, students can easily measure their preparedness for success. **Order now!**

HIGHLIGHTS:

- 10 Model Test Papers based on latest NEET syllabus
- Last 15 years' solved test papers of AIPMT / NEET
- Includes NEET 2017 solved paper
- Detailed solutions for self-assessment and to practice time management



Scan now with your smartphone or tablet*



Available at all leading book shops throughout India.
For more information or for help in placing your order:
Call 0124-6601200 or email: info@mtg.in

*Application to read QR codes required

Visit
www.mtg.in
for latest offers
and to buy
online!

Oxidation states


- B and Al show +3 oxidation states only but Ga, In and Tl show +1 and +3 oxidation states due to inert pair effect.

Inert pair effect

The two s-electrons in the outer shell tend to remain paired and will not participate in compound formation. This effect is called inert pair effect. The inert pair effect increases gradually in Ga, In and Tl compounds *e.g.*, Ga^+ compounds are unstable, In^+ compounds are moderately stable, whereas Tl^+ compounds are most stable.

Reactivity towards air

- $4M_{(s)} + 3O_{2(g)} \xrightarrow{\Delta} 2M_2O_{3(s)}$ ($M = \text{B, Al, Ga, In, Tl}$)
- $2M_{(s)} + N_{2(g)} \xrightarrow{\Delta} 2MN_{(s)}$ ($M = \text{B, Al}$)
- On moving down the group, the acidic character of oxides decreases and the basic character increases.

**Reactivity towards water**

- Boron is not affected by water or steam. Red hot boron decomposes steam.
 $2\text{B} + 3\text{H}_2\text{O} \longrightarrow \text{B}_2\text{O}_3 + 3\text{H}_2$
- Aluminium can decompose the steam.
 $2\text{Al} + 3\text{H}_2\text{O} \longrightarrow \text{Al}_2\text{O}_3 + 3\text{H}_2$
- Ga and In are attacked by water only in the presence of oxygen.
- Tl is attacked by moist air.
 $4\text{Tl} + 2\text{H}_2\text{O} + \text{O}_2 \longrightarrow 4\text{TlOH}$

Chemical Properties**Reactivity towards acids and alkalis**

- Boron does not react with non-oxidising acids while others react with acids *e.g.*,
 $2\text{Al}_{(s)} + 6\text{HCl}_{(aq)} \longrightarrow 2\text{Al}_{(aq)}^{3+} + 6\text{Cl}_{(aq)}^- + 3\text{H}_{2(g)}$
- Conc. HNO_3 makes Al passive due to the formation of thin layer of Al_2O_3 .
- Boron resist the action of alkalis upto 773 K.
 $2\text{B}_{(s)} + 6\text{KOH}_{(aq)} \xrightarrow{773\text{ K}} 2\text{K}_3\text{BO}_{3(s)} + 3\text{H}_{2(g)}$
- Al and Ga being amphoteric also react with aqueous alkalis.
 $2M_{(s)} + 2\text{NaOH}_{(aq)} + 6\text{H}_2\text{O}_{(l)} \longrightarrow 2\text{Na}^+[\text{M}(\text{OH})_4]^- + 3\text{H}_{2(g)}$ ($M = \text{Al, Ga}$)

Reactivity towards non-metals

- At high temperature, B reacts directly with all non-metals except H, Ge, Te and the noble gases.
 $4\text{B} + \text{C} \xrightarrow{\Delta} \text{B}_4\text{C}$
 $4\text{B} + 3\text{O}_2 \xrightarrow{\Delta} 2\text{B}_2\text{O}_3$
 $2\text{B} + 3\text{S} \xrightarrow{\Delta} \text{B}_2\text{S}_3$
- Al combines with most of the non-metals on heating.

Reactivity towards halogens

- These elements react with halogens to form trihalides (except TlI_3).
 $2M_{(s)} + 3X_{2(g)} \longrightarrow 2MX_{3(s)}$ ($X = \text{F, Cl, Br, I}$)

- Lewis acid character decreases down the group
 $\text{BX}_3 > \text{AlX}_3 > \text{GaX}_3 > \text{InX}_3$
(Due to increase in size)
- Lewis acid character of boron trihalides
 $\text{BI}_3 > \text{BBr}_3 > \text{BCl}_3 > \text{BF}_3$
(Due to $p\pi$ - $p\pi$ back bonding)


INFOSHOTS


Turning dirty aluminium foil into biofuel catalyst!

Recently, an innovative crystallization method is developed to obtain 100% pure single crystals of aluminium salts from the contaminated foil. This is the starting material for the preparation of alumina catalyst. Usually to produce this type of alumina it would have to come from bauxite ore, which is mined in countries such as West Africa, West Indies and Australia, causing huge environmental damage.

FULLY LOADED & COMPLETELY UPDATED

MTG's BITSAT Power Guide is not only the most exhaustive prep-tool, but also the only book available at present, updated as per the latest BITSAT syllabus for students aspiring for top rank in BITSAT 2018.



₹ 800



Get MTG's BITSAT Power Guide today for a real-world feel of BITSAT. Find out what's different about the BITSAT test, including its pattern of examination and key success factors. Be it with chapterwise MCQs or model test papers, check how good your chances are for glory in BITSAT 2018.

FEATURES

- Covers all 5 subjects - Physics, Chemistry, Mathematics, English & Logical Reasoning
- Chapterwise MCQs in each section for practice
- Past chapterwise BITSAT Qs (2017-2011)
- 10 Model Test Papers with detailed solutions
- Attempt all questions & tests from this book online, for free with #L

Visit www.MTG.in to buy online. Or visit a leading bookseller near you. For more information, email info@mtg.in or call 1800 300 23355 (toll-free).

ANOMALOUS BEHAVIOUR OF BORON

Boron shows anomalous behaviour as compared to the other members of the group due to

- its smallest size and high ionization energy
- highest electronegativity
- absence of *d*-orbitals in its valence shell.



Points of Differences

- B forms only covalent compounds while others form both ionic and covalent compounds.
- Trihalides of B are monomeric while other elements have halogen bridged dimeric structures.
- B has two allotropic forms; crystalline and amorphous. Other members do not show allotropy.
- The melting and boiling points of B are much higher as compared to other members.

- Boron shows a maximum covalency of four in its compounds e.g., BH_4^- while other members show a maximum covalency of six e.g., $[\text{AlF}_6]^{3-}$.

Uses	
Aluminium	
• It forms many useful alloys with Cu, Mn, Mg, Si and Zn.	
• It is used as a conductor for transmission of electricity.	
• It is used for making silvery paints for covering iron or other materials.	
Boron	
• As a moderator in nuclear reactors.	
• As an abrasive and a refractory material.	
• As rocket fuels because of high energy/mass ratio.	
• As a hardener in steel industry.	

IMPORTANT COMPOUNDS OF BORON AND ALUMINIUM

	Preparation	Properties	Structure
Borax ($\text{Na}_2\text{B}_4\text{O}_7 \cdot 10\text{H}_2\text{O}$)	$\text{Ca}_2\text{B}_6\text{O}_{11} + 2\text{Na}_2\text{CO}_3 \xrightarrow{\Delta} \text{Na}_2\text{B}_4\text{O}_7$ Colemanite Borax $+ 2\text{NaBO}_2 + 2\text{CaCO}_3 \downarrow$ $4\text{NaBO}_2 + \text{CO}_2 \longrightarrow \text{Na}_2\text{B}_4\text{O}_7$ $+ \text{Na}_2\text{CO}_3$ $4\text{H}_3\text{BO}_3 + \text{Na}_2\text{CO}_3 \longrightarrow \text{Na}_2\text{B}_4\text{O}_7$ $+ 6\text{H}_2\text{O} + \text{CO}_2$	$\text{Na}_2\text{B}_4\text{O}_7 \cdot 10\text{H}_2\text{O} \xrightarrow{\Delta} \text{Na}_2\text{B}_4\text{O}_7 + 10\text{H}_2\text{O}$ $\text{Na}_2\text{B}_4\text{O}_7 \xrightarrow{\Delta} 2\text{NaBO}_2 + \text{B}_2\text{O}_3$ Transparent glassy bead $\text{Na}_2\text{B}_4\text{O}_7 + 2\text{NaOH} \longrightarrow 4\text{NaBO}_2 + \text{H}_2\text{O}$ $\text{Na}_2\text{B}_4\text{O}_7 + \text{H}_2\text{SO}_4 \longrightarrow \text{Na}_2\text{SO}_4$ $+ \text{H}_2\text{B}_4\text{O}_7$	 <p>$2\text{Na}^+ \left[\text{HO}-\text{B}-\text{O}-\text{B}-\text{O}-\text{B}-\text{O}-\text{B}-\text{OH} \right]$ $\text{Na}_2[\text{B}_4\text{O}_5(\text{OH})_4] \cdot 8\text{H}_2\text{O}$</p>
	Uses : <ul style="list-style-type: none"> • As a water softener and a cleansing agent. • In the laboratory for borax bead test. 		
Orthoboric acid (H_3BO_3)	$\text{Na}_2\text{B}_4\text{O}_7 + 2\text{HCl} + 5\text{H}_2\text{O} \longrightarrow 4\text{H}_3\text{BO}_3 + 2\text{NaCl}$ $\text{Na}_2\text{B}_4\text{O}_7 + \text{H}_2\text{SO}_4 + 5\text{H}_2\text{O} \longrightarrow 4\text{H}_3\text{BO}_3 + \text{Na}_2\text{SO}_4$ $\text{Ca}_2\text{B}_6\text{O}_{11} + 4\text{SO}_2 + 11\text{H}_2\text{O} \longrightarrow 2\text{Ca}(\text{HSO}_3)_2 + 6\text{H}_3\text{BO}_3$ $\text{B}_2\text{H}_6 + 6\text{H}_2\text{O} \longrightarrow 2\text{H}_3\text{BO}_3 + 6\text{H}_2$ $\text{BN} + 3\text{H}_2\text{O} \longrightarrow \text{H}_3\text{BO}_3 + \text{NH}_3$	$\text{H}-\text{OH} + \text{B}(\text{OH})_3 \longrightarrow [\text{B}(\text{OH})_4]^- + \text{H}^+$ $\text{p}K_a = 9.25$ $\text{H}_3\text{BO}_3 \xrightarrow{370\text{ K}} \text{HBO}_2 + \text{H}_2\text{O}$ $\xrightarrow{410\text{ K}} \text{H}_2\text{B}_4\text{O}_7$ $\xrightarrow{-\text{H}_2\text{O}} \text{H}_2\text{B}_4\text{O}_7$ $\downarrow \text{Red Hot}$ $2\text{B}_2\text{O}_3 + \text{H}_2\text{O}$ $\text{B}(\text{OH})_3 + 3\text{C}_2\text{H}_5\text{OH} \xrightarrow{\text{Conc. H}_2\text{SO}_4} \text{B}(\text{OC}_2\text{H}_5)_3 + 3\text{H}_2\text{O}$	

	<p>Uses :</p> <ul style="list-style-type: none"> It is used in the manufacture of heat resistant borosilicate glass. The aqueous solution of boric acid is used as a mild antiseptic especially as eye wash under the name boric lotion. 		
Diborane (B ₂ H ₆)	$2\text{NaBH}_4 + \text{I}_2 \xrightarrow{\text{Diglyme}} \text{B}_2\text{H}_6 + 2\text{NaI} + \text{H}_2$ $2\text{BF}_3 + 6\text{NaH} \xrightarrow{450\text{ K}} \text{B}_2\text{H}_6 + 6\text{NaF}$ $4\text{BF}_3 \cdot \text{Et}_2\text{O} + 3\text{LiAlH}_4 \xrightarrow{\text{Diethyl ether}} 2\text{B}_2\text{H}_6 + 3\text{LiF} + 3\text{AlF}_3 + 4\text{Et}_2\text{O}$	$\text{B}_2\text{H}_6 + 3\text{O}_2 \longrightarrow \text{B}_2\text{O}_3 + 3\text{H}_2\text{O}; \quad \Delta_c H^\circ = -1976 \text{ kJ mol}^{-1}$ $\text{B}_2\text{H}_6 + 6\text{H}_2\text{O} \longrightarrow 2\text{H}_3\text{BO}_3 + 6\text{H}_2$ $\text{B}_2\text{H}_6 + 6\text{CH}_3\text{OH} \longrightarrow 2\text{B}(\text{OCH}_3)_3 + 6\text{H}_2$ $\text{B}_2\text{H}_6 + 2\text{NMe}_3 \longrightarrow 2\text{BH}_3 \cdot \text{NMe}_3$ $3\text{B}_2\text{H}_6 + 6\text{NH}_3 \xrightarrow{\text{Low temp.}} 3[\text{BH}_2(\text{NH}_3)_2]^+ [\text{BH}_4]^- \xrightarrow{473\text{ K}} 2\text{B}_3\text{N}_3\text{H}_6 + 12\text{H}_2$ <p style="text-align: center;">Borazine</p>	<p>Due to the resemblance of three centre two electron bond to a banana, it is also called banana bond.</p>
		<p>Uses :</p> <ul style="list-style-type: none"> For preparing a number of borohydrides reducing agent such as LiBH₄, NaBH₄, etc, in organic reaction. To make bullet-proof vests and light composite material for aircrafts. 	
Aluminium chloride (AlCl ₃)	$2\text{Al} + 3\text{Cl}_2 \longrightarrow 2\text{AlCl}_3$ $2\text{Al} + 6\text{HCl} \longrightarrow 2\text{AlCl}_3 + 3\text{H}_2$ $\text{Al}_2\text{O}_3 + 3\text{C} + 3\text{Cl}_2 \xrightarrow{100^\circ\text{C}} 2\text{AlCl}_3 + 3\text{CO}$ $\text{Al}(\text{OH})_3 + 3\text{HCl} \longrightarrow \text{AlCl}_3 + 3\text{H}_2\text{O}$	$\text{AlCl}_3 + 3\text{H}_2\text{O} \longrightarrow \text{Al}(\text{OH})_3 + 3\text{HCl}$ $\text{AlCl}_3 + 3\text{NH}_4\text{OH} \longrightarrow \text{Al}(\text{OH})_3 \downarrow + 3\text{NH}_4\text{Cl}$ <p style="text-align: center;">Gelatinous ppt. (Insoluble in NH₄OH)</p> $\text{AlCl}_3 + 3\text{NaOH} \longrightarrow \text{Al}(\text{OH})_3 \downarrow + 3\text{NaCl}$ <p style="text-align: center;">(Soluble in excess NaOH)</p> $\text{Al}(\text{OH})_3 + \text{NaOH} \longrightarrow \text{NaAlO}_2 + 2\text{H}_2\text{O}$ <p style="text-align: center;">(Soluble meta-aluminate)</p>	<p style="text-align: center;">(Tetrahedral)</p> <p>AlCl₃ achieves stability by forming dimer.</p>
		<p>Uses :</p> <ul style="list-style-type: none"> Anhydrous AlCl₃ is used as a catalyst in Friedel-Crafts reaction and in cracking of petroleum. It is extensively used in manufacture of dyes, drugs and perfumes. 	

GROUP-14 ELEMENTS (CARBON FAMILY)

INTRODUCTION

Group-14 is transition between metals and non-metals. C is the most versatile element. Organic chemistry is devoted to carbon containing compounds. A new element is added to group-14, *i.e.*, Flerovium (Fl) which is radioactive.

Group-14 contains the following elements with general electronic configuration $ns^2 np^2$.

Element	Electronic configuration
Carbon (6)	[He] $2s^2 2p^2$
Silicon (14)	[Ne] $3s^2 3p^2$
Germanium (32)	[Ar] $3d^{10} 4s^2 4p^2$
Tin (50)	[Kr] $4d^{10} 5s^2 5p^2$
Lead (82)	[Xe] $4f^{14} 5d^{10} 6s^2 6p^2$
Flerovium (114)	[Rn] $5f^{14} 6d^{10} 7s^2 7p^2$

Physical Properties

Covalent and ionic radii

Covalent radii of group-14 elements regularly increases with increase in atomic number. The same trend is observed with ionic radius in their +2 and +4 oxidation states.

Ionisation energy

First *I.E.* decreases from C to Sn but the decrease is not regular, decrease is very sharp from C to Si, but from Si onwards the decrease is very little due to poor shielding effect of 3*d*- and 4*d*- electrons in Ge and Sn. Pb is having a little more *I.E.* than Sn but less than Ge due to lanthanide contraction. Thus, the first *I.E.* of group-14 elements follow the order : C > Si > Ge > Sn < Pb

Electronegativity

C is the most electronegative element in this group. Electronegativity decreases from C to Si and remain constant from Si to Sn and then slightly increases for Pb.

Density

C (diamond) has more density than Si. Except carbon, the density of group-14 element increases with increase in atomic number.

Melting point and boiling point

The m.pt. and b.pt. decreases as we move down the group due to corresponding decrease in the interatomic forces of attraction. However, exceptionally the m.pt. of Sn is lower than that of Pb.

The m.pt. and b.pt. of group-14 elements are higher than the corresponding group-13 elements due to the formation of four covalent bonds with each other resulting in strong binding forces between their atoms in solid as well as in liquid state.

Metallic and non-metallic character

Metallic character increases with increase in atomic number.

C	Si	Ge	Sn	Pb
Non-metal	Metalloid		Metal	

INFOSHOTS

Carbon Nanotube Hybrid Structure!

To introduce and control the temperature coefficient of resistance (TCR) of metal matrix composite, relatively thick and short multi-walled carbon nanotubes (MWCNTs) were introduced in the metal matrix with *in-situ* formation of chromium carbide (Cr₇C₃) at the CNT/copper (Cu) interface. Incompatible properties such as electrical conductivity and TCR can be achieved simultaneously by introducing MWCNTs in the Cu matrix, with control of the interfacial resistivity using the MWCNT/Cr₇C₃-Cu system.

Your favourite MTG Books/Magazines available in JAMMU & KASHMIR at

- Kabli Book Stall - Anantnag Mob: 8803043296, 9906460029
- New Valley Book Depot - Anantnag Mob: 9469183540
- Sharjha Book Depot - Anantnag Mob: 9419040456
- Harnam Dass & Bro'S - Jammu Ph: 0191-2542175, 2574428; Mob: 9419664141
- Sahitya Sangam - Jammu
Ph: 0191-2579593, 2562191, 2579593; Mob: 9419190177
- Shiela Book Centre - Jammu Ph: 0191-2574912; Mob: 9419146803
- Kapoor Sons - Srinagar Ph: 0194-2456458; Mob: 9419425757, 9419069199
- Kashmir Book Depot - Srinagar
Ph: 2450262, 474440, 2475973; Mob: 9906726231, 9419761773
- Shah Book Centre - Srinagar Mob: 9906763627, 9419062444
- Abdullah News Agency - Srinagar
Ph: 0194-2472621, 2435057; Mob: 9419074859
- Highway Book Shop - Srinagar Mob: 9858304786
- Paradise Book House - Srinagar Mob: 9419067856

Visit "MTG IN YOUR CITY" on www.mtg.in to locate nearest book seller OR write to info@mtg.in OR call **0124-6601200** for further assistance.

Chemical Properties

Oxidation states

- C and Si both show an oxidation state of +4 while all other elements of group-14 show two oxidation states of +2 and +4 due to inert pair effect. The stability of +2 oxidation state increases from Ge to Pb *i.e.*, $\overset{2+}{\text{Ge}} < \overset{2+}{\text{Sn}} < \overset{2+}{\text{Pb}}$, while the stability of +4 oxidation state decreases from C to Pb *i.e.*, $\overset{4+}{\text{C}} > \overset{4+}{\text{Si}} > \overset{4+}{\text{Ge}} > \overset{4+}{\text{Sn}} > \overset{4+}{\text{Pb}}$



Oxidising and reducing properties

- Order of the reduction power in +2 oxidation state : $\text{C} > \text{Si} > \text{Ge} > \text{Sn} > \text{Pb}$
- Order of oxidation power in +4 oxidation state : $\text{C} < \text{Si} < \text{Ge} < \text{Sn} < \text{Pb}$

Reactivity towards oxygen

- They form oxides when heated in oxygen but do not react with oxygen at ordinary temperature.
- Group-14 elements form monoxides (MO) and dioxides (MO₂). SiO exists only at high temperature.
- Acidic nature of oxides :

CO_2	SiO_2	GeO_2	SnO_2	PbO_2	(MO ₂ type)
Acidic			Amphoteric		
CO	GeO	SnO	PbO	(MO type)	
Neutral	Acidic	Amphoteric			
- Pb can form Pb₃O₄ (trilead tetraoxide), known as red lead or sindhur, on heating PbO (litharge) with excess of air at 673 K.

$$6\text{PbO} + \text{O}_2 \xrightarrow{673\text{K}} 2\text{Pb}_3\text{O}_4$$

ANOMALOUS BEHAVIOUR OF CARBON

C differs from the rest of the members of group-14 because of

- its small atomic size
- its high electronegativity
- its property of catenation
- absence of *d*-orbitals in its valence shell.

Points of Differences

- The m.pt. and b.pt., *I.E.* and electronegativity of C is very high as compared to the other members of its family.

Reactivity towards water

- C, Si and Ge are not affected by water. Sn decomposes steam.

$$\text{Sn} + 2\text{H}_2\text{O} \xrightarrow{\Delta} \text{SnO}_2 + 2\text{H}_2$$
- Pb is unaffected by water on its surface probably because of a protective oxide film formation.
- Pb dissolved slowly in water containing dissolved oxygen. This is called plumbosolvency.

$$2\text{Pb} + 2\text{H}_2\text{O} + \text{O}_2 \longrightarrow 2\text{Pb}(\text{OH})_2$$

Reactivity towards halogen

- These elements can form halides of formula MX₂ and MX₄ (X = F, Cl, Br, I).
- Except carbon, all other members react directly with halogen.
- Stability of dihalides increases down the group.
- Except CCl₄, other tetrachlorides are easily hydrolysed by water because the central atom can accommodate the lone pair of electrons from oxygen atom of water molecule in vacant *d*-orbitals.

Reducing character

- Carbon is a powerful reducing agent, it can reduce oxides of various elements *e.g.*,

$$\text{SnO}_2 + 2\text{C} \xrightarrow{\Delta} \text{Sn} + 2\text{CO}$$

$$\text{CO}_2 + \text{C} \xrightarrow{\Delta} 2\text{CO}$$

$$\text{Fe}_2\text{O}_3 + 3\text{CO} \xrightarrow{250^\circ-700^\circ\text{C}} 2\text{Fe} + 3\text{CO}_2$$
- Si is also used as deoxidizer in the manufacture of steel in the form of ferrosilicon.
- Sn also acts as a good reducing agent.

- Carbon in its diamond form, is one of the hardest known element.
- It has maximum tendency to show catenation.
- Carbon has strong tendency to form *pπ-pπ* multiple bonds either with itself or with other elements like N, O, S, etc. Other members of the family form *pπ-dπ* bonds and that to a lesser extent.
- CO₂ is a gas while the dioxides of all other members are solids.
- Carbon shows a maximum covalency of four while other members of the family may expand their covalency beyond 4 *e.g.*, [SiCl₆]²⁻.

ALLOTROPES OF CARBON



- **Diamond** : In diamond, C-atom is sp^3 hybridised and has three dimensional network structure. Since, no free electron is available, it is a bad conductor of electricity.
- **Graphite**: C-atom is sp^2 hybridised and each carbon is directly bound to three other C-atoms. Thus, graphite has a delocalized π -electron cloud, which is responsible for its high electrical conductivity. It possesses layer structure.
- **Fullerenes** : It consists 12 pentagons and 20 hexagons folded into a sphere, so that it looks like a soccer ball, called 'bucky ball'.

All the carbon atoms are equal and they undergo sp^2 hybridisation.

USES OF CARBON

- Graphite is used as a lubricant, electrodes for dry cell, moderator for fast moving neutrons in nuclear reactor.
- Graphite is used in lead pencils.
- Diamond is used for cutting glass and other cutting and drilling tools. It is also used for making precious gems and jewellery.
- Being highly porous, activated charcoal is used in adsorbing poisonous gases, also used in water filters to remove organic contaminants and in air conditioning system to control odour.

IMPORTANT COMPOUNDS OF CARBON AND SILICON

	Preparation	Properties	Structure
Carbon monoxide (CO)	$2C + O_2 \xrightarrow{\Delta} 2CO$ $HCOOH \xrightarrow[Conc. H_2SO_4]{373 K} H_2O + CO$ $C + H_2O \xrightarrow{473 - 1273 K} \underbrace{CO + H_2}_{\text{Water gas}}$ $2C + O_2 + 4N_2 \xrightarrow{473 - 1273 K} \underbrace{2CO + 4N_2}_{\text{Producer gas}}$	$2CO + O_2 \longrightarrow 2CO_2$ $3CO + Fe_2O_3 \xrightarrow{\Delta} 2Fe + 3CO_2$ $CO + ZnO \xrightarrow{\Delta} Zn + CO_2$ $4CO + Ni \xrightarrow{80^\circ C} [Ni(CO)_4]$ $5CO + Fe \xrightarrow{180^\circ C} [Fe(CO)_5]$ CO is highly poisonous gas as it forms a complex with haemoglobin (Hb) which is 300 times more stable than O_2 -Hb complex. Thus, it prevents haemoglobin to carry oxygen from the lungs to other parts of the body.	$:C \equiv \ddot{O}:$ \longleftrightarrow $:\ddot{O} \equiv C:$ or $:C \equiv \ddot{O}:$ (Linear structure with sp hybridisation)
Uses :	<ul style="list-style-type: none"> • As a fuel in the form of water gas, producer gas etc. • In the extraction of iron as reducing agent. 		
Carbon dioxide (CO₂)	$C + O_2 \xrightarrow{\Delta} CO_2$ $CH_4 + 2O_2 \xrightarrow{\Delta} CO_2 + 2H_2O$ $CaCO_3 + 2HCl \longrightarrow CaCl_2 + CO_2 + H_2O$ $C_6H_{12}O_6 \xrightarrow{Zymase} 2C_2H_5OH + 2CO_2$	$CO_2 + 2Mg \longrightarrow 2MgO + C$ $CO_2 + H_2O \rightleftharpoons H_2CO_3$ $CO_2 + K_2CO_3 + H_2O \longrightarrow 2KHCO_3$ $CO_2 + Ca(OH)_2 \longrightarrow CaCO_3 \downarrow + H_2O$ $CO_2 + CaCO_3 + H_2O \longrightarrow Ca(HCO_3)_2$ $CO_2 + Zn \longrightarrow ZnO + CO$ $6CO_2 + 6H_2O \xrightarrow{hv} C_6H_{12}O_6 + 6O_2$	
Uses :	<ul style="list-style-type: none"> • As carbogen [mixture of O_2 ($\approx 95\%$) + CO_2 ($\approx 5-10\%$)] in artificial respiration especially for pneumonia patients and victims of CO poisoning. • As a fire extinguisher. 		
Silicon dioxide (SiO₂, Silica)	It is a covalent, three dimensional network solid, almost non-reactive due to high Si—O bond enthalpy. However, it is attacked by HF and NaOH. $SiO_2 + 2NaOH \longrightarrow Na_2SiO_3 + H_2O$ $SiO_2 + 4HF \longrightarrow SiF_4 + 2H_2O$		
Uses :	<ul style="list-style-type: none"> • Quartz (crystallographic form of silica) is extensively used as a piezoelectric material. • Silica gel is used as a drying agent and as a supporter for chromatographic materials and catalysts. • Kieselguhr, an amorphous form of silica, is used in filtration plants. 		


<p>Silicones</p> $\left[\text{O} - \begin{array}{c} \\ \text{R} \\ \\ \text{Si} \\ \\ \text{R} \end{array} - \text{O} \right]_n$ <p>where R = alkyl or phenyl group</p>	<ul style="list-style-type: none"> Synthetic organosilicon polymers containing repeated $R_2\text{SiO}$ units held by Si—O—Si linkage. They are water repelling due to non-polar alkyl groups. $2\text{CH}_3\text{Cl} + \text{Si} \xrightarrow[570\text{ K}]{\text{Cu powder}} (\text{CH}_3)_2\text{SiCl}_2 \xrightarrow[-2\text{HCl}]{+2\text{H}_2\text{O}} (\text{CH}_3)_2\text{Si}(\text{OH})_2$ $n \text{HO} - \begin{array}{c} \\ \text{CH}_3 \\ \\ \text{Si} \\ \\ \text{CH}_3 \end{array} - \text{OH} \xrightarrow[\text{Polymerisation}]{-n\text{H}_2\text{O}} \left[\text{O} - \begin{array}{c} \\ \text{CH}_3 \\ \\ \text{Si} \\ \\ \text{CH}_3 \end{array} - \text{O} - \begin{array}{c} \\ \text{CH}_3 \\ \\ \text{Si} \\ \\ \text{CH}_3 \end{array} \right]_n$ <p style="text-align: center;">Silicone straight chain</p> <p>The chain length of polymer can be controlled by adding $(\text{CH}_3)_3\text{SiCl}$ which blocks the ends as shown below :</p> $n \text{HO} - \begin{array}{c} \\ \text{CH}_3 \\ \\ \text{Si} \\ \\ \text{CH}_3 \end{array} - \text{OH} + \text{HO} - \begin{array}{c} \\ \text{CH}_3 \\ \\ \text{Si} \\ \\ \text{CH}_3 \end{array} - \text{CH}_3 \xrightarrow[\text{Polymerisation}]{-n\text{H}_2\text{O}} \left[\text{O} - \begin{array}{c} \\ \text{CH}_3 \\ \\ \text{Si} \\ \\ \text{CH}_3 \end{array} - \text{O} - \begin{array}{c} \\ \text{CH}_3 \\ \\ \text{Si} \\ \\ \text{CH}_3 \end{array} - \text{CH}_3 \right]_n$ <p style="text-align: center;">Silicone</p>	$\left[\begin{array}{c} \text{R} \\ \\ \text{Si} \\ \\ \text{R} \end{array} - \text{O} - \begin{array}{c} \text{R} \\ \\ \text{Si} \\ \\ \text{R} \end{array} \right]_n$
---	---	---

Uses :

- They are used as sealant, greases, electrical insulators and for water proofing of fabrics. Being biocompatible they are also used in surgical and cosmetic implants.

Silicates

Silicates are discrete SiO_4^{4-} tetrahedra or a number of such units joined together by sharing of oxygen.




ZEOLITES

- Zeolites are hydrated 3-dimensional aluminosilicates which are formed by replacing some of the silicon atoms by Al^{3+} ions. Their general formula is $M_{x/n}[(\text{AlO}_2)_x(\text{SiO}_2)_y] \cdot m\text{H}_2\text{O}$.
- Zeolites are used as catalysts in petrochemical industries. They are also used as molecular sieves and softening of water by ion-exchange method.

SPEED PRACTICE

- Which gas is evolved when PbO_2 is treated with conc. HNO_3 ?
(a) NO_2 (b) O_2 (c) N_2 (d) N_2O
- In group-13, electronegativity first decreases from B to Al and then increases marginally down the group. This is because of
(a) non-metallic nature of B
(b) discrepancies in atomic size of elements
(c) ability of B and Al to form $p\pi - p\pi$ multiple bonds
(d) irregular trend in electronegativity throughout the periodic table.
- Which of the following statements is incorrect?
(a) The carbon dioxide molecule behaves as a non-polar molecule even though two of its resonating structures, $\overset{+}{\text{O}}=\text{C}-\overset{-}{\text{O}}$ and $\overset{+}{\text{O}}\equiv\text{C}-\overset{-}{\text{O}}$, are polar.
(b) Carbon dioxide is the anhydride of the unstable dibasic acid $\text{O}=\text{C}(\text{OH})_2$.
(c) The carbon dioxide molecule is linear because the carbon atom utilises its sp -orbitals to form σ -bonds.
(d) The carbon atom is sp^2 -hybridised in the CO_2 molecule as well as in the molecule of its hydrate H_2CO_3 .
- TlI_3 , a soluble ionic compound, in aqueous medium will give
(a) Tl^{3+} and I^- ions (b) Tl^+ , I^- and I_2
(c) Tl^{3+} and I_3^- ions (d) Tl^+ and I_3^- ions.
- Among the following, the correct statement(s) is(are)
(a) $\text{Al}(\text{CH}_3)_3$ has the three-centre two-electron bonds in its dimeric structure
(b) BH_3 has the three-centre two-electron bonds in its dimeric structure
(c) the Lewis acidity of BCl_3 is greater than that of AlCl_3
(d) AlCl_3 has the three-centre two-electron bonds in its dimeric structure. (JEE Advanced 2017)
- Which of the following halides is least stable and has doubtful existence?
(a) Cl_4 (b) GeI_4 (c) SnI_4 (d) PbI_4
- Incomplete combustion of petrol or diesel oil in automobile engines can be best detected by testing the fuel gases for the presence of
(a) carbon monoxide and water vapour
(b) carbon monoxide
(c) nitrogen dioxide
(d) sulphur dioxide.
- In BX_3 , B - X distance is shorter than what is expected theoretically because ($X = \text{F}, \text{Cl}, \text{Br}, \text{I}$)
(a) of sp^3 hybridisation of B
(b) B - X has a double bond character due to back-bonding
(c) dimerisation takes place in BX_3
(d) of large size of X, B - X distance decreases.
- An inorganic compound (A), composed of two most occurring elements in the earth's crust, have polymeric tetrahedral network structure. With carbon, compound (A) produces a poisonous gas (B) which is a stable diatomic molecule. Compounds (A) and (B) can be
(a) $\text{SiO}_2, \text{CO}_2$ (b) SiO_2, CO
(c) SiC, CO (d) SiO_2, N_2
- The increasing order of atomic radii of the following group 13 elements is
(a) $\text{Al} < \text{Ga} < \text{In} < \text{Tl}$
(b) $\text{Ga} < \text{Al} < \text{In} < \text{Tl}$
(c) $\text{Al} < \text{In} < \text{Ga} < \text{Tl}$
(d) $\text{Al} < \text{Ga} < \text{Tl} < \text{In}$ (JEE Advanced 2016)
- In carbon-60, all carbon atoms are
(a) sp^2 -hybridised with a truncated icosahedron shape
(b) sp^3 -hybridised with a square antiprism shape
(c) sp^2 -hybridised with a diamond-like shape
(d) sp^2 -hybridised with a graphite-like shape.
- AlF_3 is soluble in HF only in presence of KF. It is due to the formation of
(a) $\text{K}_3[\text{AlF}_3\text{H}_3]$ (b) $\text{K}_3[\text{AlF}_6]$
(c) AlH_3 (d) $\text{K}[\text{AlF}_3\text{H}]$ (NEET Phase-II 2016)

13. An aqueous solution of borax is
 (a) neutral (b) amphoteric
 (c) basic (d) acidic.
14. Which of the following structures correctly represents the boron trifluoride molecule?



15. Consider the following standard electrode potentials (E° in volts) in aqueous solution,

Element	M^{3+} / M	M^+ / M
Al	-1.66	+0.55
Tl	+1.26	-0.34

Based on these data, which of the following statements is correct?

- (a) Tl^+ is more stable than Al^+ .
 (b) Tl^{3+} is more stable than Al^{3+} .
 (c) Al^+ is more stable than Al^{3+} .
 (d) Tl^+ is more stable than Al^{3+} .
- (JEE Main online 2017)
16. Even though thermodynamically favourable, the conversion of diamond into graphite does not occur normally since
 (a) the activation energy of the process is high.
 (b) the activation energy of the process is low.
 (c) the change of entropy is zero.
 (d) the change of enthalpy is zero.
17. $AlCl_3$ achieves stability by forming a dimer. In trivalent state the compound is hydrolysed in water. The product formed is
 (a) $AlCl_3 \cdot 4H_2O$ (b) $AlCl_3 \cdot 6H_2O$
 (c) $AlCl_3 \cdot 2H_2O$ (d) $Al_2O_3 + HCl$

18. Select incorrect statement.
 (a) Cyanamide ion, $(CN_2)^{2-}$ is isoelectronic with CO_2 and has the same linear structure.
 (b) Mg_2C_3 reacts with water to form propyne.

- (c) CaC_2 has C_2^{2-} and it contains one sigma and two pi bonds.
 (d) Al_4C_3 is an example of methanide carbide and it contains C_4^{4-} .

19. Which of the following statements is incorrect?
 (a) $B(OH)_3$ partially reacts with water to form H_3O^+ and $[B(OH)_4]^-$, and behaves like a weak acid.
 (b) $B(OH)_3$ behaves like a strong monobasic acid in the presence of sugars, and this acid can be titrated against NaOH solution using phenolphthalein as an indicator.
 (c) $B(OH)_3$ does not donate a proton and hence does not form any salt with NaOH.
 (d) $B(OH)_3$ reacts with NaOH, forming $Na[B(OH)_4]$.
20. CCl_4 does not undergo hydrolysis but $SiCl_4$ is readily hydrolysed because
 (a) carbon cannot expand its octet but silicon can expand
 (b) electronegativity of carbon is higher than silicon
 (c) I.E. of carbon is higher than that of silicon
 (d) carbon forms double and triple bonds but not silicon.

21. Boric acid is an acid because its molecule
 (a) contains replaceable H^+ ion
 (b) gives up a proton
 (c) accepts OH^- from water releasing proton
 (d) combines with proton from water molecule.
- (NEET Phase-II 2016)

22. Amongst the halides
 (1) BCl_3 (2) $AlCl_3$ (3) $GaCl_3$ (4) $InCl_3$
 the order of decreasing Lewis acid character is
 (a) 1, 2, 3, 4 (b) 4, 3, 2, 1
 (c) 3, 4, 2, 1 (d) 2, 3, 4, 1
23. $(COOH)_2 \xrightarrow{\text{Conc. } H_2SO_4} A + B + H_2O$, concentration of A can be analysed by using iodine pentoxide, then incorrect statement about 'B' is
 (a) solid B sublimates at 194.65 K
 (b) B reacts with ammonia to form urea
 (c) supercritical B is used as solvent
 (d) B is linear and neutral oxide.
24. What happens when a mixture of cobalt oxide and borax is heated in a flame on a loop of platinum wire?
 (a) A transparent white bead is formed.
 (b) A bright pink coloured $NaBO_2$ bead is formed.
 (c) A blue coloured $Co(BO_2)_2$ bead is formed.
 (d) A red coloured $Co(BO_2)_2$ bead is formed.

25. The pair in which phosphorus atoms have a formal oxidation state of +3 is
 (a) orthophosphorous and pyrophosphorous acids
 (b) pyrophosphorous and hypophosphoric acids
 (c) orthophosphorous and hypophosphoric acids
 (d) pyrophosphorous and pyrophosphoric acids.

(JEE Main 2016)

26. X reacts with aqueous NaOH solution to form Y and H₂. Aqueous solution of Y on heating to 323-333 K and on passing CO₂ into it, Na₂CO₃ and Z are formed. When Z is heated to 1200 °C, Al₂O₃ is formed. X, Y and Z respectively are
 (a) Al, AlCl₃, NaAlO₂ (b) Zn, Na₂ZnO₂, Al(OH)₃
 (c) Al, Al(OH)₃, AlCl₃ (d) Al, NaAlO₂, Al(OH)₃

27. Ge(II) compounds are powerful reducing agents whereas Pb(IV) compounds are strong oxidants. This can be due to

- (a) Pb is more electropositive than Ge
 (b) ionisation potential of lead is less than that of Ge
 (c) ionic radii of Pb²⁺ and Pb⁴⁺ are larger than those of Ge²⁺ and Ge⁴⁺
 (d) more pronounced inert pair effect in Pb than in Ge.



28. CO₂ and N₂ are non-supporters of combustion. However, for putting out fires CO₂ is preferred over N₂ because CO₂

- (a) does not burn
 (b) forms non-combustible products with burning substances
 (c) is denser than nitrogen
 (d) is a more reactive gas.

29. Alum helps in purifying water by

- (a) forming Si complex with clay particles
 (b) sulphate part which combines with dirt and removes it
 (c) coagulating the mud particles
 (d) making mud water soluble.


30. $\text{NaBH}_4 + \text{I}_2 \xrightarrow[\text{solution}]{\text{in diglyme}}$ X + Y + Z



X, Y, Z, P, Q and R in the reactions are


X	Y	Z	P	Q	R
(a) Na ₂ B ₄ O ₇	NaI	HI	HF	LiF	AlF ₃
(b) B ₂ H ₆	NaI	H ₂	NaF	LiF	AlF ₃
(c) B ₂ H ₆	BH ₃	NaI	B ₃ N ₃ H ₆	Al ₂ F ₆	AlF ₃
(d) BH ₃	B ₂ H ₆	H ₂	B ₃ N ₃ H ₆	LiF	AlF ₃

SOLUTIONS




(a) : Name	Formula	Oxidation state
Orthophosphorous acid	H ₃ PO ₃	+3
Pyrophosphorous acid	H ₄ P ₂ O ₅	+3
Hypophosphoric acid	H ₄ P ₂ O ₆	+4
Pyrophosphoric acid	H ₄ P ₂ O ₇	+5


3. (d): Carbon atom is sp-hybridised in CO₂ molecule and sp²-hybridised in H₂CO₃.



5. (a, b, c) : (a) Al₂(CH₃)₆ has the three centre-two electron bonds.




- (b) Dimer of BH₃ has 3-center 2-electron bond.



- (c) The size of element increases down the group, thus, Lewis acid character of BCl₃ is greater than that of AlCl₃.


- (d) AlCl₃ has dative bond in its dimeric structure, not 3-center 2-electron bond.



6. (d): It is because of inert pair effect.

7. (b): Incomplete combustion of petrol or diesel gives carbon monoxide gas.

8. (b): Due to back-bonding, B - X bond in BX₃ has a double bond character which, results in shorter B - X distance.



9. (b): The most abundant elements in the earth's crust are oxygen and silicon. They form SiO₂ which has polymeric tetrahedral network structure. This on heating with carbon forms SiC and CO.

BEST TOOLS FOR SUCCESS IN

JEE Main



10 Very Similar Practice Test Papers

16 Years JEE MAIN 2017-2015(Offline & Online)-2013 & AIEEE (2012-2002)



Available at all leading book shops throughout India.
For more information or for help in placing your order:
Call 0124-6601200 or email: info@mtg.in


Visit
www.mtg.in
for latest offers
and to buy
online!

10. (b): The increasing order of atomic radii of group 13 elements is $\text{Ga} < \text{Al} < \text{In} < \text{Tl}$.

Atomic radius of Ga is slightly lower than that of Al due to the presence of d -electrons in Ga which do not shield the nucleus effectively.

11. (a): In carbon-60, all carbon atoms are sp^2 -hybridised with a truncated icosahedron shape.

12. (b): AlF_3 is insoluble in anhydrous HF because the F^- ions are not available in hydrogen bonded HF but, in the presence of little amount of KF it forms soluble complex, $\text{K}_3[\text{AlF}_6]$.



13. (c): An aqueous solution of borax is basic in nature due to the hydrolysis of $\text{B}_4\text{O}_7^{2-}$ ions.


14. (a): BF_3 molecule involves extensive back bonding from fluorine to boron.

15. (a): Al^{3+} is more stable than Tl^{3+} because of negative $E_{\text{Al}^{3+}/\text{Al}}^\circ$ value.

Tl^+ is more stable than Al^+ because of negative $E_{\text{Tl}^+/\text{Tl}}^\circ$ value.



16. (a)

17. (b): AlCl_3 in aqueous solution forms $\text{AlCl}_3 \cdot 6\text{H}_2\text{O}$.



18. (d)

19. (c): Boric acid being acidic in nature forms salt (sodium metaborate) with NaOH.




20. (a)


21. (c): Boric acid behaves as a Lewis acid, by accepting a pair of electrons from OH^- ion of water thereby releasing a proton.

22. (a): Lewis acid strength of group-13 halides follow the order: $\text{BCl}_3 > \text{AlCl}_3 > \text{GaCl}_3 > \text{InCl}_3$.

23. (d): CO_2 is an acidic oxide.



24. (c): $\text{Na}_2\text{B}_4\text{O}_7 \cdot 10\text{H}_2\text{O} \xrightarrow{\Delta} \text{Na}_2\text{B}_4\text{O}_7 \xrightarrow{\Delta} 2\text{NaBO}_2 + \text{B}_2\text{O}_3$




25. (b)

26. (d): $2\text{Al} + 2\text{NaOH} + 2\text{H}_2\text{O} \longrightarrow 2\text{NaAlO}_2 + 3\text{H}_2$


(X)

(Y)



(Y)

(Z)






(Z)

27. (d): Ge^{4+} is more stable than Ge^{2+} as a result of which Ge^{2+} has a tendency to be oxidised to Ge^{4+} , so Ge^{2+} compounds act as powerful reducing agents. But Pb^{2+} is more stable than Pb^{4+} because of pronounced inert pair effect as a result of which Pb^{4+} has a tendency to get reduced to Pb^{2+} so Pb^{4+} compounds act as strong oxidising agents.

28. (c): CO_2 is denser than N_2 and air thus surrounds the igniting material more effectively.

29. (c): Alums help in purifying water by coagulating the negatively charged mud particles by Al^{3+} ions provided by it.



◆◆


NEET
ONLINE TEST SERIES

Practice Part Syllabus/ Full Syllabus
40 Mock Test

Now on your android & IOS Smart phones
with the same login of web portal.

Log on to <http://pcmbtoday.com>

EXAMINER'S MIND CLASS XI



The questions given in this column have been prepared strictly on the basis of NCERT Chemistry for Class XI. This year JEE (Main & Advanced)/NEET/AIIMS have drawn their papers heavily from NCERT books.

Section - I	Q. 1 to 10 Only One Option Correct Type MCQs.
Section - II	Q. 11 to 13 More than One Options Correct Type MCQs.
Section - III	Q. 14 to 17 Paragraph Type MCQs having Only One Option Correct.
Section - IV	Q. 18 & 19 Matching List Type MCQs having Only One Option Correct.
Section - V	Q. 20 to 22 Assertion Reason Type MCQs having Only One Option Correct. Mark the correct choice as : (a) If both assertion and reason are true and reason is the correct explanation of assertion. (b) If both assertion and reason are true but reason is not the correct explanation of assertion. (c) If assertion is true but reason is false. (d) If both assertion and reason are false.
Section - VI	Q. 23 to 25 Integer Value Correct Type Questions having Single Digit Integer Answer, ranging from 0 to 9 (both inclusive).

EQUILIBRIUM


SECTION - I

Only One Option Correct Type


- In which of the following reactions, the equilibrium remains unaffected on addition of small amount of argon at constant volume?
 - $\text{H}_{2(g)} + \text{I}_{2(g)} \rightleftharpoons 2\text{HI}_{(g)}$
 - $\text{PCl}_{5(g)} \rightleftharpoons \text{PCl}_{3(g)} + \text{Cl}_{2(g)}$
 - $\text{N}_{2(g)} + 3\text{H}_{2(g)} \rightleftharpoons 2\text{NH}_{3(g)}$
 - All of these
- a moles of PCl_5 are heated in a closed container to equilibrate $\text{PCl}_{5(g)} \rightleftharpoons \text{PCl}_{3(g)} + \text{Cl}_{2(g)}$ at a pressure of P atm. If x moles of PCl_5 dissociate at equilibrium, then
 - $\frac{x}{a} = \frac{K_p}{K_p + P}$
 - $\frac{x}{a} = \left(\frac{K_p + P}{K_p} \right)^{1/2}$
 - $\frac{x}{a} = \left(\frac{K_p}{P} \right)^{1/2}$
 - $\frac{x}{a} = \left(\frac{K_p}{K_p + P} \right)^{1/2}$
- At 350 K, K_p for the reaction given below is $3.0 \times 10^{10} \text{ bar}^{-1}$ at equilibrium. What will be the value of K_c at this temperature?

$$2\text{N}_{2(g)} + \text{O}_{2(g)} \rightleftharpoons 2\text{N}_2\text{O}_{(g)}$$
 - $7.4 \times 10^{11} \text{ L mol}^{-1}$
 - $8.715 \times 10^{10} \text{ L mol}^{-1}$
 - 0.08 L mol^{-1}
 - $8.715 \times 10^{11} \text{ L mol}^{-1}$
- What is the correct sequence, to complete the paragraph? According to Le-Chatelier's principle, if pressure is increased,
 - if pressure is decreased, the system will try to increase the pressure
 - the system will try to reduce the pressure
 - by favouring that reaction which is accompanied by a decrease in the number of moles
 - by favouring that reaction which is accompanied by an increase in the number of moles.
 - 2 3 1 4
 - 1 2 3 4
 - 1 4 3 2
 - 1 3 2 4
- Dissociation constant values of few acids are given below. Arrange them in order of increasing acidic strength. (K_a values : $\text{H}_2\text{SO}_3 = 1.3 \times 10^{-2}$, $\text{HNO}_2 = 4 \times 10^{-4}$, $\text{CH}_3\text{COOH} = 1.8 \times 10^{-5}$, $\text{HCN} = 4 \times 10^{-10}$)
 - $\text{HCN} < \text{CH}_3\text{COOH} < \text{HNO}_2 < \text{H}_2\text{SO}_3$
 - $\text{CH}_3\text{COOH} < \text{HNO}_2 < \text{HCN} < \text{H}_2\text{SO}_3$
 - $\text{CH}_3\text{COOH} < \text{HCN} < \text{H}_2\text{SO}_3 < \text{HNO}_2$
 - $\text{HNO}_2 < \text{H}_2\text{SO}_3 < \text{CH}_3\text{COOH} < \text{HCN}$


6. Solid $\text{Ba}(\text{NO}_3)_2$ is gradually dissolved in $1.0 \times 10^{-4} \text{ M}$ Na_2CO_3 solution. At what concentration of Ba^{2+} the precipitate will begin to form? (K_{sp} for $\text{BaCO}_3 = 5.1 \times 10^{-9}$)
- (a) $4.1 \times 10^{-5} \text{ M}$ (b) $5.1 \times 10^{-5} \text{ M}$
 (c) $8.1 \times 10^{-8} \text{ M}$ (d) $8.1 \times 10^{-7} \text{ M}$
7. Which of the following reflects that the reaction is independent of temperature?



8. For the equilibrium,



equilibrium constant is found to be 1.732 at 298 K. Now if in a vessel at 298 K, a mixture of these two gases is taken and the point p is represented in the given figure, predict what will happen?



- (a) Immediately equilibrium will set-up.
 (b) The given reaction will go in the forward direction till it attains equilibrium.
 (c) The given reaction will go in the backward direction till it attains equilibrium.
 (d) Cannot be predicted.
9. Calculate the degree of dissociation of PCl_5 , when the density at 230°C is 70.
- (a) 97.8% (b) 48.9%
 (c) 4.89% (d) 24.45%
10. The correct order of increasing $[\text{H}_3\text{O}^+]$ in the following aqueous solutions is
- (a) $0.01 \text{ M H}_2\text{S} < 0.01 \text{ M H}_2\text{SO}_4 < 0.01 \text{ M NaCl} < 0.01 \text{ M NaNO}_2$
 (b) $0.01 \text{ M NaCl} < 0.01 \text{ M NaNO}_2 < 0.01 \text{ M H}_2\text{S} < 0.01 \text{ M H}_2\text{SO}_4$


- (c) $0.01 \text{ M NaNO}_2 < 0.01 \text{ M NaCl} < 0.01 \text{ M H}_2\text{S} < 0.01 \text{ M H}_2\text{SO}_4$
 (d) $0.01 \text{ M H}_2\text{S} < 0.01 \text{ M NaNO}_2 < 0.01 \text{ M NaCl} < 0.01 \text{ M H}_2\text{SO}_4$

SECTION - II

More than One Options Correct Type

11. For the reaction, $\text{N}_2\text{O}_4(\text{g}) \rightleftharpoons 2\text{NO}_2(\text{g})$, the value of K is 50 at 400 K and 1700 at 500 K. Which of the following options are correct?
- (a) The reaction is endothermic.
 (b) The reaction is exothermic.
 (c) If $\text{NO}_2(\text{g})$ and $\text{N}_2\text{O}_4(\text{g})$ are mixed at 400 K at partial pressures 20 bar and 2 bar respectively, more $\text{N}_2\text{O}_4(\text{g})$ will be formed.
 (d) The entropy of the system remains constant.
12. The thermal decomposition of $\text{CaCO}_3(\text{s})$ is studied under different conditions,
 $\text{CaCO}_3(\text{s}) \rightleftharpoons \text{CaO}(\text{s}) + \text{CO}_2(\text{g})$
 For this equilibrium, the correct statements are
- (a) ΔH is dependent on T
 (b) K is independent of the initial amount of CaCO_3
 (c) K is dependent on the pressure of CO_2 at a given T
 (d) ΔH is independent of the catalyst, if any.

13. The equilibrium constants of the reactions,



are K_1 and K_2 respectively. The relationship between K_1 and K_2 is

- (a) $K_1 = K_2$ (b) $K_2 = \sqrt{K_1}$
 (c) $K_1 = K_2^2$ (d) $K_1 = \sqrt{K_2}$

SECTION - III

Paragraph Type

Paragraph for Questions 14 and 15

A number of acid-base concepts have been put forward from time to time. Arrhenius laid emphasis on H^+ ions and OH^- ions, whereas Bronsted and Lowry considered acids and bases as proton donors and acceptors and also suggested the concept of conjugate acid-base pairs. Lewis concept was the electronic concept. The first two concepts could also explain the strengths of acids and bases. Arrhenius could explain in terms of dissociation constants of the acids and bases whereas Bronsted-Lowry could do it by suggesting that a strong acid has a weak conjugate base and vice-versa.

14. In the reaction $I_2 + I^- \longrightarrow I_3^-$, the Lewis base is
 (a) I_2 (b) I^- (c) I_3^- (d) none.
15. The successive dissociation constants K_{a_1} , K_{a_2} and K_{a_3} of orthophosphorous acid (H_3PO_3) could be
 (a) 7×10^{-7} , 1.6×10^{-2} , 0
 (b) 1.6×10^{-2} , 7×10^{-7} , 0
 (c) 1.6×10^{-2} , 7×10^{-7} , 1×10^{-12}
 (d) 1×10^{-12} , 7×10^{-7} , 1.6×10^{-2}

Paragraph for Questions 16 and 17

Two solids X and Y dissociate into gaseous products at a certain temperature as follows :

- (i) $X_{(s)} \rightleftharpoons A_{(g)} + C_{(g)}$ and
 (ii) $Y_{(s)} \rightleftharpoons B_{(g)} + C_{(g)}$

At a given temperature, total pressure over solid 'X' is 40 mm of Hg and total pressure over solid 'Y' is 60 mm of Hg. Now, answer the following questions :

16. Ratio of K_p for reaction (i) to that of reaction (ii), is
 (a) 4 : 9 (b) 2 : 3 (c) 3 : 2 (d) 2 : 1
17. The ratio of moles of A and B in the vapour state over a mixture of solids X and Y, is
 (a) 2 : 3 (b) 2 : 5 (c) 4 : 9 (d) 1 : 1

SECTION - IV

Matching List Type

18. Match the List I with List II and select the correct answer using the codes given below the lists :

List I		List II	
(P) $2NO \rightleftharpoons N_2 + O_2$	(i) $K_p = K_c/RT$		
(Q) $N_2 + 3H_2 \rightleftharpoons 2NH_3$	(ii) $K_p = K_cRT$		
(R) $PCl_3 + Cl_2 \rightleftharpoons PCl_5$	(iii) $K_p = K_c$		
(S) $2O_3 \rightleftharpoons 3O_2$	(iv) $K_p = K_c(RT)^{-2}$		

P	Q	R	S
(a) (iii)	(iv)	(i)	(ii)
(b) (i)	(ii)	(iii)	(iv)
(c) (iv)	(iii)	(i)	(ii)
(d) (ii)	(i)	(iv)	(iii)

MPP-8 CLASS XII ANSWER KEY

1. (a) 2. (d) 3. (d) 4. (b) 5. (b)
 6. (b) 7. (a) 8. (c) 9. (b) 10. (a)
 11. (c) 12. (a) 13. (c) 14. (d) 15. (b)
 16. (a) 17. (b) 18. (d) 19. (b) 20. (b,d)
 21. (a,b) 22. (a,c) 23. (c,d) 24. (1) 25. (3)
 26. (1) 27. (d) 28. (b) 29. (d) 30. (c)

19. Match the List I with List II and select the correct answer using the codes given below the lists :

List I		List II	
(P) Solubility of Hg_2Cl_2 in 0.1 M NaCl solution.	(i) $10 K_{sp}$		
(Q) Solubility of PbI_2 in 0.01 M KI solution.	(ii) $100 K_{sp}$		
(R) Solubility of Ag_2CrO_4 in 0.25 M K_2CrO_4 solution.	(iii) $10000 K_{sp}$		
(S) Solubility of calcium oxalate in 0.1 M oxalic acid solution.	(iv) $\sqrt{K_{sp}}$		

P	Q	R	S
(a) (ii)	(iii)	(iv)	(i)
(b) (iii)	(ii)	(i)	(iv)
(c) (ii)	(i)	(iv)	(iii)
(d) (i)	(iii)	(iv)	(ii)

SECTION - V

Assertion Reason Type

20. **Assertion** : The second dissociation constant of fumaric acid is greater than maleic acid.
Reason : Higher the dissociation constant of acid more is acidic character.
21. **Assertion** : The ionisation of hydrogen sulphide in water is low in the presence of hydrochloric acid.
Reason : Hydrogen sulphide is a weak acid.
22. **Assertion** : The pH of an aqueous solution of acetic acid remains unchanged on addition of sodium acetate.
Reason : The ionisation of acetic acid is increased by addition of sodium acetate.

SECTION - VI

Integer Value Correct Type

23. In 1 L saturated solution of $AgCl$ [$K_{sp}(AgCl) = 1.6 \times 10^{-10}$], 0.1 mol of $CuCl$ [$K_{sp}(CuCl) = 1.0 \times 10^{-6}$] is added. The resultant concentration of Ag^+ in the solution is 1.6×10^{-x} . The value of x is
24. Amongst the following, the total number of compounds whose aqueous solution turns red litmus paper blue is
 KCN K_2SO_4 $(NH_4)_2C_2O_4$
 NaCl $Zn(NO_3)_2$ $FeCl_3$
 K_2CO_3 NH_4NO_3 LiCN
25. Find the total number of diprotic acids among the following :
 H_3PO_4 , H_2SO_4 , H_3PO_3 , H_2CO_3 , $H_2S_2O_7$, H_3BO_3 , H_3PO_2 , H_2CrO_4 , H_2SO_3

THE p-BLOCK ELEMENTS

SECTION - I

Only One Option Correct Type

- Which of the following elements is not likely to act as central atom in MF_6^{3-} ?
(a) B (b) Al
(c) Ga (d) In
- An aqueous solution of boric acid is found to be weakly acidic in nature. This acidic character arises due to which of the following reasons?
(a) It is a protic acid which donates protons in aqueous solution.
(b) It is a Lewis acid which accepts OH^- from water and leaves H^+ to make the solution acidic.
(c) It gives metaboric acid when dissolved in water.
(d) It is prepared by reaction of borax with sulphuric acid hence, it behaves as an acid.
- Which of the following statements is false?
(a) Silicon carbide has a three dimensional structure with each silicon and carbon atoms being tetrahedrally surrounded by four atoms of the other kind.
(b) Carbon can form $C \equiv S$ bond because C has the ability to form $d\pi - d\pi$ bonds.
(c) Boron nitride has a structure similar to graphite.
(d) Graphite conducts electricity because of the availability of delocalised π -electrons.
- A solid element Y conducts electricity and forms two chlorides, YCl_n (a colourless volatile liquid) and YCl_{n-2} (a colourless solid). Then the element Y is
(a) C (b) Sn
(c) B (d) Al
- Aluminium oxide is not reduced by chemical reactions since
(a) reducing agent contaminate
(b) the process pollute the environment
(c) aluminium oxide is highly stable
(d) aluminium oxide is reactive.
- An alkali metal hydride (NaH) reacts with diborane in 'A' to give a tetrahedral compound 'B' which is extensively used as reducing agent in organic synthesis. The compound 'A' and 'B' respectively are
(a) CH_3COCH_3 and $B_3N_3H_6$
(b) $(C_2H_5)_2O$ and $NaBH_4$
(c) C_2H_6 and C_2H_5Na
(d) C_6H_6 and $NaBH_4$
- Predict X and Y in the following reaction?
 $3B_2H_6 + 6X \xrightarrow{\text{Low temp.}} 3[BH_2(X)_2]^+ [BH_4]^- \xrightarrow{473\text{ K}} Y + 12H_2$
(a) $X = NH_3$, $Y = B_3N_3H_6$
(b) $X = CO$, $Y = BH_3 \cdot CO$
(c) $X = NaH$, $Y = NaF$
(d) $X = NF_3$, $Y = B_3N_3$
- Which of the following statements is incorrect?
(a) Silicones are polymeric organosilicon compounds.
(b) SiF_4 has tetrahedral shape.
(c) Pb^{2+} is less stable than Sn^{2+} .
(d) Mg_2C_3 is an allylide type of ionic carbide.
- The catenation tendency of C, Si and Ge is in the order $Ge < Si < C$. The bond energies (in kJ mol^{-1}) of C-C, Si-Si and Ge-Ge are respectively
(a) 260, 297, 348 (b) 297, 260, 348
(c) 348, 260, 297 (d) 348, 297, 260
- Unlike the other elements of its group, carbon does not form CX_2 type molecules because
(a) energetically this is not possible
(b) carbon undergoes catenation
(c) it is non-metallic
(d) carbon does not contain d-orbital.

SECTION - II

More than One Options Correct Type

- Me_3SiCl is used during polymerisation of organo-silicones because
(a) the chain length of organosilicone polymers can be controlled by adding Me_3SiCl
(b) Me_3SiCl blocks the end terminal of silicone polymer
(c) Me_3SiCl improves the quality and yield of the polymer
(d) Me_3SiCl acts as a catalyst during polymerisation.
- Which of the following statements are correct?
(a) PbO_2 is a strong oxidising agent.
(b) CO_2 is non-poisonous and neutral in character.
(c) $GeCl_2$ is a strong reductant.
(d) $B(OH)_3$ gives weak alkaline solution.

13. Which of the following statements are correct?
- Group-13 elements show only one oxidation state *i.e.*, +3.
 - Tl(III) salts undergo disproportionation.
 - SiO₂ is covalent solid.
 - B(OMe)₃ imparts green colour to Bunsen flame.

SECTION - III

Paragraph Type

Paragraph for Questions 14 and 15

The maximum or the highest oxidation state shown by an element of *p*-block is equal to the total number of valence electrons. In Groups-13 and 14, the group oxidation state is the most stable for the lighter elements of the group but as we move down in a group, a lower oxidation state, which is two unit less than the group oxidation state, becomes progressively more stable for heavier elements. This is due to inert pair effect.

14. Which among the following is the strongest oxidising agent?
- SnO₂
 - SiO₂
 - GeO₂
 - PbO₂
15. Which of the following is the most acidic oxide?
- CO₂
 - SiO₂
 - PbO₂
 - SnO₂

Paragraph for Questions 16 and 17

Boron halides are electron-deficient compounds and act as Lewis acids. In boron halides, there is double bond character between boron and halogen atom because of the back bonding. Even the weakest bases will form adducts with the trihalides of boron. The rehybridisation of boron that accompanies adduct formation results in a loss of B—X double bond character.

16. In the adduct formation of trimethyl amine with boron halide,
- $$BX_3 + :N(CH_3)_3 \longrightarrow X_3B : N(CH_3)_3$$
- the enthalpy change is more negative in the case of
- BF₃
 - BCl₃
 - BBr₃
 - all are equal.
17. BF₃ exists as monomer but not as dimer because
- of back bonding
 - BF₃ is ionic compound
 - of steric hindrance
 - none of these.

SECTION - IV

Matching List Type

18. Match the List I with List II and select the correct answer using the codes given below the lists :

List I		List II	
(P) Diborane	(i)	Used as a flux for soldering metals	
(Q) Gallium	(ii)	Banana bonds	
(R) Borax	(iii)	Low melting and high boiling point, useful for measuring high temperatures	
(S) Aluminosilicate	(iv)	Used as a catalyst in petrochemical industries	

P	Q	R	S
(a) (ii)	(i)	(iii)	(iv)
(b) (ii)	(iii)	(i)	(iv)
(c) (i)	(iii)	(iv)	(ii)
(d) (iv)	(ii)	(i)	(iii)

19. Match the List I with List II and select the correct answer using the codes given below the lists :

List I		List II	
(P) Aluminosilicate glass	(i)	Cheap laboratory apparatus	
(Q) Calcium silicate	(ii)	Optical glasses	
(R) Lead glass	(iii)	Domestic glass for windows	
(S) Soda glass	(iv)	Low coefficient of expansion	

P	Q	R	S
(a) (iv)	(iii)	(ii)	(i)
(b) (i)	(iv)	(iii)	(ii)
(c) (ii)	(i)	(iv)	(iii)
(d) (iii)	(ii)	(i)	(iv)

SECTION - V

Assertion Reason Type

20. **Assertion :** If aluminium atoms replace a few silicon atoms in three dimensional network of silicon dioxide, the overall structure acquires a negative charge.
- Reason :** Aluminium is trivalent while silicon is tetravalent.

21. Assertion : Although aluminium is above hydrogen in electrochemical series, it is stable in air and water.

Reason : The thin protective layer of its oxide (Al_2O_3) on the surface, protects the aluminium.

22. Assertion : In CO_2 molecule, C-atom undergoes sp^2 hybridisation.

Reason : CO_2 molecule has net dipole moment.

SECTION - VI

Integer Value Correct Type

23. The total number of protons donated by one molecule of boric acid (H_3BO_3) is

24. The compound $(\text{CH}_3)_n\text{Si}(\text{Cl})_{4-n}$ on hydrolysis gives a branched chain silicone. The value of n is

25. The oxidation state of C in CO is x . The oxidation state of C in COCl_2 is y . Then, the value of $y - x$ is

SOLUTIONS

EQUILIBRIUM

1. (d): The equilibrium will remain unaffected in all three reactions on addition of small amount of argon (inert gas) at constant volume.

2. (d): $\text{PCl}_5(\text{g}) \rightleftharpoons \text{PCl}_3(\text{g}) + \text{Cl}_2(\text{g})$
 $1 - \alpha \qquad \qquad \alpha \qquad \qquad \alpha$
 (α = degree of dissociation)

Total moles = $1 - \alpha + \alpha + \alpha = 1 + \alpha$

$$K_p = \frac{p_{\text{PCl}_3} \times p_{\text{Cl}_2}}{p_{\text{PCl}_5}} = \frac{\left[\frac{\alpha}{1+\alpha} P \right] \left[\frac{\alpha}{1+\alpha} P \right]}{\frac{1-\alpha}{1+\alpha} P}$$

$$= \frac{\alpha^2 P}{1-\alpha^2} \Rightarrow \alpha = \left(\frac{K_p}{K_p + P} \right)^{1/2} \quad \dots \text{(i)}$$

Given, a moles of PCl_5 are heated then x moles of it dissociate at equilibrium.

Hence, $\alpha = \frac{x}{a}$

Thus, equation (i) becomes,

$$\frac{x}{a} = \left(\frac{K_p}{K_p + P} \right)^{1/2}$$

3. (d): $K_p = K_c (RT)^{\Delta n}$; $\Delta n = 2 - 3 = -1$
 $T = 350 \text{ K}$, $R = 0.083 \text{ L bar K}^{-1} \text{ mol}^{-1}$

$$K_c = \frac{K_p}{(RT)^{\Delta n}}$$

$$K_c = \frac{3 \times 10^{10} \text{ bar}^{-1}}{(0.083 \text{ L bar K}^{-1} \text{ mol}^{-1} \times 350 \text{ K})^{-1}}$$

$$= 3 \times 10^{10} \text{ bar}^{-1} \times 0.083 \text{ L bar K}^{-1} \text{ mol}^{-1} \times 350 \text{ K}$$

$$= 8.715 \times 10^{11} \text{ L mol}^{-1}$$

4. (a)

5. (a): Acidic strength $\propto K_a$

6. (b): K_{sp} for $\text{BaCO}_3 = [\text{Ba}^{2+}][\text{CO}_3^{2-}]$

Given, $[\text{CO}_3^{2-}] = 1 \times 10^{-4} \text{ M}$ (from Na_2CO_3)

$$K_{sp} = 5.1 \times 10^{-9}$$

$$\therefore 5.1 \times 10^{-9} = [\text{Ba}^{2+}] \times 10^{-4}$$

$$\Rightarrow [\text{Ba}^{2+}] = 5.1 \times 10^{-5} \text{ M}$$

Thus, at $[\text{Ba}^{2+}] = 5.1 \times 10^{-5} \text{ M}$, BaCO_3 will start precipitating.

7. (c): $\Delta H = 0$, it means reaction is independent of temperature.


8. (c): From given information,

$$\tan 75^\circ = 3.73 = \frac{\text{conc. of isobutane}}{\text{conc. of } n\text{-butane}} > K_c$$

So, reaction will proceed in backward direction.

9. (b): Degree of dissociation (α) = $\frac{D-d}{(n-1)d}$

where n = no. of moles of gaseous products produced by 1 mole of reactant.



1 mol 1 mol 1 mol

i.e., $n = 2$

$$D(\text{vapour density}) = \frac{\text{Mol. wt.}}{2} = \frac{208.5}{2} = 104.25$$

$$\alpha = \frac{104.25 - 70}{(2-1)70} = 0.489$$


Percentage dissociation = $0.489 \times 100 = 48.9\%$

10. (c): H_2SO_4 is a strong acid having $\text{pH} < 7$. NaNO_2 on hydrolysis gives alkaline solution of $\text{pH} > 7$. NaCl is a neutral salt so, $\text{pH} = 7$ and H_2S is a weak acid so, pH will be slightly lesser than 7.


11. (a, c): (a) For an endothermic reaction, K increases with increase in temperature.

$$\text{(c) At } 400 \text{ K, } Q = \frac{(p_{\text{NO}_2})^2}{(p_{\text{N}_2\text{O}_4})} = \frac{(20)^2}{2} = 200$$


As $Q > K$, hence, reaction proceeds in backward direction.



The equilibrium constant (K) is independent of the initial amount of CaCO_3 whereas at a given temperature, it is independent of pressure of CO_2 . ΔH is independent of catalyst and it depends on temperature.




$$\therefore K_1 = \frac{[\text{NH}_3]^2}{[\text{N}_2][\text{H}_2]^3}$$




$$\therefore K_2 = \frac{[\text{NH}_3]}{[\text{N}_2]^{1/2}[\text{H}_2]^{3/2}} = \sqrt{K_1} \Rightarrow K_1 = K_2^2$$

14. (b): I^- donates an electron pair hence, it acts as a Lewis base.

15. (b): $K_{a_1} > K_{a_2}$ and $K_{a_3} = 0$ (because H_3PO_3 is a dibasic acid).





$$K_{p_1} = 20 \times 20 = 400 \text{ mm}^2 \text{ and}$$



$$K_{p_2} = 30 \times 30 = 900 \text{ mm}^2$$

$$\therefore \frac{K_{p_1}}{K_{p_2}} = \frac{4}{9}$$

17. (c): In mixture, solids X and Y dissociate simultaneously,




$$K_{p_1} = p_A \times p_C = p_1(p_1 + p_2) \quad \dots(i)$$

$$K_{p_2} = p_B \times p_C = p_2(p_1 + p_2) \quad \dots(ii)$$

On dividing eqn. (i) by (ii), we get


$$\therefore \frac{K_{p_1}}{K_{p_2}} = \frac{p_1}{p_2} = \frac{4}{9} = \frac{y_A}{y_B}$$

18. (a)




$$K_{sp} = [\text{Hg}_2^{2+}][\text{Cl}^-]^2 = s(0.1 + 2s)^2 \approx (0.1)^2 s$$

$$\therefore s = 100 K_{sp}$$




$$K_{sp} = [\text{Pb}^{2+}][\text{I}^-]^2 = s(0.01 + 2s)^2 \approx s(0.01)^2 = 10^{-4} s$$

$$\therefore s = 10^4 K_{sp}$$



$$K_{sp} = [\text{Ag}^+]^2[\text{CrO}_4^{2-}] = (2s)^2(0.25 + s) \approx (2s)^2(0.25)$$

$$\text{or } s = \sqrt{K_{sp}}$$




$$K_{sp} = [\text{Ca}^{2+}][\text{C}_2\text{O}_4^{2-}] = s(0.1 + s) \approx 0.1s \text{ or } s = 10 K_{sp}$$

20. (a): Both fumaric and maleic acids have two ionisable H^+ ions *i.e.*, protons. The maleate monoanion shows intramolecular H-bonding, whereas fumarate monoanion shows intermolecular H-bonding. Thus, fumarate monoanion requires more energy to give fumarate dianion. Hence, second dissociation of fumaric acid is more than that of maleic acid.

21. (b)

22. (d): Addition of a small amount of CH_3COONa to acetic acid increases the pH value because CH_3COONa is a strong electrolyte and thus, ionises to give CH_3COO^- and Na^+ ions. Due to common ion effect, it suppresses the ionisation of CH_3COOH which decreases $[\text{H}^+]$ and hence, pH value increases.

23. (7): Let the solubility of AgCl and CuCl be a mol litre $^{-1}$ and b mol litre $^{-1}$ respectively.



$$\therefore K_{sp} \text{ of AgCl} = [\text{Ag}^+][\text{Cl}^-] = 1.6 \times 10^{-10} = a(a + b) \quad \dots(i)$$

$$\text{Similarly, } K_{sp} \text{ of CuCl} = [\text{Cu}^+][\text{Cl}^-] = 1.0 \times 10^{-6} = b(a + b) \quad \dots(ii)$$




Dividing eqn. (i) by (ii), we get

$$\frac{a}{b} = 1.6 \times 10^{-4} \text{ or } a = 1.6 \times 10^{-4} \times b$$




Substituting the value of a in eqn. (i), we get

$$\begin{aligned} 1.6 \times 10^{-10} &= 1.6 \times 10^{-4} b(1.6 \times 10^{-4} b + b) \\ \Rightarrow 10^{-6} &= b^2(1.6 \times 10^{-4} + 1) \\ \Rightarrow b &= 10^{-3} [\because 1.6 \times 10^{-4} \ll 1] \Rightarrow a = 1.6 \times 10^{-7} \\ [\text{Ag}^+] &= a = 1.6 \times 10^{-7}; \text{Comparing with } 1.6 \times 10^{-x} \\ \Rightarrow x &= 7 \end{aligned}$$

24. (3): Salts of strong base and weak acid give basic solution.



25. (6): Diprotic acids are those which furnish two protons per molecule like, H_2SO_4 , H_3PO_3 , H_2CO_3 , $\text{H}_2\text{S}_2\text{O}_7$, H_2CrO_4 , H_2SO_3 .



THE *p*-BLOCK ELEMENTS

- (a): Boron cannot expand its coordination number beyond 4 due to absence of *d*-orbitals. Therefore, it can not act as central metal atom in MF_6^{3-} .
- (b): It reacts with water to accept OH^- ions and releases H_3O^+ ions to make acidic solution.
 $\text{B(OH)}_3 + 2\text{H}_2\text{O} \longrightarrow \text{H}_3\text{O}^+ + [\text{B(OH)}_4]^-$; $pK_a = 9.25$
- (b): Carbon does not have *d*-orbitals to form *dπ-dπ* bonds.
- (b): Element Y is tin which conducts electricity and can exist in +2 and +4 oxidation state. SnCl_4 is colourless volatile liquid whereas SnCl_2 is colourless ionic solid.
- (c)
- (b): $2\text{NaH} + \text{B}_2\text{H}_6 \xrightarrow{(\text{C}_2\text{H}_5)_2\text{O(A)}} 2\text{NaBH}_4$
Sodium borohydride
(B)
- (a): $3\text{B}_2\text{H}_6 + 6\text{NH}_3 \xrightarrow{(X)} 3[\text{BH}_2(\text{NH}_3)_2]^+[\text{BH}_4]^-$
Heat
↓
 $2\text{B}_3\text{N}_3\text{H}_6 + 12\text{H}_2$
(Y)
- (c): Stability of +2 oxidation state increases down the group-14 due to inert pair effect thus, Pb^{2+} is more stable than Sn^{2+} .
- (d): Catenation tendency decreases with the decrease in *M-M* bond energy. Hence, bond

energies of C-C, Si-Si and Ge-Ge are 348, 297 and 260 kJ/mol respectively.

- (a): The elements of group-14 show tetravalency by sharing four of its valence electrons. They have O.S. of +4. Ge, Sn and Pb also show +2 O.S. The stability of dihalides increases down the group. The stability of dihalides follow the sequence :
 $\text{CX}_2 \ll \text{SiX}_2 \ll \text{GeX}_2 \ll \text{SnX}_2 \ll \text{PbX}_2$
- (a, b): The chain length of the polymer can be controlled by adding Me_3SiCl which blocks the ends of silicone polymer.
- (a, c): In PbO_2 , oxidation state of Pb is +4, passes readily into more stable oxidation state +2 (due to inert pair effect) and hence, is a strong oxidising agent. On the contrary, GeCl_2 is a strong reducing agent.
- (c, d)
- (d): PbO_2 is the strongest oxidising agent as it readily changes from +4 to +2 oxidation state.
- (a)
- (c)
- (a): Due to *pπ-pπ* back bonding, B-F bond possess some double bond character.
- (b)
- (a)
- (a)
- (a)
- (d): In CO_2 molecule, carbon atom undergoes *sp* hybridisation. The two *sp* hybridised orbitals of C-atom overlap with two *p*-orbitals of O-atoms to make 2 σ bonds while other two electrons of C-atom are involved in *pπ-pπ* bonding with oxygen atom. Hence, it is linear in shape with equal C-O bond lengths and no dipole moment.

$$:\ddot{\text{O}} \equiv \text{C} \equiv \ddot{\text{O}}: \therefore \mu = 0$$
- (0): H_3BO_3 is not protonic acid. It is a Lewis acid as it can accept a pair of electrons from OH^- ion of water.
- (1): To give branched chain silicone, the number of Cl atoms must be three to give $\text{CH}_3\text{Si(OH)}_3$, which on condensation gives branched silicone.
- (2): Oxidation state of C in CO is +2
 $\therefore x = +2$
 Oxidation state of C in COCl_2 is +4
 $\therefore y = +4$
 Now, $y - x = 4 - 2 = 2$



ACE

YOUR WAY CBSE XI



Series 6

CHAPTERWISE PRACTICE PAPER : THE *p*-BLOCK ELEMENTS | ORGANIC CHEMISTRY-SOME BASIC PRINCIPLES AND TECHNIQUES

Time Allowed : 3 hours

Maximum Marks : 70

GENERAL INSTRUCTIONS

- (i) All questions are compulsory.
- (ii) Q. no. 1 to 5 are very short answer questions and carry 1 mark each.
- (iii) Q. no. 6 to 10 are short answer questions and carry 2 marks each.
- (iv) Q. no. 11 to 22 are also short answer questions and carry 3 marks each.
- (v) Q. no. 23 is a value based question and carries 4 marks.
- (vi) Q. no. 24 to 26 are long answer questions and carry 5 marks each.
- (vii) Use log tables if necessary, use of calculators is not allowed.

1. What is the maximum covalency of silicon in its compounds?
 2. Give the IUPAC name of the saturated hydrocarbon having the following carbon skeleton.

$$\begin{array}{ccccccc} \text{C} & - & \text{C} & - & \text{C} & - & \text{C} & - & \text{C} & - & \text{C} \\ & & | & & | & & | & & | & & \\ \text{C} & - & \text{C} & - & \text{C} & & & & & & \end{array}$$
 3. What is inert pair effect?
 4. How many σ bonds and π bonds are present in the second member of the alkene series?
 5. What is catenation?
 6. Like CO, its analogue SiO is not stable. Why?
 7. By mistake, an alcohol (boiling point 97°C) was mixed with a hydrocarbon (boiling point 68°C). Suggest a suitable method to separate the two compounds. Explain the reason for your choice.
 8. How does NaBH_4 react with iodine? Also discuss the structure of the main product formed.
 9. What are reactive intermediates? How are they generated by bond fission?
 10. Will CCl_4 give white precipitate of AgCl on heating it with silver nitrate? Give reason for your answer.
- OR**
- In DNA and RNA, nitrogen atom is present in the ring system. Can Kjeldahl method be used for the estimation of nitrogen present in these? Give reason.
11. What happens when
 - (i) Borax is heated strongly
 - (ii) Boric acid is added to water
 - (iii) Aluminium is treated with HCl ?
 12. Identify the reagents shown in brackets in the following equations as nucleophiles or electrophiles :
 - (a) $\text{CH}_3\text{COOH} + (\text{NH}_3) \xrightarrow{\Delta} \text{CH}_3\text{CONH}_2 + \text{H}_2\text{O}$
 - (b) $\text{CH}_3\text{COCH}_3 + (\text{CN}^-) \longrightarrow (\text{CH}_3)_2\text{C}(\text{CN})(\text{OH})$
 - (c) $\text{C}_6\text{H}_6 + (\text{CH}_3\overset{+}{\text{C}}\text{O}) \longrightarrow \text{C}_6\text{H}_5\text{COCH}_3$
- OR**
- How many cyclic and acyclic isomers are possible for the molecular formula $\text{C}_3\text{H}_6\text{O}$? Draw the structures also.

13. A non-metallic element of group 13, used in making bullet proof vests is extremely hard solid of black colour. It can exist in many allotropic forms and has unusually high melting point. Its trifluoride acts as Lewis acid towards ammonia. The element exhibits maximum covalency of four. Identify the element and write the reaction of its trifluoride with ammonia. Explain why does the trifluoride act as a Lewis acid.
14. Explain the terms inductive and electromeric effects. Which electron displacement effect explains the following orders of acidity of the carboxylic acids?
 (a) $\text{Cl}_3\text{CCOOH} > \text{Cl}_2\text{CHCOOH} > \text{ClCH}_2\text{COOH}$
 (b) $\text{CH}_3\text{CH}_2\text{COOH} > (\text{CH}_3)_2\text{CHCOOH} > (\text{CH}_3)_3\text{CCOOH}$
15. How will you prepare the following in steps not exceeding three (give reactions only)?
 (i) Silicon from SiCl_4
 (ii) Linear silicon containing methyl groups only from Si
 (iii) Hydrofluorosilicic acid from SiO_2
16. What is the difference between distillation, distillation under reduced pressure and steam distillation?
17. Rationalise the given statements and give chemical reactions :
 (i) Lead (II) chloride reacts with Cl_2 to give PbCl_4 .
 (ii) Lead (IV) chloride is highly unstable towards heat.
 (iii) Lead is known not to form an iodide, PbI_4 .
18. Arrange the following :
 (i) $\text{C}_6\text{H}_5\overset{+}{\text{C}}\text{HCH}_3$, $\text{C}_6\text{H}_5\overset{+}{\text{C}}\text{HCH}=\text{CH}_2$, $\text{C}_6\text{H}_5\overset{+}{\text{C}}\text{H}_2\text{CH}_2$, $\text{C}_6\text{H}_5\overset{+}{\text{C}}(\text{CH}_3)_2$ in order of increasing stability.
 (ii) $\text{CH}_3\overset{+}{\text{C}}\text{H}_2$, $\text{C}_6\text{H}_5\overset{+}{\text{C}}\text{H}_2$, $(\text{CH}_3)_3\overset{+}{\text{C}}$, $\text{CH}_2=\overset{+}{\text{C}}\text{HCH}_2$ in order of decreasing stability.
 (iii) $\text{HC}\equiv\text{C}^-$, $\text{CH}_2=\overset{-}{\text{C}}\text{H}$, $\text{CH}_3\overset{-}{\text{C}}\text{H}_2$, CH_3 , $(\text{CH}_3)_2\overset{-}{\text{C}}\text{H}$, $\text{C}_6\text{H}_5\overset{-}{\text{C}}\text{H}_2$ in order of increasing stability.
19. Draw the structure of boric acid showing hydrogen bonding. In which form it is present in water? What is the hybridisation of boron in this species?
20. What is meant by hybridisation? Compound $\text{CH}_2=\text{C}=\text{CH}_2$ contains *sp* or *sp*² hybridised carbon atoms. Will it be a planar molecule?
21. H_3O^+ or R_4N^+ neither acts as an electrophile nor as a nucleophile. Explain why.
22. Which of the following is not hydrolysed by water and why?
 BF_3 , BCl_3 and BBr_3
23. Students of Class XI, were given different compounds for the detection of extra elements present in the compound by their science teacher. For this, all the students prepared sodium extract and used it for detection. Pratiksha added a small amount of freshly prepared ferrous sulphate to a part of sodium extract and warmed it. She then added few drops of FeCl_3 and acidified the resultant solution with dil. HCl. She got a bluish green colouration. Her fellow mate, Ritwik followed the same sequence of steps as followed by Pratiksha but he got a blood red colouration. Ritwik thought he has performed the test incorrectly. Pratiksha explained him the reason for different colouration. Now, answer the following questions :
 (a) What did Pratiksha explain him?
 (b) Write the reactions involved in both the cases.
 (c) What are the values shown by Pratiksha?
24. Describe the general trends in the following properties of the elements in Group 13 and 14.
 (i) Atomic size (ii) Ionisation enthalpy
 (iii) Metallic character (iv) Oxidation states
 (v) Nature of halides
- OR**
- When BCl_3 is treated with water, it hydrolyses and forms $[\text{B}(\text{OH})_4]^-$ only whereas AlCl_3 in acidified aqueous solution forms $[\text{Al}(\text{H}_2\text{O})_6]^{3+}$ ion. Why? Explain what is the hybridisation of boron and aluminium in these species?
25. Draw the resonance structures for the following compounds. Show the electron shift using curved arrow notation.
 (i) $\text{C}_6\text{H}_5\text{OH}$ (ii) $\text{C}_6\text{H}_5\text{NO}_2$
 (iii) $\text{CH}_3\text{CH}=\overset{+}{\text{C}}\text{HCHO}$
 (iv) $\text{C}_6\text{H}_5-\overset{+}{\text{C}}\text{H}_2$ (v) $\text{CH}_3\text{CH}=\overset{+}{\text{C}}\text{HCH}_2$
- OR**
- Answer the following questions :
 (a) Benzoic acid is an organic compound. Its crude sample can be purified by crystallisation from hot water. What characteristic differences in the properties of benzoic acid and the impurity make this process of purification suitable?

- (b) An organic compound contains 69% carbon and 4.8% hydrogen, the remainder being oxygen. Calculate the masses of carbon dioxide and water produced when 0.20 g of this substance is subjected to complete combustion.

26. Answer the following questions :

- (a) Explain why BF_3 exists but BH_3 does not.
 (b) Gallium has higher ionization enthalpy than aluminium. Why?
 (c) AlF_3 is high melting solid but AlCl_3 is low melting. Explain.


OR

Answer the following :

- (a) Aluminium dissolves in mineral acids and aqueous alkalis and thus shows amphoteric character. A piece of aluminium foil is treated with dilute hydrochloric acid or dilute sodium hydroxide solution in a test tube and on bringing a burning matchstick near the mouth of the test tube, a pop sound indicates the evolution of hydrogen gas. The same activity when performed with concentrated nitric acid, reaction does not proceed. Explain the reason.
 (b) Identify the compounds A, X and Z in the following reactions :
 (i) $\text{A} + 2\text{HCl} + 5\text{H}_2\text{O} \longrightarrow 2\text{NaCl} + \text{X}$
 (ii) $\text{X} \xrightarrow[>370\text{ K}]{\Delta} \text{HBO}_2 \xrightarrow{\text{Red heat}} \text{Z}$

SOLUTIONS


1. Silicon shows maximum covalency in its compounds equal to six e.g., Na_2SiF_6 .
 2. 5-Ethyl-2, 3, 4-trimethyloctane
 3. In the elements of 4th, 5th and 6th period of the *p*-block, ns^2 -electrons remain more tightly held by the nucleus due to poor shielding of *d*- and *f*-electrons and hence, do not participate in bonding. This is called inert pair effect.
 4. The second member of the alkene series is propene. The structural formula of the propene is



5. Property of self-linking of atoms to form chains or rings of various sizes is called catenation. Carbon

has the maximum tendency for catenation due to high bond strength of C—C bonds.


6. CO has the following resonating structures :




Therefore, CO has $p\pi$ - $p\pi$ multiple bonds. This is because C being small in size and having high electronegativity than Si forms strong $p\pi$ - $p\pi$ bond. Silicon on the otherhand, because of its larger size and lower electronegativity has no tendency to form $p\pi$ - $p\pi$ bond and hence, does not form stable SiO.

7. The difference in boiling points of two liquids is more than 25°C. Hence, simple distillation can be used and since at the boiling point of low boiling liquid, the vapours would consist entirely of only low boiling liquid without any contamination of vapours of high boiling liquid and *vice-versa*. Thus, both the liquids can be distilled without any decomposition.




8. When NaBH_4 reacts with iodine diborane is produced :



In the structure of diborane four hydrogen atoms, two on the left and two on the right, known as terminal hydrogens and two boron atoms lie in the same plane. Two hydrogen atoms forming bridges, one above and other below, lie in a plane perpendicular to the rest of molecule.



9. Reactive intermediates are short-lived, high-energy, highly reactive molecules generated in a chemical reaction. These molecules quickly convert into a more stable product e.g., carbocation, carbanion and free radical etc. Carbocation and carbanion are generated by the heterolytic bond fission while free radical is produced by homolytic bond fission.




10. No. AgNO_3 solution is ionic in nature. It contains Ag^+ ion which when reacts with Cl^- ions, produces a white ppt. of AgCl . While CCl_4 is a covalent compound and does not produce Cl^- ions. Therefore, when AgNO_3 is added to it, it does not produce a white ppt. of AgCl .


OR

Nitrogen atom present in DNA and RNA cannot be estimated by Kjeldahl method. DNA and RNA have nitrogen in the heterocyclic rings. Nitrogen present in rings, azo and nitro groups cannot be converted into ammonium sulphate. Thus, Kjeldahl method cannot be used for the estimation of nitrogen present in DNA and RNA.



11. (i) On heating, borax first loses water molecules and swells up. On further heating it turns into a transparent liquid which solidifies into glass like material called borax bead.




- (ii) It acts as Lewis acid by accepting electrons from a hydroxyl ion.



- (iii) Hydrogen gas is liberated.




Here, NH_3 is acting as a nucleophile.



Here, CN^- is acting as a nucleophile.

The carbonyl carbon is a positive centre, since the oxygen withdraws the carbonyl electron and leaves the carbon atom electron deficient.





CH_3CO^+ is an electrophile.

Benzene being an electron rich molecule, is most suitable for an electrophilic attack.



OR

Seven following isomers are possible :


- (a) Acyclic isomers :



- (b) Cyclic isomers :




13. The only non-metallic element of group 13 is boron. It is an extremely hard black substance and is used in making bullet proof vests. It exists in many allotropic forms and has unusually high melting point. Since B does not have vacant *d*-orbitals, therefore, it can exhibit maximum covalency of four.




Since B in BF_3 has only six electrons in its valence shell, therefore, it needs two more electrons to complete its octet. Thus, BF_3 acts as a Lewis acid.

14. **Inductive effect** : Polarisation of σ -bond due to the electron donating or withdrawing nature of groups attached is called inductive effect.


e.g. Attachment of Cl, an electronegative atom to an ethane molecule causes the carbons to carry a δ^+ charge.



Electromeric effect : In the presence of an attacking reagent, the π -electron cloud in a multiple bonded compound tends to get polarized. Such an effect is termed as temporary effect and the original condition is restored if the reagent is removed.








(a) The given order of acidity can be explained by $-I$ effect of chlorine atom. More the no. of Cl atoms, higher is the acidic strength.




(b) The given order of acidity can be explained by $+I$ effect of $-\text{CH}_3$ group.

More the number of $-\text{CH}_3$ groups on the α -carbon, lower will be the acidic strength.



SiF_4 thus formed dissolves in HF to form hydrofluorosilicic acid.



16. Differences between distillation, distillation under reduced pressure and steam distillation may be summarized as :




Criterion for difference	Distillation	Distillation under reduced pressure	Steam distillation
Used for	Separation of volatile liquids from non-volatile impurities and the liquids having sufficient difference in their boiling points.	Purify liquids having very high boiling points or those, which decompose at or below their boiling points.	To separate substances which are immiscible with water and are steam volatile.
Method	Simple heating of the mixture in a round bottom flask fitted with condenser allows the lower boiling liquid to be collected first and the higher boiling one is collected later.	Since, one of the component decomposes before reaching its boiling point the heating is carried out under reduced pressure so that boiling occurs at a lower temperature and the component is collected in pure form.	Mixture of steam and volatile organic compound is condensed and collected. The compound is later separated from water using a separating funnel.
Example	A mixture of chloroform (b.pt. 334 K) and aniline (b.pt. 457 K).	Separation of glycerol from spent lye in soap industry.	Separation of aniline from aniline-water mixture.

17. (i) $\text{PbCl}_2 + \text{Cl}_2 \nrightarrow \text{PbCl}_4$
Lead is more stable in +2 oxidation state than in +4 state due to inert pair effect. Thus, the reaction is not feasible.


(ii) $\text{PbCl}_4 \xrightarrow{\Delta} \text{PbCl}_2 + \text{Cl}_2$
Pb is more stable in its +2 oxidation state due to inert pair effect. As a result, when subjected to heat, Pb (IV) goes to Pb (II) state.

(iii) $\text{Pb} + 2\text{I}_2 \longrightarrow \text{PbI}_4$
 I^- is a good reducing agent and therefore,

reduces Pb (IV) to Pb (II) easily. That is why, PbI_4 does not exist.




19. Boric acid has a layer structure in which planar BO_3 units are joined by hydrogen bonds.



Structure of boric acid; the dotted lines represent hydrogen bonds

It is present in water as $[\text{B}(\text{OH})_4]^-$ unit. Boron has sp^3 hybridisation in $[\text{B}(\text{OH})_4]^-$ unit.

20. Intermixing of atomic orbitals of same atom with slightly different energies resulting in the formation of same number of new orbitals with equal energies and shape is known as hybridisation.




Central carbon atom is sp hybridised and its two unhybridised p -orbitals are perpendicular to each other. The p -orbitals in one plane overlap with one of the p -orbitals of left terminal carbon atom and the p -orbital in other plane overlaps with p -orbital of right side terminal carbon atom. As a result the position of two terminal carbon atoms and the hydrogen atoms attached to them are fixed in planes perpendicular to each other. Therefore, the pair of hydrogen atoms attached to terminal carbon atoms are present in different planes and hence molecule is not planar.

21. $\text{H}_3\ddot{\text{O}}^+$ has a lone pair of electrons but due to the presence of +ve charge, it cannot donate its electron pair and hence it does not act as a nucleophile. R_4N^+ , however, does not have a lone pair of electrons, therefore, it does not act as a nucleophile. $\text{H}_3\ddot{\text{O}}^+$ has 8 electrons in the valence shell. It cannot expand its valence shell beyond 8 due to the absence of d -orbitals. Therefore, it does not act as an electrophile. Similarly, R_4N^+ also has 8 electrons in the valence shell. Like O, N also cannot expand

its valence shell beyond 8 and hence it also does not act as an electrophile.




Thus, H_3O^+ or R_4N^+ neither acts as a nucleophile nor as an electrophile.

22. BF_3 is not easily hydrolysed by water. It forms an adduct $\text{F}_3\text{B} \leftarrow \text{OH}_2$ whereas BCl_3 and BBr_3 are hydrolysed to boric acid and HCl or HBr respectively. This is because the $\text{B}-\text{F}$ bond in BF_3 is very strong due to extensive $p\pi-p\pi$ back bonding. As a result, it is not hydrolysed by water. The $\text{B}-\text{F}$ bond energy is far larger than $\text{B}-\text{OH}$ bond energy and cannot be compensated.





However, in BCl_3 and BBr_3 , the corresponding $\text{B}-\text{Cl}$ and $\text{B}-\text{Br}$ bond energy is relatively less than $\text{B}-\text{F}$ because of inefficient $p\pi-p\pi$ back bonding. Therefore, these get hydrolysed.



23. (a) Pratiksha explained Ritwik that he has performed the experiment correctly. The compound she got contained only nitrogen whereas Ritwik's compound had both nitrogen and sulphur.



Sodium hexacyanoferrate (II)



In second case :



- (c) Knowledge and teaching aptitude are the values shown by Pratiksha.
24. (i) **Atomic size** : Atomic radii of group 13 elements increase down the group with exception $\text{Ga} < \text{Al}$ due to presence of extra ten d -electrons which offer poor screening effect. In group 14, there is a considerable

increase in radius from C to Si, thereafter from Si → Pb a small increase is seen due to presence of completely filled *d*- and *f*- orbitals.

- (ii) **Ionisation enthalpy** : For group 13, the trend is $B > Al < Ga > In < Tl$. This is due to increase in size and low screening effect of *d*- and *f*- electrons.

Group 14 elements have higher I.E. than group 13 elements. First I.E. decreases from C to Sn. The exceptions of increase in I.E. from Sn → Pb is due to poor shielding effect of *d*- and *f*- orbitals.

- (iii) **Metallic character** : Metallic character increases from boron to aluminium then decreases down the group for group 13 elements. Due to smaller size group 14 elements are less metallic. Metallic character increases gradually down the group. C (non-metal), Si, Ge (metalloid) Sn, Pb (metal).



- (iv) **Oxidation states** : For group 13 both +1 and +3 oxidation states are observed. +1 oxidation state becomes more stable as we move down the group due to inert pair effect. Boron does not show +3 oxidation states.

For group 14 common oxidation states are +4 and +2. Tendency to show +2 oxidation state increases down the group.

- (v) **Nature of halides** : Group 13 elements form trihalides (except TlI_3). Due to electron deficient nature BCl_3 accepts electrons and forms adduct. $AlCl_3$ achieves stability by forming a dimer. Group 14 elements form halides with formula MX_2 and MX_4 . Except CCl_4 other halides are easily hydrolysed. Stability of dihalides increases down the group.


OR

When BCl_3 reacts with water, it first undergoes hydrolysis to form boric acid, $B(OH)_3$. Due to small size and high electronegativity of B, $B(OH)_3$ polarizes H_2O molecule accepting an OH^- ion to form $[B(OH)_4]^-$ species and releasing a proton.



Since B lies in the 2nd period, it has only one *s*- and three *p*-orbitals but no *d*-orbitals. In other words, at the maximum, it can have four pairs of electrons in the valence shell, *i.e.*, its maximum coordination number is 4. That is why, it accepts one OH^- ion forming $[B(OH)_4]^-$ in which boron


is sp^3 -hybridized. In contrast, $AlCl_3$, undergoes hydrolysis in acidified aqueous solution to form $[Al(H_2O)_6]^{3+}$





This may be explained as follows : (i) In acidic medium, the conc. of OH^- ions is much lower than that of H^+ ions, therefore, Al^{3+} ions coordinate with H_2O molecules and not with OH^- ions.

(ii) Due to the presence of vacant *d*-orbitals in Al^{3+} ions, it can expand its coordination number from 4 to 6 and hence forms $[Al(H_2O)_6]^{3+}$ in which hybridization of Al is sp^3d^2 .

E.C. of Al : $[Ne]3s^23p^1$







Formation of $[Al(H_2O)_6]^{3+}$



MPP-8 CLASS XI ANSWER KEY

1. (b) 2. (d) 3. (d) 4. (a) 5. (a)
6. (c) 7. (d) 8. (c) 9. (b) 10. (d)
11. (c) 12. (c) 13. (b) 14. (a) 15. (c)
16. (a) 17. (d) 18. (d) 19. (b) 20. (b,c)
21. (a,b,c) 22. (a,b,c,d) 23. (b,d) 24. (3) 25. (6)
26. (2) 27. (c) 28. (c) 29. (c) 30. (b)



OR

(a) Benzoic acid can be purified by hot water because of the following characteristics :

- Benzoic acid is more soluble in hot water and less soluble in cold water.
- Impurities present in benzoic acid are either insoluble in water or are more soluble in water to such an extent that they remain in solution as the mother liquor upon crystallisation.

(b) $\% \text{C} = \frac{12}{44} \times \frac{\text{Mass of CO}_2 \text{ formed}}{\text{Mass of substance taken}} \times 100$

$$69 = \frac{12}{44} \times \frac{\text{Mass of CO}_2 \text{ formed}}{0.2} \times 100$$


$$\therefore \text{Mass of CO}_2 \text{ formed} = \frac{69 \times 44 \times 0.2}{12 \times 100} = 0.506 \text{ g}$$

$$\% \text{H} = \frac{2}{18} \times \frac{\text{Mass of H}_2\text{O formed}}{\text{Mass of substance taken}} \times 100$$

$$4.8 = \frac{2}{18} \times \frac{\text{Mass of H}_2\text{O formed}}{0.2} \times 100$$

$$\therefore \text{Mass of H}_2\text{O formed} = \frac{4.8 \times 18 \times 0.2}{2 \times 100} = 0.0864 \text{ g}$$

26. (a) In BF_3 , the vacant $2p$ -orbital of boron accepts π electron density by overlapping with filled $2p_z$ orbital of F. Therefore, the stability of BF_3 is due to back bonding.





However, hydrogen atoms do not have free electrons to form back bonding with boron and therefore, BH_3 is not stable. However, to satisfy the electron deficiency of boron, it dimerizes to form B_2H_6 .


- Gallium has higher ionization enthalpy than aluminium because of higher effective nuclear charge. This is due to additional ten d -electrons which do not shield the nuclear charge effectively so that the outer electrons are strongly held.
- AlF_3 is high melting solid because it is ionic in nature. On the other hand, AlCl_3 is covalent in nature and hence is a low melting solid.

OR



(a) Al being amphoteric dissolves both in acids and alkalis evolving H_2 gas which burns with a pop sound.



With conc. HNO_3 , Al becomes passive and the reaction does not proceed. This passivity is due to the formation of a thin protective layer of its oxide (Al_2O_3) on the surface of the metal which prevents further action.



- (i) $\text{Na}_2\text{B}_4\text{O}_7 + 2\text{HCl} + 5\text{H}_2\text{O} \longrightarrow 2\text{NaCl} + 4\text{H}_3\text{BO}_3$
Borax (A) Boric acid (X)



MPP-8 MONTHLY Practice Problems

Class XI

This specially designed column enables students to self analyse their extent of understanding of specified chapter. Give yourself four marks for correct answer and deduct one mark for wrong answer. Self check table given at the end will help you to check your readiness.



The *p*-Block Elements

Total Marks : 120

Time Taken : 60 Min.

NEET / AIIMS

Only One Option Correct Type

- 2 g of aluminium is treated separately with excess of dilute H_2SO_4 and excess of NaOH . The ratio of the volumes of hydrogen evolved is
(a) 2 : 3 (b) 1 : 1 (c) 2 : 1 (d) 1 : 2
- First ionisation enthalpy for the elements of group-13 follows the order
(a) $\text{B} > \text{Al} > \text{Ga} > \text{In} > \text{Tl}$
(b) $\text{B} < \text{Al} < \text{Ga} < \text{In} < \text{Tl}$
(c) $\text{B} < \text{Al} > \text{Ga} < \text{In} > \text{Tl}$
(d) $\text{B} > \text{Al} < \text{Ga} > \text{In} < \text{Tl}$
- The stability of +1 oxidation state increases in the sequence
(a) $\text{Tl} < \text{In} < \text{Ga} < \text{Al}$ (b) $\text{In} < \text{Tl} < \text{Ga} < \text{Al}$
(c) $\text{Ga} < \text{In} < \text{Al} < \text{Tl}$ (d) $\text{Al} < \text{Ga} < \text{In} < \text{Tl}$
- Boron cannot form which one of the following anions?
(a) BF_6^{3-} (b) BH_4^-
(c) $\text{B}(\text{OH})_4^-$ (d) BO_2^-
- Which is not correct?
(a) GeCl_2 is more stable than GeCl_4 .
(b) $\text{Ge}(\text{OH})_2$ is amphoteric.
(c) GeO_2 is weakly acidic.
(d) GeCl_4 in HCl forms $[\text{GeCl}_2]^{2-}$ ion.
- Graphite is a soft solid lubricant extremely difficult to melt because graphite
(a) is an allotropic form of carbon
(b) is a non-crystalline substance
(c) has carbon atoms arranged in large plates of rings of strongly bound carbon atoms with weak interplate bonds
(d) has molecules of variable molecular masses like polymers.
- The transformation of metallic tin into grey powder is due to
(a) the interaction with nitrogen of the air at very low temperature
(b) the interaction with water vapour contained in the humid air
(c) a change in the partial pressure of oxygen in air
(d) a change in the crystalline structure of tin.
- Which of the following is incorrect for the fact, 'carbon differs from other elements of the group'?
(a) It has significant tendency to form long chains.
(b) It has unique ability to form multiple bonds.
(c) It has *d*-orbital in penultimate shell.
(d) Maximum coordination number of carbon is 4.
- In aluminium extraction by the Baeyer's process, alumina is extracted from bauxite by sodium hydroxide at high temperature and pressure.
$$\text{Al}_2\text{O}_{3(s)} + 2\text{OH}^-_{(aq)} \longrightarrow 2\text{AlO}_2^-_{(aq)} + \text{H}_2\text{O}_{(l)}$$

Solid impurities such as Fe_2O_3 and SiO_2 are removed and then $\text{Al}(\text{OH})_4^-$ is reprecipitated.
$$2\text{Al}(\text{OH})_4^- \longrightarrow \text{Al}_2\text{O}_3 \cdot 3\text{H}_2\text{O} + 2\text{OH}^-$$

In the industrial world
(a) carbon dioxide is added to precipitate the alumina
(b) temperature and pressure are dropped and the supersaturated solution is seeded
(c) both (a) and (b) are practised
(d) water is evaporated.
- In diborane, the number of electrons that account for bonding in the bridges is
(a) six (b) two (c) eight (d) four.

11. Which of the following statements is incorrect with respect to the structure of diborane?
- There are four terminal hydrogen atoms and two bridged hydrogen atoms.
 - The terminal hydrogen atoms and boron atoms lie in a plane.
 - There are five two-centred and five three-centred bonds.
 - Two bridged hydrogen atoms lie above and below the plane.
12. What is not true about borax?
- Molecular formula is $\text{Na}_2\text{B}_4\text{O}_7 \cdot 10\text{H}_2\text{O}$.
 - Crystalline borax contains tetranuclear units of $[\text{B}_4\text{O}_5(\text{OH})_4]^{2-}$.
 - It hydrolyses to give an acidic solution.
 - It is a white crystalline solid.

Assertion & Reason Type

Directions : In the following questions, a statement of assertion is followed by a statement of reason. Mark the correct choice as:

- If both assertion and reason are true and reason is the correct explanation of assertion.
- If both assertion and reason are true but reason is not the correct explanation of assertion.
- If assertion is true but reason is false.
- If both assertion and reason are false.

13. **Assertion :** Silicones are water repelling in nature.
Reason : Silicones are organosilicon polymers, which have $(\text{R}_2\text{SiO})_n$ as repeating unit.

14. **Assertion :** Borazole is aromatic in nature.
Reason : Nitrogen contributes π -electrons to the system.

15. **Assertion :** Molten AlBr_3 is poor conductor of electricity.
Reason : AlBr_3 being ionic in nature provides Al^{3+} and Br^- ions.

JEE MAIN / JEE ADVANCED

Only One Option Correct Type

16. Diborane, B_2H_6 , reacts with water to form boric acid and hydrogen. What is the pH of the solution which results when 1.104 g of B_2H_6 reacts with 100 mL water? Assume the final volume to be 100 mL.
(Given : K_a of $\text{H}_3\text{BO}_3 = 8 \times 10^{-10}$, $\text{p}K_a = 9.1$)
- 4.6
 - 6.2
 - 8.9
 - 7.2

17. Which one of the following is the correct statement?
- $\text{B}_2\text{H}_6 \cdot 2\text{NH}_3$ is known as 'inorganic benzene'.
 - Boric acid is a protonic acid.
 - Beryllium exhibits coordination number six.
 - Chlorides of both beryllium and aluminium have bridged chloride structures in solid phase.

18. X reacts with NaOH (aqueous solution) to form Y and H_2 . Aqueous solution of Y is heated to 323-333 K and on passing CO_2 into it, Na_2CO_3 and Z are formed. When Z is heated to 1200 °C, Al_2O_3 is formed. X, Y and Z respectively are

- Al, AlCl_3 , NaAlO_2
- Zn, Na_2ZnO_2 , $\text{Al}(\text{OH})_3$
- Al, $\text{Al}(\text{OH})_3$, AlCl_3
- Al, NaAlO_2 , $\text{Al}(\text{OH})_3$

19. $\text{NH}_3 + \text{CO}_2 \xrightarrow{\text{pressure}/\Delta} X \xrightarrow{\Delta} Y$

X and Y are respectively

- $(\text{NH}_4)_2\text{CO}_3$, $\text{NH}_2\text{COONH}_4$
- $\text{NH}_2\text{COONH}_4$, $\text{NH}_2 - \text{CO} - \text{NH}_2$
- NH_2COOH , $\text{NH}_2 - \text{CO} - \text{NH}_2$
- $(\text{NH}_4)_2\text{CO}_3$, $\text{NH}_2 - \text{CO} - \text{NH}_2$

More than One Options Correct Type

20. Which of the following statements are correct?
- Fullerenes have dangling bonds.
 - Fullerenes are cage-like molecules.
 - Graphite is thermodynamically most stable allotrope of carbon.
 - Graphite is slippery and hard and therefore used as a dry lubricant in machines.

21. In the following reaction,
 $2X + \text{B}_2\text{H}_6 \longrightarrow [\text{BH}_2(\text{X})_2]^+ [\text{BH}_4]^-$
X can be

Your favourite MTG Books/Magazines available in UTTARAKHAND at

- National Book House - Dehradun Ph: 0135-2659430; Mob: 9897830283
- Army Book Depot - Dehradun Ph: 0135-2756683; Mob: 9897927902
- Om Vidya Educational Book - Dehradun Mob: 9897833882
- Career Zone - Haldwani Ph: 05946-262051; Mob: 9412128075, 94123833167
- World Vision Publication - Haldwani Mob: 9927932200, 9027240169
- Diamond Stationers - Haridwar Ph: 0133-4252043; Mob: 9358398035, 9359763348
- Consul Book Depot - Nainital Ph: 0592-235164; Mob: 9412084105
- Indra Book Emporium - Roorkee Ph: 0133-276105; Mob: 8126293314
- Cambridge Book Depot - Roorkee Ph: 272341, 272345; Mob: 9719190955

Visit "MTG IN YOUR CITY" on www.mtg.in to locate nearest book seller OR write to info@mtg.in OR call

0124-6601200 for further assistance.

- (a) NH_3 (b) CH_3NH_2
 (c) $(\text{CH}_3)_2\text{NH}$ (d) $(\text{CH}_3)_3\text{N}$

22. Select the correct statements.

- (a) Borax is used as a buffer.
 (b) 1 M borax solution reacts with equal volumes of 2 M HCl solution.
 (c) A solution of borax can be titrated with a mineral acid using methyl orange as indicator.
 (d) Coloured bead obtained in borax bead test contains metaborate.

23. Select the correct statements about silicates.

- (a) Cyclic silicate having three Si atoms contains six Si — O — Si linkages.
 (b) Chain silicates contain $(\text{SiO}_3)_n^{2n-}$ ions which are formed by linking SiO_4^{4-} units linearly.
 (c) $(\text{Si}_2\text{O}_5)_n^{2n-}$ is the formula of double chain silicate.
 (d) SiO_4^{4-} units polymerize to form silicate because Si atom has less tendency to form π -bond with oxygen.

Integer Answer Type

24. How many moles of NO_2 are produced when one mole of B reacts with HNO_3 ?
 25. The coordination number of Al in the crystalline state of AlCl_3 is
 26. How many silicon atoms are present in the anion of a pyrosilicate?

Comprehension Type

Boron is the first member of group 13 of the periodic table and is the only non-metal of this group. It has a small size, high ionization enthalpy and high electronegativity due to which it cannot lose its valence electrons to form B^{3+} ions. Instead, it can form three covalent bonds by sharing its three valence electrons. Consequently, boron compounds, especially the hydrides and halides are electron deficient and behave as Lewis acids.

27. The compound which is not a Lewis acid is

- (a) BF_3 (b) AlCl_3
 (c) $\text{BeCl}_2 \cdot 2\text{H}_2\text{O}$ (d) SnCl_4

28. If a molecule of MX_3 has zero dipole moment, the sigma bonding orbitals used by M (atomic number < 21) are

- (a) pure p (b) sp-hybrid
 (c) sp^2 -hybrid (d) sp^3 -hybrid.

Matrix Match Type

29. Match the Column I with Column II and choose the correct answer using the codes given below.

Column I	Column II
P. Does not show inert pair effect	1. InCl
Q. Shows $p\pi - p\pi$ back bonding	2. Al
R. Shows disproportionation reaction	3. $\text{N}(\text{SiH}_3)_3$
S. Shows $p\pi - d\pi$ back bonding	4. BF_3

P	Q	R	S
(a) 4	1	3	2
(b) 1	4	2	3
(c) 2	4	1	3
(d) 4	1	2	3

30. Match the species given in Column I with the properties mentioned in Column II and choose the correct answer using the codes given below.

Column I	Column II
(P) BF_4^-	1. Strong oxidising agent
(Q) AlCl_3	2. Lewis acid
(R) SnO	3. Can be further oxidised
(S) PbO_2	4. Tetrahedral shape

P	Q	R	S
(a) 4	1	4	2
(b) 4	2	3	1
(c) 1	4	2	3
(d) 3	2	1	4



Keys are published in this issue. Search now! ☺

SELF CHECK

No. of questions attempted
 No. of questions correct
 Marks scored in percentage


Check your score! If your score is

> 90%	EXCELLENT WORK !	You are well prepared to take the challenge of final exam.
90-75%	GOOD WORK !	You can score good in the final exam.
74-60%	SATISFACTORY !	You need to score more next time.
< 60%	NOT SATISFACTORY!	Revise thoroughly and strengthen your concepts.


A mechanistic approach to any reaction classifies the reaction according to mechanism rather than by functional group. It explains the stereochemistry involved in a particular reaction, which can either be regioselective, stereoselective or stereospecific.

I Hydroboration – Oxidation Reaction

Step - I : Hydroboration




Step - II : Oxidation




- Rate of formation of the C—B bond > Rate of formation of C—H bond.
- Formation of four-centred transition state.
- Hydroboration is regioselective.
- Hydroboration is a *syn*-addition across the alkene.
- In step-II, boron goes backward and forward between planar neutral structure and anionic tetrahedral structure.
- In step-II, cleavage of O—O single bond is the driving force.
- In step-I, new C—B bond and in step-II new C—O bond are formed.
- The net result of this reaction is addition of water across the double bond.

II Oxymercuration – Demercuration Reaction

Step - I : Oxymercuration




Step - II : Demercuration




- In step-I, *i.e.*, oxymercuration, —OH and Hg²⁺ are added across the alkene.
- Oxymercuration is regioselective and stereospecific.
- Stereochemically, oxymercuration is an *anti*-addition.
- In step-I, water attacks at the more substituted end of the mercurinium ion (transition state).
- Driving force for demercuration is a weak C—Hg bond.
- To replace Hg with H, NaBH₄, a reducing agent is used.
- Oxymercuration-reduction is a popular laboratory technique with Markovnikov selectivity while avoiding carbocation intermediate and thus, rearrangement which can lead to complex product mixture.

III Friedel–Crafts Alkylation Reaction

Step - I : Formation of Intermediate (Carbocation)



Step - II : Reaction with Electrophile (Alkylation)



- This reaction follows S_N1 pathway.
- Species having capability to form carbocation are used.
- Carbocation can be generated by
 - protonation of an alkene.
 - the acid-catalysed decomposition of a tertiary alcohol.
 - Lewis acid catalysed decomposition of a *tert*-alkyl chloride.
- Carbocation rearrangement (to more stable carbocation) takes place whenever possible.
- Friedel–Crafts alkylation with alkyl halides proceeds via a carbocation and chiral alkyl halides are expected to give racemic arene products. The extent of racemisation depends on the Lewis acid and the reaction conditions.

The f-Block Elements

These elements are also called rare earth metals due to their extremely low natural occurrence. Mostly, these are prepared artificially in laboratory. All the actinoids are radioactive hence, can be used in nuclear power plants or as weapons.

Lanthanoids (4f-series)

La	57	[Xe] 5d ¹ 6s ²	Gd	64	[Xe] 4f ⁷ 5d ¹ 6s ²
Ce	58	[Xe] 4f ¹ 5d ¹ 6s ²	Tb	65	[Xe] 4f ⁹ 6s ²
Pr	59	[Xe] 4f ³ 6s ²	Dy	66	[Xe] 4f ¹⁰ 6s ²
Nd	60	[Xe] 4f ⁴ 6s ²	Ho	67	[Xe] 4f ¹¹ 6s ²
Pm	61	[Xe] 4f ⁵ 6s ²	Er	68	[Xe] 4f ¹² 6s ²
Sm	62	[Xe] 4f ⁶ 6s ²	Tm	69	[Xe] 4f ¹³ 6s ²
Eu	63	[Xe] 4f ⁷ 6s ²	Yb	70	[Xe] 4f ¹⁴ 6s ²
			Lu	71	[Xe] 4f ¹⁴ 5d ¹ 6s ²

Oxidation States

+3, *e.g.*, Gd³⁺, Lu³⁺ and +2 and +4. *e.g.*, Eu²⁺, Ce⁴⁺
Lanthanoids show limited number of oxidation states because the energy gap between 4*f*- and 5*d*-subshells is large.

Lanthanoid Contraction

The steady decrease in the atomic/ionic radii of lanthanoids with increasing atomic no. is called *lanthanoid contraction*.

Cause of Lanthanoid Contraction

As the atomic number increases in lanthanoid series, the new electron goes to fill 4*f*-orbitals. The nuclear charge, however, increases by one at each step. Thus, there is a gradual increase in the effective nuclear charge experienced by the outer electrons. Consequently, the attraction of the nucleus for the electrons in the outermost shell increases which results in gradual decrease in size of lanthanoids with increase in atomic number.

Consequences of Lanthanoid Contraction

- Similarity in properties of second and third transition series elements.
- Due to lanthanoid contraction, the differences in physical properties enable the separation of individual lanthanoid elements by ion exchange method.
- Due to lanthanoid contraction the basic strength decreases from La(OH)₃ to Lu(OH)₃.

Magnetic Properties

All the lanthanoid ions except La³⁺, Lu³⁺, Ce⁴⁺ and Yb²⁺ are paramagnetic. Magnetic moment ($\mu = \sqrt{4S(S+1) + L(L+1)}$)

Uses

- Used for the production of alloys *e.g.*, Mischmetal, which contains lanthanoid metals ($\approx 95\%$) and iron ($\approx 5\%$).
- La₂O₃ is used in Crooke's lenses.

Actinoids (5f-series)

Ac	89	[Rn] 6d ¹ 7s ²	Cm	96	[Rn] 5f ⁷ 6d ¹ 7s ²
Th	90	[Rn] 6d ² 7s ²	Bk	97	[Rn] 5f ⁹ 7s ²
Pa	91	[Rn] 5f ² 6d ¹ 7s ²	Cf	98	[Rn] 5f ¹⁰ 7s ²
U	92	[Rn] 5f ³ 6d ¹ 7s ²	Es	99	[Rn] 5f ¹¹ 7s ²
Np	93	[Rn] 5f ⁴ 6d ¹ 7s ²	Fm	100	[Rn] 5f ¹² 7s ²
Pu	94	[Rn] 5f ⁶ 7s ²	Md	101	[Rn] 5f ¹³ 7s ²
Am	95	[Rn] 5f ⁷ 7s ²	No	102	[Rn] 5f ¹⁴ 7s ²
			Lr	103	[Rn] 5f ¹⁴ 6d ¹ 7s ²

Oxidation States

- General oxidation state is +3.
- Besides +3 state, other possible oxidation states are +2, +4, +5, +6 and +7.
- Actinoids show a large number of oxidation states because of very small energy gap between 5*f*, 6*d* and 7*s* subshells.

Actinoid Contraction

The steady decrease in ionic radii with increase in atomic number is referred to as *actinoid contraction*. The contraction is greater from element to element in actinoid series as compared to lanthanoid contraction. This is due to poor shielding of 5*f*-electrons.

Colour of the Ions

The actinoid cations are generally coloured due to *f-f* transition, *e.g.*, U³⁺ (5*f*³) : Red, Np³⁺ (5*f*⁴) : Purple, Pu³⁺ (5*f*⁵) : Violet, Am³⁺ (5*f*⁶) : Pink, U⁴⁺ (5*f*²) : Pink.

Magnetic Properties

The actinoid elements or ions are paramagnetic due to the presence of unpaired electrons. However, the magnetic properties of actinoids are higher than lanthanoids.

Uses

- Thorium is used in atomic reactors as fuel rods and in the treatment of cancer.
- Uranium is used as nuclear fuel. Its salts are used in glass industry (for imparting green colour), textile industry and also in medicines.
- Plutonium is used as fuel for atomic reactors as well as in atomic bombs.

NEET | JEE

ESSENTIALS

Class
XII

Maximise your chance of success, and high rank in NEET, JEE (Main and Advanced) by reading this column. This specially designed column is updated year after year by a panel of highly qualified teaching experts well-tuned to the requirements of these Entrance Tests.


Unit 6

Aldehydes and Ketones | Carboxylic Acids

ALDEHYDES AND KETONES



INTRODUCTION

- Organic compounds containing carbonyl group (>C=O) are known as carbonyl compounds. Both aldehydes and ketones come under carbonyl compounds, having general formula, $\text{C}_n\text{H}_{2n}\text{O}$.




NOMENCLATURE

- Common System :** In the common system, aldehydes are named according to the name of the corresponding carboxylic acid to which they are oxidised. Suffix '-ic acid' is replaced by 'aldehyde'.
Acetic acid $\xrightarrow[-\text{ic acid}]{+\text{aldehyde}}$ Acetaldehyde
In the common system, ketones (aliphatic or aromatic) are named by using the names of alkyl or aryl groups (in the alphabetical order) present in the molecule by adding the suffix 'ketone'.
- IUPAC System :** The name of aldehyde is derived by replacing the terminal '-e' of the corresponding alkane by '-al'.
In ketones, the name of the individual members are derived by replacing the terminal 'e' of the corresponding alkane by 'one'. Thus, the ketones are known as *alkanones*.


Formula	Common Name	IUPAC Name
HCHO	Formaldehyde	Methanal
CH ₃ CHO	Acetaldehyde	Ethanal
$\begin{array}{c} \text{3} \quad \text{2} \quad \text{1} \\ \text{CH}_3\text{CHCHO} \\ \\ \text{CH}_3 \end{array}$	Iso-butyraldehyde	2-Methyl propanal
$\text{CH}_2=\overset{2}{\text{CH}}-\overset{1}{\text{CHO}}$	Acrolein	Prop-2-enal
	Benzene carb-aldehyde	Benzaldehyde
	Phthaldehyde	Benzene-1,2-dicarbaldehyde
CH ₃ COCH ₃	Dimethyl ketone or acetone	Propanone
$\overset{1}{\text{CH}_3}\overset{2}{\text{CO}}\overset{3}{\text{CH}_2}\overset{4}{\text{CH}_2}\overset{5}{\text{CH}_3}$	Methyl n-propyl ketone	Pentan-2-one
$\begin{array}{c} \text{1} \quad \text{2} \quad \text{3} \quad \text{4} \quad \text{5} \\ \text{CH}_3\text{CCH}_2\text{CCH}_3 \\ \quad \\ \text{O} \quad \text{O} \end{array}$	Acetylacetone	Pentane-2,4-dione

ELECTRONIC STRUCTURE OF CARBONYL GROUP

- The carbonyl carbon is sp^2 hybridised thus, joined with three atoms by three sigma bonds. The fourth valence electron of carbon remains in its p -orbital and forms a π -bond by overlapping with p -orbital of oxygen. Since, these bonds utilise sp^2 -orbitals, they lie in the same plane, having planar structure and these bonds are 120° apart to one another.





Due to electronegativity difference of carbon and oxygen, carbonyl group is polar.





PREPARATION


- From alcohols :**







- From carboxylic acids :**





- From alkenes :**



- From alkynes :**




- From gem-dihalides :**




INFOSHOTS

Photoinduced annulation of 1-aryl-2-(furan/thiophen-2-yl)butane-1,3-diones

A photoinduced direct oxidative annulation of 1-aryl-2-(furan/thiophen-2-yl)butane-1,3-diones and ethyl-2-(furan-2-yl)-3-oxo-3-(aryl-2-yl)propanoates in EtOH without the need for any transition metals and oxidants provided access to highly functionalized polyheterocyclic 1-(5-hydroxynaphtho[2,1-b]furan-4-yl)ethanones and 1-(5-hydroxyfuran/thieno/pyrrolo[3,2-e]benzofuran-4-yl)ethanones. The phenomenon of excited-state intramolecular proton transfer (ESIPT) was observed for both 1-(5-hydroxynaphtho[2,1-b]furan-4-yl)ethanone and 1-(5-hydroxyfuran/thieno/pyrrolo[3,2-e]benzofuran-4-yl)ethanone analogues.



• From Grignard Reagents :



Methods of Preparation

For aldehydes

$$\text{R-C}\equiv\text{N} + 2[\text{H}] \xrightarrow{\text{SnCl}_2/\text{HCl}} \text{R-CH=NH}\cdot\text{HCl} \xrightarrow[\text{H}_2\text{O}]{\text{Boiling}} \text{RCHO} + \text{NH}_4\text{Cl}$$

Alkyl cyanides (Stephen reduction)

$$\text{R-CH=CH}_2 + \text{CO} + \text{H}_2 \xrightarrow[373 \text{ K, Pressure}]{[\text{Co}(\text{CO})_4]_2} \text{R-CH}_2\text{-CH}_2\text{-CHO}$$

Alkene

$$\text{R-C(=O)-Cl} + \text{H}_2 \xrightarrow[\text{Boiling xylene}]{\text{Pd/BaSO}_4, \text{S}} \text{R-C(=O)-H}$$

Acid chloride (Rosenmund reduction)

For ketones

$$\begin{array}{c} \text{R} \\ \diagdown \\ \text{CHOH} \\ \diagup \\ \text{R} \end{array} + \begin{array}{c} \text{CH}_3 \\ \diagdown \\ \text{C=O} \\ \diagup \\ \text{CH}_3 \end{array} \xrightarrow{[(\text{CH}_3)_3\text{CO}]_3\text{Al}} \begin{array}{c} \text{R} \\ \diagdown \\ \text{C=O} \\ \diagup \\ \text{R} \end{array} + \begin{array}{c} \text{CH}_3 \\ \diagdown \\ \text{CH-OH} \\ \diagup \\ \text{CH}_3 \end{array}$$


sec-Alcohol + Acetone (Excess) → Ketone + Propyl alcohol (Oppenauer Oxidation)

$$\text{CH}_3\text{-C(=O)-CH}_2\text{-COOC}_2\text{H}_5 \xrightarrow[\Delta]{\text{H}_2\text{O/H}^+} \text{CH}_3\text{-C(=O)-CH}_3 + \text{CO}_2 + \text{C}_2\text{H}_5\text{OH}$$


Acetoacetic ester

PREPARATION OF AROMATIC ALDEHYDES AND KETONES


• Commercial Method :




• Etard Reaction :




• Gattermann-Koch Reaction :




• Reimer-Tiemann Reaction







• Friedel-Crafts Acylation :



• Fries Rearrangement :



Presenting India's No.1 NEET Guides



MTG's Complete NEET Guides are India's best selling PMT books!! Rich in theoretical knowledge with a vast question bank comprising a wide variety of problems and exercises, these guidebooks ensure students are ready to compete in the toughest of medical entrance tests. 100% NCERT based, the guidebooks have been updated to match the syllabus and the exam pattern for medical entrance exams. No wonder these guidebooks emerged as the bestsellers in a short period of time.

HIGHLIGHTS:

- 100% NCERT based
- Comprehensive Chapterwise theory complemented with concept maps, flowcharts and easy-to-understand illustrations
- Last 10 years' questions (2008-2017) of AIPMT/NEET
- Chapterwise Topicwise MCQs with detailed explanations and solutions
- NEET 2017 Solved Paper included
- Over 50% of questions that appeared in NEET 2017 were from MTG's Complete NEET Guides



Scan now with your smartphone or tablet*



Available at all leading book shops throughout India.
For more information or for help in placing your order:
Call 0124-6601200 or e-mail:info@mtg.in

*Application to read QR codes required

Visit
www.mtg.in
for latest offers
and to buy
online!

DESIGN YOUR OWN DEGREE.

CHOOSE YOUR OWN CURRICULUM.

DETERMINE YOUR OWN CAREER.

Introducing Inter Disciplinary Experiential Active Learning (IDEAL)



IDEAL gives you the opportunity to select a discipline that will give you a career edge, and also study something you are passionate about. With IDEAL, you can now pursue a Specialization as a part of your Major degree or a Minor in another discipline, earning you extra credits at no additional cost. For instance, you can couple your Major in Mechanical Engineering with a Minor in Finance, or specialize in Artificial Intelligence while majoring in Computer Science. Chart your IDEAL path enriched further by industry internships and technology-enabled active learning that will help you in getting jobs. Begin your journey with SRMJEEE.

Take the SRMJEEE (B.Tech) 2018.
Online applications: Nov. 1, 2017 onwards.

ONLINE EXAM DATES

APRIL 16-30, 2018

TO APPLY

Visit: applications.srmuniv.ac.in
Make e-payment OR send the filled application form with DD for ₹1100. For eligibility and fee details, visit the respective websites.

FOR QUERIES

Call: +91 44 2745 5510
+91 44 4743 7500
Email: admissions.india@srmuniv.ac.in



SRM
UNIVERSITY
(Under Section 3 of UGC Act 1956)
www.srmuniv.ac.in

KATTANKULATHUR
RAMAPURAM
DELHI-NCR GHAZIABAD
VADAPALANI










SRM
UNIVERSITY
— AP, Amaravati
www.srmmap.edu.in



SRM
UNIVERSITY
— Delhi-NCR, Sonapat
www.srmuniversity.ac.in



CBSE CHAMPION Chapterwise -Topicwise Solved Papers



CBSE CHAMPION Chapterwise -Topicwise Solved Papers Series contains topicwise questions and solutions asked over last decade in CBSE-Board examination.

Questions are supported with topicwise graphical analysis of previous years CBSE Board questions as well as comprehensive and lucid theory. The questions in each topic have been arranged in descending order as per their marking scheme. Questions from Delhi, All India, Foreign and Compartment papers are included. This ensures that all types of questions that are necessary for Board exam preparation have been covered.

Important feature of these books is that the solutions to all the questions have been given according to CBSE marking scheme. CBSE sample paper and practice papers are also supplemented.

Examination papers for Class-10 and 12 Boards are based on a certain pattern. To excel, studying right is therefore more important than studying hard, which is why we created this series.



Available at all leading book shops throughout India.
For more information or for help in placing your order:
Call 0124-6601200 or email info@mtg.in


PHYSICAL PROPERTIES

- Due to intermolecular dipole-dipole interactions, they have higher b.pt. than non-polar compounds and weakly polar compounds. Among isomeric aldehydes and ketones, ketones have slightly higher boiling points.
- Lower aldehydes and ketones are soluble in water due to hydrogen bonding with water. But solubility decreases as the size of hydrocarbon part increases.

CHEMICAL PROPERTIES


Nucleophilic Addition Reactions

- In carbonyl group, due to high electronegativity of oxygen, carbon atom acquires a small positive charge and behaves as an electrophile. Therefore, the positively charged carbon is readily attacked by a nucleophile. However, during addition reaction, nucleophilic attack precedes the electrophilic attack because the anion produced is more stable than the cation.




- **Relative reactivities of aldehydes and ketones:** Aldehydes are more reactive than ketones in nucleophilic addition reactions. This is because alkyl groups are electron donating (+I-effect) and

Oxidation and Reduction Reactions




thus, increase the electron density on the carbonyl carbon in ketones. Also, in ketones, the steric effect of alkyl groups decreases the reactivity. Reactivity order decreases as the steric hindrance and +I-effect of alkyl group increases.

$\text{CH}_2\text{O} > \text{CH}_3\text{CHO} > (\text{CH}_3)_2\text{CO} > [(\text{CH}_3)_2\text{CH}]_2\text{CO}$
 Due to +R effect of benzene ring, aromatic aldehydes and ketones are less reactive than the corresponding aliphatic aldehydes and ketones. The decreasing order of reactivity is :
 $\text{C}_6\text{H}_5\text{CHO} > \text{C}_6\text{H}_5\text{COCH}_3 > \text{C}_6\text{H}_5\text{COC}_6\text{H}_5$






Nucleophilic Addition - Elimination Reactions





Miscellaneous Reactions


- Aldol condensation** : Aldehydes and ketones containing at least one α -hydrogen atom undergo self condensation in presence of dilute alkali to form aldol or ketol respectively.




- Cannizzaro reaction** : Aldehydes which do not contain an α -hydrogen atom undergo disproportionation when treated with concentrated alkali solution.




- Cross Cannizzaro reaction** :




Mechanism of Aldol Condensation




- Intramolecular Cannizzaro reaction** :



- Tischenko reaction** :




• **Polymerisation reactions :**



Substitution Reactions of Aromatic Aldehydes and Ketones

- Aldehydic and ketonic groups are electron withdrawing, therefore, they are deactivating and *m*-directing.



Distinction tests for aldehydes and ketones

Test	Aldehydes	Ketones
1. With Tollens' reagent	Form silver mirror	No reaction
2. With Fehling's solution	Give red precipitate (Aliphatic aldehyde)	No reaction
3. With Schiff's reagent	Give pink colour	No reaction
4. Reduction with LiAlH ₄	Primary alcohols are formed	Secondary alcohols are formed.
5. Action with alcohol in the presence of dry HCl.	Form acetals easily	Do not form ketals easily.

Uses

- 40% solution of formaldehyde in water is called formalin. It is used for preserving biological specimens.
- Acetaldehyde is used in the manufacturing of a number of organic compounds such as acetic acid, ethyl acetate, etc.
- Benzaldehyde is used as a starting material for the synthesis of organic compounds such as cinnamic acid, benzoyl chloride, etc.
- Acetone is extensively used as a solvent in industries and laboratories.
- Acetone is used in the preparation of chloroform, acetic anhydride. It is also one of the constituents of liquid nail polish.

Your favourite MTG Books/Magazines available in GOA at

- Golden Heart Emporium - Goa
Ph: 0832-2725208, 3257383, 2730874; Mob: 8322725208, 9370273479
- Universal Traders - Goa Ph: 0832-2315985; Mob: 9404150150
- Success Stationers - Margao Mob: 9850398314


Visit "MTG IN YOUR CITY" on www.mtg.in to locate nearest book seller OR write to info@mtg.in OR call

0124-6601200 for further assistance.

CARBOXYLIC ACIDS



INTRODUCTION

- Organic compounds containing $-\text{COOH}$ as a functional group are called carboxylic acids. A large number of carboxylic acids are found naturally. Some higher members of aliphatic carboxylic acids (C_{12} - C_{18}) are known as fatty acids because these are obtained by the hydrolysis of oils and fats.




NOMENCLATURE

- The common names of carboxylic acids are based on their source of origin.
- In IUPAC system, the name of the acids is derived from the corresponding alkane by replacing the terminal 'e' by suffix '-oic acid'.

Monocarboxylic acid	Common Name	IUPAC Name
HCOOH	Formic acid	Methanoic acid
CH_3COOH	Acetic acid	Ethanoic acid
$\begin{array}{c} 3 & 2 & 1 \\ \text{CH}_3 & \text{CH} & \text{COOH} \\ & & \\ & \text{CH}_3 & \end{array}$	<i>Iso</i> -butyric acid (α -Methyl propionic acid)	2-Methyl propanoic acid
$\begin{array}{c} 5 & 4 & 3 & 2 & 1 \\ \text{CH}_3 & \text{CH}_2 & \text{CH}_2 & \text{CH}_2 & \text{COOH} \end{array}$	Valeric acid	Pentanoic acid
$\begin{array}{c} 3 & 2 & 1 \\ \text{CH}_3 & -\text{CH} & -\text{COOH} \\ & & \\ & \text{OH} & \end{array}$	α -Hydroxy-propionic acid or Lactic acid	2-Hydroxy-propanoic acid
$\begin{array}{c} 3 & 2 & 1 \\ \text{CH}_2 = & \text{CH} & \text{COOH} \end{array}$	Acrylic acid	Prop-2-enoic acid
	Benzoic acid	Benzene-carboxylic acid or benzoic acid
	Salicylic acid	2-Hydroxybenzoic acid

ELECTRONIC STRUCTURE OF CARBOXYL GROUP

- In carboxylic acids, the bonds to the carboxyl carbon lie in one plane and are separated by about 120° .
- The carboxyl carbon is less electrophilic than carbonyl carbon because of the possible resonance structures.



PREPARATION

- $$\text{RMgX} \xrightarrow[\text{(ii) H}_2\text{O/H}^+]{\text{(i) CO}_2/\text{dry ether}} \text{RCOOH}$$

(Carbonylation of Grignard reagent)
- $$\text{RCH}_2\text{OH} \xrightarrow[\text{[O]}]{\text{K}_2\text{Cr}_2\text{O}_7/\text{H}^+} \text{RCOOH}$$

(Oxidation of 1° alcohol)
- $$\text{RCOCl} \xrightarrow{\text{H}_2\text{O}} \text{RCOOH}$$

(Hydrolysis of acid chlorides)
- $$\text{RCONH}_2 \xrightarrow[\Delta]{\text{H}_2\text{O/H}^+ \text{ or } \text{OH}^-} \text{RCOOH} + \text{NH}_3$$

(Hydrolysis of acidamides)

$$\text{RCONH}_2 + \text{HNO}_2 \longrightarrow \text{RCOOH} + \text{N}_2 + \text{H}_2\text{O}$$
- $$\text{R}-\text{C}\equiv\text{N} \xrightarrow[\Delta]{2\text{H}_2\text{O/HCl (dil.)}} \text{RCOOH} + \text{NH}_4\text{Cl}$$

(Hydrolysis of alkyl cyanides)
- $$\text{H}_2\text{C}=\text{CH}_2 + \text{CO} + \text{H}_2\text{O} \xrightarrow[\text{under pressure}]{\text{H}_3\text{PO}_4, 623 \text{ K}} \text{H}_3\text{CCH}_2\text{COOH}$$

(Koch reaction)
- $$\text{CH}_3-\overset{\text{O}}{\parallel}{\text{C}}-\text{R} \xrightarrow{\text{NaOI}} \text{CHI}_3 + \text{RCOONa}$$

(Haloform reaction)

$$\text{CH}_3\text{CHOH}-\text{R} \xrightarrow{\text{NaOI}} \text{CHI}_3 + \text{RCOONa}$$

(Haloform reaction)
- $$\begin{array}{c} \text{COOH} \\ | \\ \text{COOH} \end{array} \xrightarrow{\Delta} \text{HCOOH} + \text{CO}_2$$



(Thermal decomposition of α -dioic acids)

$$\text{CH}_2(\text{COOH})_2 \xrightarrow{\Delta} \text{CH}_3\text{COOH} + \text{CO}_2$$

Malonic acid
- $$\text{R}-\text{CCl}_3 + 4\text{KOH} \longrightarrow \text{RCOOK} + 3\text{KCl} + \text{H}_2\text{O}$$

(Hydrolysis of 1,1,1-trihalogen derivative of alkane)

- Aromatic acids are obtained by the oxidation of alkyl benzenes.




PHYSICAL PROPERTIES

- Smell :** C₁ – C₃ aliphatic acids have pungent smell. C₄ – C₉ have unpleasant smell. Butyric acid has a smell of rancid butter. Higher aliphatic and aromatic acids have no smell.
- State :** C₁ – C₉ are colourless liquids. The higher acids are colourless waxy solids. Most of the aromatic acids exist as colourless crystalline solids.
- Miscibility :** C₁ – C₄ are freely miscible with water due to intermolecular hydrogen bonding. Solubility of higher members decreases due to increased hydrophobic interactions of the hydrocarbon part.
- Boiling points :** Monocarboxylic acids have higher boiling points as compared to aldehydes, ketones and even alcohols of comparable molecular masses due to the presence of greater intermolecular hydrogen bonding.
- Melting points :** Acids containing even number of carbon atoms have higher melting points as compared to their next lower and higher homologues which contain odd number of carbon atoms.

ACIDIC STRENGTH

- Carboxylic acids ionise to give proton in aqueous solution.




K_a is dissociation constant of the acid which varies with temperature.

- The acid strength is also measured in terms of $\text{p}K_a$ values (negative logarithm of K_a).


$$\text{p}K_a = -\log K_a$$

i.e., more the value of $\text{p}K_a$, less acidic is the carboxylic acid indicating the lower value of K_a and smaller the value of $\text{p}K_a$, more acidic is the carboxylic acid indicating the higher value of K_a .

- The acidic nature can also be explained on the basis of resonating structures of carboxylate ion.






- By resonating structure (B), it is clear that proton is removed to give carboxylate ion as there is positive charged oxygen atom of O–H group.




EFFECT OF SUBSTITUENTS ON ACIDITY

- If electron withdrawing group is present then there will be dispersal of negative charge of the carboxylate ion as a result it will be more stable than the acid not having electron withdrawing groups.
- Acidity decreases as the number of electron withdrawing group decreases.
i.e., $\text{Cl}_3\text{CCOOH} > \text{Cl}_2\text{CHCOOH} > \text{ClCH}_2\text{COOH} > \text{CH}_3\text{COOH}$
- The acidic strength mainly depends upon inductive effect. If +I-effect of alkyl group increases, acidity decreases.
 $\text{HCOOH} > \text{CH}_3\text{COOH} > (\text{CH}_3)_2\text{CHCOOH} > (\text{CH}_3)_3\text{CCOOH}$
- If –I-effect of halogen decreases, acidity decreases.
 $\text{FCH}_2\text{COOH} > \text{ClCH}_2\text{COOH} > \text{BrCH}_2\text{COOH} > \text{ICH}_2\text{COOH}$
- Acidity decreases as the distance of the halogen atom from carboxyl group increases.
 $-\text{CHCOOH} > -\text{CHCH}_2\text{COOH} > -\text{CHCH}_2\text{CH}_2\text{COOH}$




CHEMICAL REACTIONS


- Reactions involving –OH group :




- Reactions involving proton of —OH group :




- Reaction involving —COOH group :



- Reaction involving —R group :



- Ring substitution in aromatic acids : —COOH group is deactivating and *meta*-directing.



DISTINCTION BETWEEN ALCOHOLS, PHENOLS AND CARBOXYLIC ACIDS

Test	Alcohols	Phenols	Carboxylic acids
1. Action on litmus	No reaction	Turn blue litmus red	Turn blue litmus red
2. Reaction with sodium bicarbonate (NaHCO ₃)	No reaction	No reaction	Give effervescence due to the evolution of CO ₂
3. Reaction with sodium hydroxide (NaOH)	No reaction	Form salts	Form salts
4. Reaction with FeCl ₃	No reaction	Give characteristic colours violet, blue, red, etc.	Carboxylic acids give coloured precipitates e.g., CH ₃ COOH gives buff coloured ppt.

USES


- Methanoic acid is used in leather tanning and as a coagulating agent for rubber latex in rubber industry.

- Methanoic acid is used as a medicine for gout and neuritis.
- Ethanoic acid is used in the manufacture of rayon, plastics, rubber and in silk industries.
- Ethanoic acid is used as vinegar in food industry.
- Esters of benzoic acids are used in perfumery.

INFOSHOTS

Palladium-catalyzed diastereoselective synthesis of 3-arylbutanoic acid derivatives


The first palladium-catalyzed diastereoselective conjugate addition of arylboronic acids to chiral imides is reported. The catalytic system employing 4-*tert*-butyloxazolidin-2-one as the chiral auxiliary in a mixed solvent system of MeOH/H₂O (1:3) under an air atmosphere provides the optically active 3-arylbutanoic acid derivatives in excellent yields with high diastereoselectivity.



SPEED PRACTICE

1. The reaction of a Grignard reagent with a carboxylic acid does not give a secondary alcohol. This is because
- Grignard reagents only react with aldehydes, ketones, esters and epoxides
 - the carboxylic acid is too sterically hindered to react
 - the carboxylic acid is not electrophilic enough to react
 - Grignard reagent is a base, so an acid-base reaction occurs.


2. Identify A in the given reaction,
- $$\text{CH}_3\text{C}(=\text{O})\text{CH}_2\text{CH}_2\text{CH}_2\text{Cl} \xrightarrow{\text{CH}_3\text{MgBr}} \text{A}$$




3. Consider the reactions,
- $$\text{C}_2\text{H}_6\text{O} \xrightarrow[573\text{ K}]{\text{Cu}} \text{A} \xrightarrow[\text{Silver mirror observed}]{[\text{Ag}(\text{NH}_3)_2]^+}$$
- $$\text{A} \xrightarrow[-\text{OH}, \Delta]{\text{NH}_2\text{NHCONH}_2} \text{Z}$$
- $$\text{A} \xrightarrow[-\text{OH}, \Delta]{\text{NH}_2\text{NHCONH}_2} \text{Y}$$

Identify A, X, Y and Z.


- A-Methoxymethane, X-Ethanol, Y-Ethanoic acid, Z-Semicarbazide
 - A-Ethanal, X-Ethanol, Y-But-2-enal, Z-Semicarbazone
 - A-Ethanol, X-Acetaldehyde, Y-Butanone, Z-Hydrazone
 - A-Methoxymethane, X-Ethanoic acid, Y-Acetate ion, Z-Hydrazine (NEET 2017)
4. Structure of $\text{C}_8\text{H}_8\text{Cl}_2$, which on aqueous alkali hydrolysis gives a product which does not give positive iodoform test, but gives silver mirror test, is




5. In the given reaction,
-
- The product (A) is



6. Enolisation is maximum in case of




7. Consider the following reaction sequence,



The correct statement regarding X is that

- it has an aldehyde functional group
- it has both aldehyde and ketone functional groups
- it has a ketone functional group
- it has a ketone and an acid functional group.

8. The correct sequence of reagents for the following conversion will be





- (a) CH_3MgBr , $[\text{Ag}(\text{NH}_3)_2]^+\text{OH}^-$, $\text{H}^+/\text{CH}_3\text{OH}$
 (b) $[\text{Ag}(\text{NH}_3)_2]^+\text{OH}^-$, CH_3MgBr , $\text{H}^+/\text{CH}_3\text{OH}$
 (c) $[\text{Ag}(\text{NH}_3)_2]^+\text{OH}^-$, $\text{H}^+/\text{CH}_3\text{OH}$, CH_3MgBr
 (d) CH_3MgBr , $\text{H}^+/\text{CH}_3\text{OH}$, $[\text{Ag}(\text{NH}_3)_2]^+\text{OH}^-$

(JEE Main 2017)



9. The IUPAC name of $\text{CH}_3-\overset{\text{O}}{\parallel}{\text{C}}-\text{CH}_2-\text{CH}_2-\text{CHO}$ is

- (a) 2-oxopentanal (b) 4-oxopentanal
 (c) 4-formylbutan-2-one
 (d) 5-formylbutan-2-one.

10. In the given sequence of reactions,




A and B are




- (c) both (a) and (b)
 (d) none of the above is correct.

11. The correct statement(s) about the following reactions sequence is(are)




- (a) R is steam volatile
 (b) Q gives dark violet colouration with 1% aqueous FeCl_3 solution
 (c) S gives yellow precipitate with 2, 4-dinitrophenylhydrazine
 (d) S gives dark violet colouration with 1% aqueous FeCl_3 solution. [JEE Advanced 2017]

12. Among the following four hydrogens which one is most acidic?




- (a) (i) (b) (ii) (c) (iii) (d) (iv)

13. The compounds A, B and C in the reaction sequence,




- (a) iodoform, ethylene, ethyl alcohol
 (b) iodoform, acetylene, acetaldehyde
 (c) iodoform, propyne, acetone
 (d) iodoform, 2-propanol, propanone.

14. The correct statement about the following reaction sequence is



- (a) 'R' gives an aldol condensation reaction on heating with NaOH solution
 (b) step '4' is an aromatic nucleophilic substitution reaction
 (c) the compound 'Q' gives a yellow precipitate in acetone
 (d) the end product is a mixture of three compounds.


15. The correct order of strength of following carboxylic acids is



- (a) I > II > III (b) II > III > I
 (c) III > II > I (d) II > I > III


(NEET Phase-II 2016)

16. Identify the product(s).




- (a) $\begin{array}{c} \text{COONa} \\ | \\ \text{COONa} \end{array}$ (b) $\begin{array}{c} \text{CH}_2\text{OH} \\ | \\ \text{COONa} \end{array}$
 (c) $\begin{array}{c} \text{CHO} \\ | \\ \text{COONa} \end{array}$ (d) both (a) and (b)


17. In the following reaction,



the product formed is





18. Choose the correct alternate for the given reaction sequence :



- | A | R ₁ | B | R ₂ |
|---|----------------|---|----------------|
| (a) CH ₃ CH ₂ NH ₂ | Schmidt | CH ₃ CH(Br)COOH | HVZ |
| (b) CH ₃ CH ₂ CONH ₂ | HVZ | CH ₃ CH ₂ COBr | Schmidt |
| (c) CH ₃ CH ₂ NH ₂ | HVZ | CH ₃ CH ₂ COBr | Schmidt |
| (d) CH ₃ CH ₂ CH ₂ NH ₂ | Schmidt | CH ₃ CH ₂ NH ₂ | HVZ |


19. What would be the major product in the following reaction?



20. Bouveault-Blanc reduction reaction involves

- (a) reduction of an acyl halide with H₂/Pd
 (b) reduction of an anhydride with LiAlH₄
 (c) reduction of an ester with Na/C₂H₅OH
 (d) reduction of a carbonyl compound with Na/Hg and HCl. (JEE Main Online 2016)

21. Correct statement for the given reaction is




- (a) X and Y are structural isomers

- (b) both X and Y gives only one monochlorinated product

(c) X on oxidative ozonolysis gives gluteric acid

(d) X and Y on reaction with H₂N-OH give oximes.

22. Acids with their pK_a values are given below,



Which group has maximum electron withdrawing tendency?

- (a) -NO₂ (b) -H (c) -CN (d) -OH

23. The correct statement regarding a carbonyl compound with a hydrogen atom on its alpha carbon, is

(a) a carbonyl compound with a hydrogen atom on its alpha-carbon rapidly equilibrates with its corresponding enol and this process is known as carbonylation


(b) a carbonyl compound with a hydrogen atom on its alpha-carbon rapidly equilibrates with its corresponding enol and this process is known as keto-enol tautomerism

(c) a carbonyl compound with a hydrogen atom on its alpha-carbon never equilibrates with its corresponding enol

(d) a carbonyl compound with a hydrogen atom on its alpha-carbon rapidly equilibrates with its corresponding enol and this process is known as aldehyde-ketone equilibration.

(NEET Phase-I 2016)

24. Consider the following aldol condensation reaction,





the nucleophile is

- (a) CH₂=CH-OH (b) CH₂=CH-O⁻

- (c) $\bar{\text{C}}\text{H}_2-\text{CHO}$ (d) CH₃-C(=O)-H

25. What is the final product (B) of the given reaction sequence?






26. In the given Cannizzaro reaction,


$$2\text{PhCHO} \xrightarrow{\text{OH}^-} \text{PhCH}_2\text{OH} + \text{PhCOO}^-$$
 the slowest step is
 (a) the attack of OH^- at the carbonyl group
 (b) the transfer of hydride to the carbonyl group
 (c) the abstraction of proton from carboxylic acid
 (d) the deprotonation of PhCH_2OH .
27. In the following species, the one which is likely to be the intermediate during benzoin condensation of benzaldehyde, is
 (a) $\text{Ph}-\text{C}\equiv\overset{+}{\text{O}}$ (b) $\text{Ph}-\overset{+}{\text{C}}\begin{matrix} \text{OH} \\ \text{CN} \end{matrix}$
 (c) $\text{Ph}-\overset{-}{\text{C}}\begin{matrix} \text{OH} \\ \text{CN} \end{matrix}$ (d) $\text{Ph}-\overset{-}{\text{C}}=\text{O}$

(JEE Advanced 2015)

28. Acetone molecules on condensation can not give
 (a) mesityl oxide (b) phorone
 (c) crotonaldehyde (d) mesitylene.
29. The suitable reagent for the following conversion is





- (a) LiAlH_4 (b) $\text{CH}_2=\text{O}/\text{OH}^-$
 (c) $\text{Na}/\text{C}_2\text{H}_5\text{OH}$ (d) H_2/Ni
30. Bromination of PhCOMe in acidic medium produces mainly




SOLUTIONS



1. (d)






3. (b)



Final product will give positive silver mirror test but not iodoform test.




6. (b)



9. (b)


10. (a)



'Q' is steam volatile due to intramolecular hydrogen bonding while 'R' undergoes intermolecular hydrogen bonding hence, has higher boiling point. 'Q' gives dark violet colouration with 1% aqueous FeCl₃ solution due to the presence of phenolic group while 'S' gives yellow precipitate with 2, 4-dinitrophenyl hydrazine due to the presence of aldehydic group (-CHO).



12. (b)

13. (b)



15. (b)


16. (d): The given reaction is Cannizzaro reaction hence, two products are obtained.




18. (a)

19. (a)

20. (c): Bouveault-Blanc reduction involves reduction of an ester in presence of Na/C₂H₅OH.



21. (b):




22. (a)

23. (b)

24. (a): In acid catalysed aldol condensation, enol acts as a nucleophile and can attack on the carbonyl carbon of other molecule.


25. (d)

26. (b)




28. (c)

29. (b):



30. (d): Reaction in acidic medium proceeds upto monobromination stage. ♦♦

EXAMINER'S MIND CLASS XII



The questions given in this column have been prepared strictly on the basis of NCERT Chemistry for Class XII. This year JEE (Main & Advanced)/NEET/AIIMS have drawn their papers heavily from NCERT books.



Section - I	Q. 1 to 10 Only One Option Correct Type MCQs.
Section - II	Q. 11 to 13 More than One Options Correct Type MCQs.
Section - III	Q. 14 to 17 Paragraph Type MCQs having Only One Option Correct.
Section - IV	Q. 18 & 19 Matching List Type MCQs having Only One Option Correct.
Section - V	Q. 20 to 22 Assertion Reason Type MCQs having Only One Option Correct. Mark the correct choice as : (a) If both assertion and reason are true and reason is the correct explanation of assertion. (b) If both assertion and reason are true but reason is not the correct explanation of assertion. (c) If assertion is true but reason is false. (d) If both assertion and reason are false.
Section - VI	Q. 23 to 25 Integer Value Correct Type Questions having Single Digit Integer Answer, ranging from 0 to 9 (both inclusive).

CHEMICAL KINETICS

SECTION - I


Only One Option Correct Type

1. For the first order decomposition reaction of N_2O_5 , it is observed that




which one of the following is correct?

- (a) $k = k'$ (b) $k = 2k'$
 (c) $k = k'/2$ (d) $k = k'^2$
2. The given figure depicts the change in concentration of species X and Y as a function of time. The point of intersection of the two curves represents
- (a) $t_{1/2}$
 (b) $t_{3/4}$
 (c) $t_{2/3}$
 (d) data insufficient to predict.



3. If in a reaction, $A \rightarrow \text{Products}$, the concentrations of reactant A are $C_0, aC_0, a^2C_0, a^3C_0\dots$ after time interval $0, t, 2t, 3t\dots$ where 'a' is constant ($0 < a < 1$). What is the order of reaction?
 (a) Zero order (b) Second order
 (c) First order (d) Third order
4. A first order reaction is 50% completed in 1.26×10^{14} s. How much time would it take for 100% completion?
 (a) 1.26×10^{15} s (b) 2.52×10^{14} s
 (c) 2.52×10^{28} s (d) Infinite
5. For a chemical reaction, $k = PZe^{-E_a/RT}$. Which of the following processes will increase the rate of reaction?
 (a) Lowering of E_a
 (b) Lowering of P
 (c) Lowering of Z
 (d) Independent of all the above factors.
6. If a homogeneous catalytic reaction can take place through three alternative paths as depicted in the given graph, the catalytic efficiency of paths P, Q and R would be




- (a) $P > Q > R$ (b) $Q > P > R$
 (c) $P > R > Q$ (d) $R > Q > P$

7. The activation energies of two reactions are E_1 and E_2 ($E_1 > E_2$). If the temperature of the system is increased from T_1 to T_2 , the rate constant changes from k_1 to k'_1 in the first reaction and k_2 to k'_2 in the second reaction. Predict which of the following expressions is correct?

- (a) $\frac{k'_1}{k_1} = \frac{k'_2}{k_2}$ (b) $\frac{k'_1}{k_1} > \frac{k'_2}{k_2}$
 (c) $\frac{k'_1}{k_1} < \frac{k'_2}{k_2}$ (d) $\frac{k'_1}{k_1} = \frac{k'_2}{k_2} = 1$

8. Cyclopropane rearranges to form propene as shown



This follows first order kinetics. The rate constant is $2.714 \times 10^{-3} \text{ sec}^{-1}$. The initial concentration of cyclopropane is 0.29 M. What will be the concentration of cyclopropane after 100 s?

- (a) 0.035 M (b) 0.22 M
 (c) 0.145 M (d) 0.0018 M
9. In a reaction between A and B, the initial rate of reaction r_0 was measured for different initial concentrations of A and B as given below :

A/mol L ⁻¹	0.20	0.20	0.40
B/mol L ⁻¹	0.30	0.10	0.05
r_0 /mol L ⁻¹ s ⁻¹	5.07×10^{-5}	5.07×10^{-5}	1.43×10^{-4}


The order of the reaction with respect to A is

- (a) 1.5 (b) 0.5 (c) 1 (d) 2
10. In the reaction,
 $\text{CH}_3\text{COCH}_3(l) \longrightarrow \text{C}_2\text{H}_4(g) + \text{H}_2(g) + \text{CO}(g)$
 the initial pressure was 0.40 atm and after 10 min, it was found to be 0.50 atm. The rate constant for first order reaction is [$\log 4 = 0.6021$, $\log 3.5 = 0.5441$]
- (a) 0.0133 min^{-1} (b) 0.4 s^{-1}
 (c) 10 s^{-1} (d) 0.6 min^{-1}

SECTION - II

More than One Options Correct Type

11. Which of the following graphs is correct for a zero order reaction?



12. At 380 °C, the half-life period for the first order decomposition of H_2O_2 is 360 minute. The energy of activation of the reaction is 200 kJ mol^{-1} . What will be the time required for 75% decomposition at 450 °C?

- (a) 20.39 min (b) 30.03 min
 (c) 1223.4 s (d) 2000 s

13. Which of the following statements are correct?

- (a) Increase in concentration of reactant increases the rate of a zero order reaction.
 (b) Rate constant, k is equal to collision frequency 'A' if $E_a = 0$.
 (c) $\log_e k$ vs T is a straight line.
 (d) $\log_e k$ vs $\frac{1}{T}$ is a straight line.

SECTION - III

Paragraph Type

Paragraph for Questions 14 and 15

A secondary alkyl halide (A) hydrolyses with alkali (B) in aqueous medium simultaneously via $\text{S}_{\text{N}}1$ and $\text{S}_{\text{N}}2$ pathways with rate constants k_1 and k_2 , respectively.

From kinetic data, it was found that a plot of $\frac{-1}{[A]} \frac{d[A]}{dt}$

vs $[B]$ is straight line with slope equal to $2.7 \times 10^{-4} \text{ L mol}^{-1} \text{ min}^{-1}$ and intercept equal to $1.02 \times 10^{-3} \text{ L mol}^{-1} \text{ min}^{-1}$. Minimum initial concentration of $[A] = 0.2 \text{ M}$ and $[B]$ i.e., $[\text{OH}^-] = 0.5 \text{ M}$.

14. The value of overall rate constant for the hydrolysis of A (in $\text{L mol}^{-1} \text{ min}^{-1}$) is
 (a) 2.7×10^{-4} (b) 1.02×10^{-3}
 (c) 1.29×10^{-3} (d) none of these.

15. The initial rate of consumption of secondary alkyl halide (in $M \text{ min}^{-1}$) is
- (a) 0.36×10^{-3} (b) 2.89×10^{-5}
 (c) 2.31×10^{-4} (d) 2.31×10^{-3}

Paragraph for Questions 16 and 17

Temperature has a significant effect on the rate of reaction. Quantitatively, the effect of temperature is explained by Arrhenius equation, $k = Ae^{-E_a/RT}$ in which, E_a represents the activation energy of the reaction. Collision theory explains how the rate of reaction depends upon the energy factor and orientation factor *i.e.*, for any collision to be effective, the energy of the colliding molecules should be greater than threshold value and should have proper orientation at the time of collision. Mathematically, collision theory follows, $k = PZ_{AB} e^{-E_a/RT}$.




16. The temperature coefficient for most of the reactions lies between
- (a) 1 and 3 (b) 2 and 3
 (c) 1 and 4 (d) 2 and 4
17. When $\log k$ is plotted against $1/T$, where k is rate constant and T is temperature, a straight line is obtained with a slope of -7000 K . The energy of activation of the reaction is nearly
- (a) 114 kJ mol^{-1} (b) 124 kJ mol^{-1}
 (c) 134 kJ mol^{-1} (d) 144 kJ mol^{-1}

SECTION - IV

Matching List Type

18. Match the List-I with List-II and select the correct answer using the codes given below the Lists :

List I



List II

p. Pseudo first order

q. Zero order

r. Second order

s. First order

- | A | B | C | D |
|----------|------|------|---|
| (a) p | q | r, p | s |
| (b) p | p, r | q | r |
| (c) p | q | r | s |
| (d) r, s | p | q | r |

19. Match the List-I with List-II and select the correct answer using the codes given below the Lists :

List I

List II

- | | |
|--|---------------------|
| A. Mathematical expression for rate of reaction | p. Rate constant |
| B. Rate of reaction for zero order reaction is equal to | q. Rate law |
| C. Units of rate constant for zero order reaction is same as that of | r. Slowest step |
| D. Order of a complex reaction is determined by | s. Rate of reaction |

A B C D

- (a) q p s r
 (b) p q s r
 (c) p q r s
 (d) r p q s

SECTION - V

Assertion Reason Type

20. **Assertion :** A catalyst does not alter the heat of reaction.
Reason : Catalyst increases the rate of reaction.
21. **Assertion :** For molecules to interact, they must approach each other so closely to collide with each other.
Reason : Rearrangement of chemical bonds occurs during collision.
22. **Assertion :** The order of a reaction can have a fractional value.
Reason : The order of reaction cannot be written from the balanced chemical equation.

SECTION - VI

Integer Value Correct Type


23. A reactant (A) forms two products, $A \xrightarrow{k_1} B$; Activation energy E_{a_1}
 $A \xrightarrow{k_2} C$; Activation energy E_{a_2}
 If $E_{a_2} = 2E_{a_1}$, then k_1 and k_2 are related as $k_1 = x \times k_2 e^{E_{a_1}/RT}$. Find out the value of 'x'.
24. The $t_{1/2}$ for the decomposition of CH_3CHO at constant temperature and at initial pressure of 340 mm of Hg and 170 mm of Hg were 410 s and 820 s respectively. Calculate the order of the reaction.
25. For the reaction, $Cl_2 + 2I^- \longrightarrow 2Cl^- + I_2$, the initial concentration of I^- was 0.20 mol L^{-1} and the concentration after 20 min was 0.18 mol L^{-1} . Then, the rate of formation of I_2 in $\text{mol L}^{-1} \text{ min}^{-1}$ is written as $y \times 10^{-4}$. The value of 'y' is

AMINES AND BIOMOLECULES

SECTION - I


Only One Option Correct Type

1. The most reactive amine towards dilute hydrochloric acid is




2. Two aldopentoses 'X' and 'Y' give the same osazone derivative. 'X' is oxidised to an optically active aldaric acid by dilute nitric acid. Ruff degradation of 'Y' gave a tetrose which was similarly oxidised to an optically active aldaric acid.

Assign the structures of 'X' and 'Y' from the following.




- (a) X = I, Y = IV (b) X = IV, Y = I
(c) X = III, Y = II (d) X = II, Y = III

3. A compound of molecular formula $\text{C}_3\text{H}_9\text{N}$ when reacts with benzene sulphonyl chloride gives a product soluble in dilute NaOH solution. The compound should be




4. Which statement is incorrect regarding peptide bond?




- (c) C-N bond length in proteins is smaller than usual bond length of C-N bond.

(d) Both (b) and (c)


5. Propionitrile reacts with 'A' followed by hydrolysis to give 'B'. The compound 'B' on addition of HCN gives 'C'. The compound 'C' on reduction gives 1-amino-2-ethyl-4-methyl-2-pentanol. Identify 'A'.




6. What will be the yields of the given reaction?




7. Which of the following chemical units is certainly to be found in an enzyme?



8. An electric current is passed through an aqueous solution of a mixture of alanine (isoelectric point 6.0), glutamic acid (3.2) and arginine (10.7) buffered at pH 6. What is the fate of the three acids?



9. When the imidazole ring of histidine is protonated, tendency of nitrogen to be protonated (proton migrates from -COOH) is in the order of




- (a) $\beta > \gamma > \alpha$ (b) $\gamma > \beta > \alpha$
 (c) $\gamma > \alpha > \beta$ (d) $\beta > \alpha > \gamma$
10. Which of the following statements are incorrect?
 I. Albumin is a simple protein.
 II. Amino acid alanine contains an acidic side chain.
 III. Insulin is a hormone.
 IV. Muscles contain the protein keratin.
- (a) I, II (b) III, IV
 (c) I, III (d) II, IV

SECTION - II


More than One Options Correct Type

11. Amino acids are classified as acidic, basic or neutral depending upon the relative number of amino and carboxyl groups in their molecule. Which of the following are acidic?
- (a) $(\text{CH}_3)_2\text{CH}-\underset{\substack{| \\ \text{NH}_2}}{\text{CH}}-\text{COOH}$
- (b) $\text{HOOC}-\text{CH}_2-\text{CH}_2-\underset{\substack{| \\ \text{NH}_2}}{\text{CH}}-\text{COOH}$
- (c) $\text{H}_2\text{N}-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{COOH}$
- (d) $\text{HOOC}-\text{CH}_2-\underset{\substack{| \\ \text{NH}_2}}{\text{CH}}-\text{COOH}$
12. Under which of the following reaction conditions, aniline gives *p*-nitro derivative as the major product?
- (a) Acetyl chloride/pyridine followed by conc. H_2SO_4 + conc. HNO_3
 (b) Acetic anhydride/pyridine followed by conc. H_2SO_4 + conc. HNO_3
 (c) Dil. HCl followed by conc. H_2SO_4 + conc. HNO_3
 (d) NaNO_2 + HCl at 0-5 °C

13. The product of the reaction,



is/are?




SECTION - III

Paragraph Type

Paragraph for Questions 14 and 15

RCONH_2 is converted into RNH_2 by means of Hoffmann bromamide degradation.



RCONHBr is formed from which this reaction has derived its name. Electron donating group at phenyl activates the reaction. Hoffmann degradation reaction is an intramolecular reaction.

14. How can the conversion of (i) to (ii) be brought about?
- (a) KBr (b) $\text{KBr} + \text{CH}_3\text{ONa}$
 (c) $\text{KBr} + \text{KOH}$ (d) $\text{Br}_2 + \text{KOH}$
15. Which is the rate determining step in Hoffmann bromamide degradation?
- (a) Formation of (vi) (b) Formation of (ii)
 (c) Formation of (iii) (d) Formation of (iv)

Paragraph for Questions 16 and 17

All monosaccharides (both aldoses and ketoses) containing five or six carbon atoms can have two cyclic structures namely furanose and pyranose forms. Due to cyclic nature, the carbon atom of the aldehyde/keto group of the monosaccharide becomes chiral and consequently each monosaccharide whether in pyranose or in furanose form exists in two stereoisomeric forms called α - and β -anomer. In case, the two stereoisomers differ in configuration at a chiral carbon other than the anomeric carbon, then they are called epimers.

16. α -D-(+)-glucose and β -D-(+)-glucose are
 (a) enantiomers (b) geometrical isomers
 (c) epimers (d) anomers.
17. Glucose and galactose are
 (a) anomers (b) C₂-epimers
 (c) C₃-epimers (d) C₄-epimers.

SECTION - IV

Matching List Type

18. Match the reactions given in List I with the statements given in List II and select the correct answer using the codes given below the lists:

List I		List II	
P. Ammonolysis	1. Amine with lesser number of carbon atoms.		
Q. Gabriel phthalimide synthesis	2. Detection test for primary amines.		
R. Hoffmann Bromamide reaction	3. Reaction of phthalimide with KOH and R—X.		
S. Carbylamine reaction	4. Reaction of alkylhalides with NH ₃ .		

P	Q	R	S
(a) 3	4	1	2
(b) 2	4	3	1
(c) 1	3	4	2
(d) 4	3	1	2

19. Match the enzymes given in List I with their catalytic reaction given in List II and select the correct answer using the codes given below the lists:

List I (Enzymes)		List II (Reactions)	
P. Invertase	1. Decomposition of urea into NH ₃ and CO ₂		
Q. Maltase	2. Conversion of glucose into ethyl alcohol		
R. Pepsin	3. Hydrolysis of maltose into glucose		
S. Urease	4. Hydrolysis of cane sugar		
T. Zymase	5. Hydrolysis of proteins into peptides		

P	Q	R	S	T
(a) 4	3	5	1	2
(b) 2	5	4	3	1
(c) 4	3	1	2	5
(d) 5	1	3	4	2

SECTION - V

Assertion Reason Type

20. **Assertion** : *N, N*-Diethylbenzene sulphonamide is insoluble in alkali.
Reason : Sulphonyl group is attached to electronegative nitrogen atom.
21. **Assertion** : In presence of enzyme, substrate molecule can be attacked by the reagent effectively.
Reason : Active sites of enzymes hold the substrate molecule in a suitable position.
22. **Assertion** : In strongly acidic solutions, aniline becomes less reactive towards electrophilic reagents.
Reason : The amino group being completely protonated in strongly acidic solution, the lone pair of electrons on the nitrogen is no longer available for resonance.

SECTION - VI


Integer Value Correct Type

23. In *D*-talose, the total number of chiral centres is
24. Amongst the following, the total number of amides which undergo Hoffmann bromamide reaction are : C₆H₅CONH₂, C₆H₅CONHCH₃, C₆H₅CON(CH₃)₂, CH₃CONH₂, CH₃CONHCH₃, CH₃CON(CH₃)₂, (CH₃)₂CHCONHCH₃, (CH₃)₂CHCONH₂, (CH₃)₂CHCON(CH₃)₂
25. Amongst the following, the maximum number of water soluble vitamins are A, B₁, B₂, B₆, H, B₁₂, C, D, E and K.


SOLUTIONS

CHEMICAL KINETICS

1. (b): For the reaction,



$$\text{Rate} = -\frac{d[\text{N}_2\text{O}_5]}{dt} = k[\text{N}_2\text{O}_5]$$



$$\text{Rate} = -\frac{1}{2} \frac{d[\text{N}_2\text{O}_5]}{dt} = k'[\text{N}_2\text{O}_5]$$

$$\Rightarrow -\frac{d[\text{N}_2\text{O}_5]}{dt} \Rightarrow 2k'[\text{N}_2\text{O}_5] = k[\text{N}_2\text{O}_5]$$

$$\therefore k = 2k'$$

2. (a)

3. (c)

4. (d): Reaction would be 100% complete only after infinite time which cannot be calculated.

5. (a)

6. (d): The catalytic efficiency of a catalyst is measured in terms of the decrease in activation energy of the reaction. A catalyst is more efficient if it decreases the activation energy to a larger extent. Thus, the correct order of catalytic efficiency is
- $R > Q > P$
- .

7. (b):
- $\log \frac{k'_1}{k_1} = \frac{E_1}{2.303R} \left(\frac{T_2 - T_1}{T_1 T_2} \right)$
- ... (i)
-
- [∵
- $T_2 > T_1$
-]

$$\log \frac{k'_2}{k_2} = \frac{E_2}{2.303R} \left(\frac{T_2 - T_1}{T_1 T_2} \right)$$
 ... (ii)

As, $E_1 > E_2$

From (i) and (ii), we get

$$\log \frac{k'_1}{k_1} > \log \frac{k'_2}{k_2} \Rightarrow \frac{k'_1}{k_1} > \frac{k'_2}{k_2}$$

8. (b):
- $k = \frac{2.303}{t} \log \frac{a}{(a-x)}$

where, $(a-x)$ is the concentration of cyclopropane left after 100 s.

$$2.714 \times 10^{-3} = \frac{2.303}{100} \log \frac{0.29}{(a-x)}$$

$$\Rightarrow \frac{0.2714}{2.303} = \log \frac{0.29}{(a-x)} \Rightarrow 0.118 = \log \frac{0.29}{(a-x)}$$

$$\Rightarrow (a-x) = 0.22 \text{ M}$$

9. (a): Let rate =
- $k[A]^x [B]^y$

From the given data,

$$5.07 \times 10^{-5} = k[0.20]^x [0.30]^y$$
 ... (i)

$$5.07 \times 10^{-5} = k[0.20]^x [0.10]^y$$
 ... (ii)

$$1.43 \times 10^{-4} = k[0.40]^x [0.05]^y$$
 ... (iii)

From eq. (i) and (ii)

$$1 = \frac{[0.30]^y}{[0.10]^y} \Rightarrow y = 0$$

By dividing eq. (iii) by eq. (ii)

$$\frac{1.43 \times 10^{-4}}{5.07 \times 10^{-5}} = \frac{[0.40]^x [0.10]^y}{[0.20]^x [0.05]^y}$$

as $y = 0$

$$\text{Thus, } 2.82 = \left(\frac{0.40}{0.20} \right)^x \Rightarrow (2)^x = 2.82 \Rightarrow x = 1.5$$

10. (a):
- $\text{CH}_3\text{COCH}_3(\text{l}) \longrightarrow \text{C}_2\text{H}_4(\text{g}) + \text{H}_2(\text{g}) + \text{CO}(\text{g})$

Initial 0.4 atm 0 0 0
pressure

After 10 min (0.4 - x) atm x atm x atm x atm

Total pressure = 0.4 - x + x + x + x = 0.50 atm

$$\Rightarrow 2x = 0.10 \text{ or } x = 0.05 \text{ atm}$$

$$k = \frac{2.303}{10} \log \frac{0.4}{0.4 - 0.05} = \frac{2.303}{10} \times 0.0580 = 0.01335 \text{ min}^{-1}$$

11. (a, d): For a zero order reaction,

$$\text{rate} = -\frac{d[A]}{dt} = k \text{ or } \frac{d[A]}{dt} = -k$$

On integrating we get, $\int d[A] = -\int k dt$

$$[A] = -kt + I$$
 ... (i)

When, $t = 0$ then $[A] = [A]_0$

$$\therefore I = [A]_0$$

On putting the value of I in eq. (i)

$$[A] = -kt + [A]_0$$

∴ A graph of $[A]$ vs t is a straight line with slope = $-k$.

12. (a, c):
- $k_1 = \frac{0.693}{t_{1/2}} = \frac{0.693}{360} = 1.925 \times 10^{-3} \text{ min}^{-1}$

$$\log_{10} \frac{k_2}{k_1} = \frac{E_a}{2.303 R} \left(\frac{T_2 - T_1}{T_1 T_2} \right)$$

$$\log_{10} \frac{k_2}{1.925 \times 10^{-3}} = \frac{200 \times 10^3}{2.303 \times 8.314} \left[\frac{723 - 653}{653 \times 723} \right]$$


$$k_2 = 0.068 \text{ min}^{-1}$$

$$t = \frac{2.303}{k_2} \log_{10} \frac{a}{a-x} \Rightarrow \frac{2.303}{0.068} \log_{10} \frac{100}{25}$$


$$\therefore t = 20.39 \text{ min} = 1223.4 \text{ s}$$

13. (b, d)

14. (c): Through
- $\text{S}_{\text{N}}1$
- pathway:



$$\text{Rate} = \frac{-d[\text{A}]}{dt} = k_1[\text{RX}]$$
 ... (i)

Through $\text{S}_{\text{N}}2$ pathway:

$$\text{Rate} = \frac{-d[A]}{dt} = k_2[RX][OH^-] \quad \dots(\text{ii})$$

Combining eq. (i) and (ii),

$$\text{Rate} = \frac{-d[A]}{dt} = k_1[RX] + k_2[RX][OH^-]$$

$$\frac{-d[A]}{dt} = k_1[A] + k_2[A][B]$$

$$\frac{-d[A]}{dt} = [A](k_1 + k_2[B]) \quad \dots(\text{iii})$$

$$\frac{1}{[A]} \frac{-d[A]}{dt} = k_1 + k_2[B] \quad \dots(\text{iv})$$

Comparing this with equation of straight line,
 $y = mx + c$

$$\therefore \text{Slope } (m) = k_2 = 2.7 \times 10^{-4} \text{ L mol}^{-1} \text{ min}^{-1}$$

$$= 0.27 \times 10^{-3} \text{ L mol}^{-1} \text{ min}^{-1}$$

$$\text{Intercept } (c) = k_1 = 1.02 \times 10^{-3} \text{ L mol}^{-1} \text{ min}^{-1}$$

$$\text{Overall rate constant} = k_1 + k_2$$

$$= (0.27 + 1.02) \times 10^{-3} = 1.29 \times 10^{-3} \text{ L mol}^{-1} \text{ min}^{-1}$$

15. (c): $[RX]_0 = [A]_0 = 0.2 \text{ M}$; $[B]_0 = [OH^-] = 0.5 \text{ M}$

$$\text{From Eq. (iii), } \frac{-d[A]}{dt} = [A](k_1 + k_2[B])$$

$$= 0.2(1.02 \times 10^{-3} + 0.27 \times 10^{-3} \times 0.5)$$

$$= 0.2 \times 10^{-3} (1.02 + 0.135)$$

$$= 0.2 \times 10^{-3} \times 1.155 = 2.31 \times 10^{-4} \text{ M min}^{-1}$$

16. (b)

17. (c): Slope = $-\frac{E_a}{2.303 R}$

$$E_a = -2.303 \times R \times \text{slope}$$

$$= -2.303 \times 8.314 \text{ J K}^{-1} \text{ mol}^{-1} \times (-7000 \text{ K})$$

$$= 134029.9 \text{ J mol}^{-1} \approx 134 \text{ kJ mol}^{-1}$$

18. (b) 19. (a) 20. (b)

21. (b) 22. (b)

23. (1): $A \xrightarrow{k_1} B$, $A \xrightarrow{k_2} C$

By Arrhenius equation,

$$k_1 = A' e^{-E_{a1}/RT} \text{ and } k_2 = A' e^{-E_{a2}/RT}$$

$$(A' \text{ is Arrhenius constant}) \quad (\because E_{a2} = 2E_{a1})$$

$$\therefore k_2 = A' e^{-2E_{a1}/RT}$$

$$\frac{k_1}{k_2} = \frac{A' e^{-E_{a1}/RT}}{A' e^{-2E_{a1}/RT}} = e^{E_{a1}/RT} \Rightarrow k_1 = k_2 e^{E_{a1}/RT}$$

On comparing this equation with the given equation, we get $x = 1$.

24. (2): $t_{1/2} \propto [CH_3CHO]^{1-n}$, where n is the order of reaction.

$$\frac{410}{820} = \left(\frac{340}{170}\right)^{1-n} \Rightarrow \log \frac{410}{820} = (1-n) \log \left(\frac{340}{170}\right)$$

$$\log(1/2) = (1-n) \log 2 \Rightarrow n = 2$$

25. (5): Rate of disappearance of $I^- =$

$$\frac{d[I^-]}{dt} = \frac{(0.20 - 0.18) \text{ mol L}^{-1}}{20 \text{ min}}$$

$$= 10^{-3} \text{ mol L}^{-1} \text{ min}^{-1}$$

$$\text{Rate of reaction} = -\frac{1}{2} \frac{d[I^-]}{dt} = +\frac{d[I_2]}{dt}$$

\therefore Rate of formation of $I_2,$

$$\frac{d[I_2]}{dt} = -\frac{1}{2} \frac{d[I^-]}{dt} = \frac{10^{-3}}{2}$$


$$= 5 \times 10^{-4} \text{ mol L}^{-1} \text{ min}^{-1}$$

Comparing it with $y \times 10^{-4}$ then $y = 5$


AMINES AND BIOMOLECULES

- (b): Secondary amines are most reactive towards dilute HCl because electron density on N is more than primary amines due to presence of two alkyl groups. Though tertiary amines have maximum electron density on N but they suffer steric hindrance.
- (c): 'X' and 'Y' have same osazone derivative, it means configuration at C-3 and C-4 is the same.


$X \xrightarrow{\text{Dil. HNO}_3} \text{optically active aldaric acid,}$



Optically active aldaric acid




'Y'




Optically active aldaric acid

3. (b):




4. (a)




6. (a)

7. (c): Since enzymes are proteins they contain -CO-NH- group.

8. (a): At pH = 6, glutamic acid exists as a dianionic species.



So it migrates to anode while arginine exists as cationic species.





So, it migrates to cathode. Alanine does not migrate to any electrode at its isoelectric point.

9. (c)

10. (d)

11. (b, d)


12. (a, b)



15. (d): The rate determining step is probably loss of Br^- to form isocyanates as this is the slowest step.

16. (d): α -D-(+)-glucose and β -D-(+)-glucose are differ in configuration at C_1 and hence are anomers.

17. (d): Glucose and galactose are differ in configuration at C_4 and hence are C_4 -epimers.



18. (d)

19. (a)

20. (b): N, N - Diethylbenzene sulphonamide is a tertiary amine which does not have acidic H attached to N-atom. Thus, it is insoluble in alkali.

21. (a): In presence of enzyme, substrate molecule can be attacked by the reagent effectively because active sites of enzymes hold the substrate molecule in a suitable position.

22. (a)

23. (4)

24. (3): Only 1° amides, i.e., $\text{C}_6\text{H}_5\text{CONH}_2$, CH_3CONH_2 , $(\text{CH}_3)_2\text{CHCONH}_2$ undergo Hoffmann bromamide reaction.

25. (6): Vitamins of B group, i.e., B_1 , B_2 , B_6 , B_{12} , H and vitamin C are water soluble.

Your favourite MTG Books/Magazines available in MANIPUR at

- Fame Book House - Imphal Mob: 9774220848, 7085556697
- Jain Book Shop - Imphal Mob: 9856031157
- Job Centre - Imphal Ph: 0385-2440140; Mob: 9856700700, 9436204282
- P.C. Jain And Co. Ph: 0385-2451756, 2225004; Mob: 9856084649

Visit "MTG IN YOUR CITY" on www.mtg.in to locate nearest book seller OR write to info@mtg.in OR call

0124-6601200 for further assistance.

ACE

YOUR WAY CBSE XII



Series 7

CHAPTERWISE PRACTICE PAPER : BIOMOLECULES | POLYMERS | CHEMISTRY IN EVERYDAY LIFE

Time Allowed : 3 hours


Maximum Marks : 70

GENERAL INSTRUCTIONS

- (i) All questions are compulsory.
- (ii) Q. no. 1 to 5 are very short answer questions and carry 1 mark each.
- (iii) Q. no. 6 to 10 are short answer questions and carry 2 marks each.
- (iv) Q. no. 11 to 22 are also short answer questions and carry 3 marks each.
- (v) Q. no. 23 is a value based question and carry 4 marks.
- (vi) Q. no. 24 to 26 are long answer questions and carry 5 marks each.
- (vii) Use log tables if necessary, use of calculators is not allowed.

1. Classify the following as addition and condensation polymers :
Terylene, bakelite, polyvinyl chloride, polythene
2. The base sequence in a fragment of one strand of DNA is :
AAT TCC CGG
What is the base sequence of the complementary strand?
3. What is the difference between Buna-N and Buna-S?
4. Name the enzyme that converts sucrose into glucose and fructose. Write chemical equation also.
5. Give one important use of each of the following :
(i) Equanil (ii) Morphine
6. Differentiate between nucleotides of DNA and RNA.
7. What are natural and synthetic polymers? Give two examples of each type.
8. What factors are responsible for the stability of α -helix structure of proteins? Why is it named as 3.6_{13} helix?
9. Why do we require artificial sweetening agents?
OR
What are fillers and what role these fillers play in soap?
10. How does the presence of double bonds in rubber molecules influence their structure and reactivity?
11. Define the following as related to proteins :
(i) Peptide linkage (ii) Primary structure
(iii) Fibrous proteins
12. What are anionic detergents? How are they prepared? Write their two main uses.
13. What is a biodegradable polymer? Give an example of a biodegradable aliphatic polyester and the reaction involved in its preparation.
14. Which moieties of nucleotides are involved in the formation of phosphodiester linkages present in dinucleotides? What does the word diester in the name of linkage indicate? Which acid is involved in the formation of this linkage?
OR
Name the sources and the main diseases caused due to lack of vitamins A, H and E.

15. Define the following with examples :
- Semi-synthetic polymers
 - Network polymers
 - Fibres
16. (a) Name the type to which the following drugs belong and give their uses.
- Luminal
 - Streptomycin
 - Analgin
- (b) Give one more example of each class.
17. What are the different types of RNA found in cells of organisms? State the functions of each type.
18. Write the names and molecular structures of the monomers of the following polymers. Give one use of each specifying the property responsible for this use.
- Glyptal
 - Nylon-6
 - Neoprene
19. Explain the role of allosteric site in enzyme inhibition.
20. Phenol and formaldehyde undergo condensation to give a polymer (A) which on heating with formaldehyde gives a thermosetting polymer (B). Name the polymers. Write the reactions involved in the formation of (B). What is the structural difference between these two polymers?
21. (i) Which one of the following is a food preservative?
Equanil, morphine, sodium benzoate
- Why is bithional added to soap?
 - Which class of drugs is used in sleeping pills?
22. (i) What is denaturation and renaturation of proteins?
- Amylase present in the saliva becomes inactive in the stomach. Explain.
23. Gopal and Gaurav both were going to play cricket match in the ground. Suddenly both of them saw an accident between a car and a motorcycle. The biker had fallen down. Both of them moved towards the biker and provided him the first aid. They put dettol over his scratches on hands and legs and gave him a painkiller.
- Why did the boys used dettol over scratches of biker?
 - What is the chemical composition of dettol?
 - What is the chemical present in antiseptic soap?
 - What are the values shown by Gopal and Gaurav?
24. Answer the following questions :
- In *E. Coli* DNA, the AT/GC ratio is 0.93. If the number of moles of adenine in the DNA sample is 4,65,000, calculate the number of moles of guanine present.
 - In an aqueous solution of glucose, it is mainly an equilibrium mixture of α -D-(+)-glucose with specific rotation $[\alpha]_D = +112^\circ$ and β -D-(+)-glucose with specific rotation, $[\alpha]_D = +18.7^\circ$. The specific rotation of the equilibrium mixture is $+52.7^\circ$. Find the percentage composition of the equilibrium mixture.
- OR**
- Explain the following :
 - Sucrose is dextrorotatory but the mixture obtained after hydrolysis is laevorotatory.
 - Amino acids behave like salts rather than simple amines or carboxylic acids.
 - How do enzymes help a substrate to be attacked by the reagent effectively?
25. (i) Name a substance which can be used as an antiseptic as well as disinfectant.
- Why are cimetidine and ranitidine better antacids than sodium bicarbonate or magnesium or aluminium hydroxide?
 - How do antiseptics differ from disinfectants? Give one example of each.
- OR**
- Answer the following questions :
- If water contains dissolved calcium bicarbonate, out of soaps and synthetic detergents which one will you use for cleaning clothes?
 - Label the hydrophilic and hydrophobic parts in the following compounds. Also, give their names.
 - $\text{CH}_3(\text{CH}_2)_{10}\text{CH}_2\text{OSO}_3^-\text{Na}^+$
 - $\text{CH}_3(\text{CH}_2)_{15}\text{N}^+(\text{CH}_3)_3\text{Br}^-$
 - $\text{CH}_3(\text{CH}_2)_{16}\text{COO}(\text{CH}_2\text{CH}_2\text{O})_n\text{CH}_2\text{CH}_2\text{OH}$
26. Answer the following questions :
- On electrolysis in acidic solution, glycine migrates towards cathode while in alkaline solution, it migrates towards anode. Explain.
 - The Fischer projection of *D*-glucose is :



- (i) Write Fischer projection for *L*-glucose.
(ii) Give the product of reaction of *L*-glucose with Tollens' reagent.

OR

- (a) Name the three major classes of carbohydrates and give two examples of each of these classes.
(b) What type of linkage is responsible for the primary structure of proteins?
(c) Name the location where protein synthesis occurs in our body.
(d) Explain the following terms :
(i) Mutarotation (ii) Avitaminoses


SOLUTIONS

1. **Addition polymers** : Polyvinyl chloride, polythene
Condensation polymers : Terylene, bakelite

2. Sequence in the complementary strand :
TTA AGG GCC

3. Buna-N is a copolymer of 1, 3-butadiene and acrylonitrile and Buna-S is a copolymer of 1, 3-butadiene and styrene.

4. Invertase.



5. (i) Equanil is a tranquilizer. It is used for controlling depression and hypertension.

(ii) Morphine is used as an analgesic in severe pain.

6. Nucleotides are monomeric units of nucleic acids. A nucleotide is made up of sugar + heterocyclic base + phosphate group.

In case of DNA nucleotide, the sugar is *D*-2-deoxyribose whereas in RNA nucleotide, it is *D*-ribose. They also differ with respect to one base, DNA contains thymine whereas, RNA contains uracil.

7. **Natural polymers** : Those polymers which are found in plants and animals are called natural polymers *e.g.*, proteins, starch, etc.

Synthetic polymers : Polymers which are prepared in laboratories are called synthetic polymers *e.g.*, plastics (polythene), synthetic fibres (nylon 6, 6), etc.

8. The stability of α -helix structure of proteins is due to intramolecular hydrogen bonding between —NH and —CO groups of the same peptide chain. The α -helix is termed as 3.6₁₃ helix since each turn of the helix has approximately 3.6 amino acids and hydrogen bonding leads to the formation of 13-membered ring.

9. Natural sweeteners, *e.g.*, sucrose and lactose add to calories intake and therefore many people prefer to use artificial sweeteners. *Ortho*-sulphobenzimide, also called saccharin, is the first popular artificial sweetening agent. It has very sweet taste and is about 550 times more sweeter than sucrose. It is excreted from the body through urine unchanged. It appears to be entirely inert and harmless when taken. Its use is of great value to diabetic persons and people who need to control intake of calories.


OR

Substances which are added to soap to change their properties in order to make them more useful for a particular application, are called fillers. For example,

(i) Sodium rosinate, sodium silicate, borax and sodium carbonate are added to laundry soaps to increase their foaming ability.

(ii) Glycerol is added to shaving soaps to prevent them from rapid drying.

10. Natural rubber is *cis*-polyisoprene and is obtained by 1, 4-polymerisation of isoprene units. In rubber molecule double bonds are located between C₂ and C₃ of each isoprene unit. These *cis*-double bonds do not allow the polymer chains to come closer for effective interactions and hence intermolecular forces are quite weak. Thus, natural rubber *i.e.*, *cis*-polyisoprene has a randomly coiled structure and hence shows elasticity.



11. (i) **Peptide linkage** : Proteins are the polymers of α -amino acids which are connected to each other by peptide bonds or peptide linkages. Chemically, peptide linkage is an amide formed between —COOH and —NH₂ groups by the elimination of a water molecule. Thus, peptide bond is a —CO—NH linkage *e.g.*, when carboxyl group of glycine combines with the amino group of alanine we get a dipeptide, glycylalanine.

(ii) **Primary structure** : Proteins may have one or more polypeptide chains. Each polypeptide in a protein has amino acids linked with each other in a specific


sequence and this sequence of amino acids is said to be the primary structure of that protein. Any change in this primary structure *i.e.*, the sequence of amino acids creates a different protein.

(iii) Fibrous proteins : Fibrous proteins consist of linear thread like molecules which tend to lie side by side to form fibres. In these proteins, polypeptide chains are parallel and are held together by hydrogen and disulphide bonds. These are usually insoluble in water, *e.g.*, keratin and myosin.

12. Anionic detergent : A detergent whose large part of the molecule is anion is called anionic detergent.


These are sodium salts of sulphonated long chain alcohols or hydrocarbons *e.g.*, sodium lauryl sulphate, $\text{CH}_3(\text{CH}_2)_{10}\text{CH}_2\text{OSO}_3^-\text{Na}^+$

Preparation : Alkyl hydrogen sulphate formed by treating long chain alcohols with concentrated sulphuric acid, are neutralised with alkali to form anionic detergents.




Uses : These are mostly used (i) for household work (ii) in toothpastes.


13. Polymers which are degraded over a period of time by microorganism like bacteria, etc. are called biodegradable polymers. Biodegradable polymers and their degraded products do not cause any serious effect on the environment. For example, PHBV (Poly- β -hydroxybutyrate-co- β -hydroxyvalerate).



14. A unit formed by the attachment of a base to C_1' position of sugar is known as nucleoside, when nucleoside is linked to phosphoric acid at C_5' position of sugar moiety, a nucleotide is formed.



Nucleotides are joined together by phosphodiester linkage between 5' and 3' carbon atoms of the pentose sugar.




The linkage resembles with two ester groups joined together. Thus, the linkage is known as diester linkage. The acid involved is phosphoric acid (H_3PO_4).

OR

	Source	Deficiency disease
Vitamin A	Ripe yellow fruits, carrots, pumpkin, spinach, milk, fish oil	Night blindness, xerophthalmia (hardening of cornea of eye) and xerosis (drying of skin)
Vitamin E	Spinach, sweet potato, nuts and seeds	Sterility in males and miscarriage in females, mild hemolytic anaemia in newborn infants
Vitamin H	Fish, milk, banana, nuts	Dermatitis, depression and loss of hairs

15. (i) Semi-synthetic polymers : These are obtained from natural polymers by chemical modifications e.g., cellulose derivatives as cellulose nitrate and cellulose acetate (rayon).



(ii) Cross-linked or Network polymers : These are usually formed from bi-functional and tri-functional monomers and contain strong covalent bonds between various linear polymer chains and form a three-dimensional network structure. They are hard and rigid e.g., bakelite, melamine etc.

(iii) Fibres : These have strong intermolecular forces like hydrogen bonding or dipole-dipole interactions. They are useful in making fibres as their molecules are long and thread-like. They possess high tensile strength, high modulus and less elasticity. These strong forces also lead to close packing of chain and imparts crystalline nature e.g., nylon-6,6 (polyamides), terylene (polyester), etc.

16. (a) (i) Luminal : This is a sedative tranquilizer. It produces sleep and is habit forming.

(ii) Streptomycin : It is an antibiotic and used to cure tuberculosis, pneumonia, etc.

(iii) Analgin : It is antipyretic and analgesic. It lowers the body temperature in fever and gives relief from pain.

(b) Tranquilizer : Veronal

Antibiotic : Penicillin

Antipyretic and analgesic : Paracetamol


17. RNA are of three types :

(i) Messenger RNA (*m*-RNA) : It functions as a messenger carrying the information in a gene to the protein synthesising machinery.

(ii) Transfer RNA (*t*-RNA) : They transfer the amino acids from cytoplasm to the protein synthesising machinery.


(iii) Ribosomal RNA (*r*-RNA) : They associate with a set of proteins to form ribosomes. These complex structures, which physically move along an *m*-RNA molecule, catalyse the assembly of amino acids into protein chains. They also bind *t*-RNAs and various molecules necessary for protein synthesis.

18. (i) Glyptal : The monomers are phthalic acid,




It is used in adherent paint because it helps in fixing paints.

(ii) Nylon-6 : The monomer is caprolactum.





It is used for making tyre cords because it has high tensile strength.

(iii) Neoprene : It is a polymer of chloroprene.



It is used for manufacturing conveyor belts in coal mines because it is non-flammable.

19. Some drugs do not bind to the active sites of the enzyme. These bind to a different site of enzyme which is called allosteric site. This binding of allosteric site changes the shape of the active site in such a way that substrate cannot recognise it. If the bond formed between an enzyme and an inhibitor is a strong covalent bond then the enzyme is blocked permanently.



Novolac (A) is linear in structure while bakelite (B) is a cross linked polymer.

21. (i) Sodium benzoate : food preservative; Morphine : narcotic analgesic; Equanil : tranquilizer.

(ii) Bithional is added to soaps to reduce the odours produced by bacterial decomposition of organic matter on the skin, due to its antiseptic properties.

(iii) Tranquilizers are used in sleeping pills because these relieve stress, reduce anxiety and fatigue by inducing sense of well-being.


22. (i) Denaturation of proteins : Protein found in a biological system with a unique three-dimensional structure and biological activity is called native protein. When there is a physical change like change in temperature or chemical change like change in pH in the native form, the hydrogen bonds get disturbed. As a result, globules unfold and helices get uncoiled and protein loses its biological activity. This is known as denaturation of protein. During denaturation, 2° and 3° structures get destroyed but 1° structure remains intact e.g., coagulation of egg white on boiling and curdling of milk.

In case of reversible denaturation process, the protein recovers its original properties and biological activity when the disruptive agent is removed. The reverse of denaturation is called renaturation.

(ii) Amylase works in the mouth where the pH is 6.8–7.2. In the stomach pH is 1–2 thus, denaturation occurs and enzyme becomes inactive.

23. (i) Dettol has an antiseptic property.

(ii) Dettol is a mixture of chloroxylenol and terpineol i.e.,



(iii) Bithional is a compound added to soaps to impart them with antiseptic properties.

(iv) Helpful nature, knowledge of chemistry and social welfare are the values shown by Gopal and Gaurav.

24. (i) The number of moles of adenine (A) must be equal to that of thymine (T).

Therefore,

$$(A + T) = 4,65,000 + 4,65,000 = 9,30,000$$

$$\text{Also, } \frac{A + T}{C + G} = 0.93 \text{ (given)}$$

$$\therefore \text{ Number of moles of } (C + G)$$

$$= \frac{9,30,000}{0.93} = 10,00,000$$

Since the number of moles of (C)

$$= \text{number of moles of (G)}$$

$$\therefore \text{ Number of moles of guanine (G)}$$

$$= \frac{10,00,000}{2} = 500,000$$

(ii) Let the percentage of α -D-glucose present at equilibrium be x .

Assuming that the amount of open-chain form of glucose is negligible at equilibrium, then

Percentage of β -D-glucose at equilibrium = $(100 - x)$

$$\therefore \frac{x \times 112}{100} + \frac{(100 - x) \times 18.7}{100} = 52.7$$

$$\text{or } 112x + 1870 - 18.7x = 52.7 \times 100$$

$$\text{or } (112 - 18.7)x = 5270 - 1870$$

$$93.3x = 3400$$


$$\Rightarrow x = 36.44\%$$

\therefore Percentage of α -D-(+)-glucose = 36.44%

and percentage of β -D-(+)-glucose = 63.56%


OR

(a) (i) Sucrose is dextrorotatory having $[\alpha]_D = +66.5^\circ$. On hydrolysis with dilute acids or enzymes, it gives equimolar D-(+)-glucose and D-(-)-fructose.



Since, D-(-)-fructose has larger specific rotation than D-(+)-glucose, the resulting mixture has specific rotation of -39.9° . Therefore, the mixture is laevorotatory.

(ii) Salt like behaviour of amino acids is due to the presence of both acidic and amino group in the same molecule. In aqueous solution, the carboxyl group loses a proton and amino group accepts a proton to form a dipolar ion or zwitter ion.



Therefore, they behave like salts rather than simple amines or carboxylic acids.

(b) Active site of enzymes hold the substrate molecule in a suitable position, so that it can be attacked by the reagent effectively.

25. (i) 0.2% solution of phenol acts as an antiseptic while 1% solution of phenol acts as a disinfectant.

(ii) If excess of NaHCO_3 , Mg(OH)_2 or Al(OH)_3 is used, it will make the stomach alkaline and trigger the production of even more HCl which may cause ulcers in the stomach. In contrast, cimetidine and ranitidine prevent the interaction of histamine with the receptor cells present in the stomach wall. This releases lesser amount of HCl.

(iii) Antiseptics are the chemical substances which are used to either kill or prevent the growth of microorganisms. These are not harmful to living tissues and can be safely applied on wounds, cuts, diseased skin surfaces. For example, dettol, savlon, furacin, soframycin, etc.

Disinfectants are the chemical substances which kill microorganisms but they cannot be applied on living tissues. In other words, they also kill microorganisms like antiseptics but are not safe for living tissues. These are commonly applied to inanimate objects such as floor, drainage system, instruments, etc. Some common examples of disinfectants are phenol (1% solution), chlorine (0.2 to 0.4 ppm), etc.

OR

(i) Calcium bicarbonate makes water hard. Soap will give precipitate with this hard water and therefore, cannot be used for cleaning clothes. On the other hand, a synthetic detergent does not give precipitate in hard water because calcium ions form soluble salts that act as cleansing agent. Therefore, synthetic detergent should be used for cleaning clothes in hard water.

(ii) (a) $\text{CH}_3(\text{CH}_2)_{10}\text{CH}_2 - \text{OSO}_3^- \text{Na}^+$
 Hydrophobic part Hydrophilic part

Name : Sodium lauryl sulphate


(b) $\text{CH}_3(\text{CH}_2)_{15} - \text{N}^+(\text{CH}_3)_3\text{Br}^-$
 Hydrophobic part Hydrophilic part

Name : Cetyltrimethylammonium bromide


(c) $\text{CH}_3(\text{CH}_2)_{16} - \text{COO}(\text{CH}_2\text{CH}_2\text{O})_n\text{CH}_2\text{CH}_2\text{OH}$
 Hydrophobic part Hydrophilic part

Name : Polyethylene glycol stearate



26. (a) Glycine exists as zwitter ion, $\text{NH}_3^+ - \text{CH}_2 - \text{COO}^-$. In acidic solution, it exists as cation (I), which migrates towards cathode on electrolysis.



In alkaline solution, glycine exists as anion (II), which on electrolysis migrates towards anode.



(b) (i) The Fischer projection of *L*-glucose is the mirror image of *D*-glucose.



OR

(a) (i) **Monosaccharides** : Glucose, fructose

(ii) **Oligosaccharides** : Sucrose, raffinose

(iii) **Polysaccharides** : Cellulose, starch

(b) Peptide linkage

(c) Ribosomes

(d) (i) **Mutarotation** : The spontaneous change in specific rotation of an optically active compound with time to an equilibrium value is called mutarotation.

Both forms of glucose are in equilibrium in the solution.
 α -*D*-glucose \rightleftharpoons Open chain form \rightleftharpoons β -*D*-glucose
 of equilibrium mixture

Sp. rotation	Sp. rotation	Sp. rotation
= +111°	= +52.5°	= +19.2°

(ii) **Avitaminoses** : Multiple deficiencies caused by lack of more than one vitamin are more common in human beings. This condition of vitamin deficiency is known as avitaminoses.



MPP-8 MONTHLY Practice Problems

Class XII

This specially designed column enables students to self analyse their extent of understanding of specified chapter. Give yourself four marks for correct answer and deduct one mark for wrong answer. Self check table given at the end will help you to check your readiness.



Aldehydes, Ketones and Carboxylic Acids



Total Marks : 120

Time Taken : 60 Min.


NEET / AIIMS

Only One Option Correct Type

- The reagent used for separation of acetaldehyde and acetophenone is
 (a) NaHSO_3 (b) $\text{C}_6\text{H}_5\text{NHNH}_2$
 (c) NH_2OH (d) NaOH/I_2
- When propionic acid is treated with aqueous sodium bicarbonate, CO_2 is liberated. The 'C' of CO_2 comes from
 (a) methyl group (b) carboxylic acid group
 (c) methylene group (d) bicarbonate group.
- An alkene on ozonolysis gives two moles of HCHO , one mole of CO_2 and one mole of CH_3COCHO . What is its structure?
 (a) $\text{CH}_2=\text{C}=\text{CHCH}_2\text{CH}_3$
 (b) $\text{CH}_2=\text{CHCH}(\text{CH}_3)\text{CH}=\text{CH}_2$
 (c) $\text{CH}_2=\text{C}=\text{C}(\text{CH}_3)_2$
 (d) $\text{CH}_2=\text{C}=\text{C}(\text{CH}_3)\text{CH}=\text{CH}_2$
- A carbonyl compound with molecular weight 86, does not reduce Fehling's solution but forms crystalline bisulphite derivatives and gives iodoform test. The possible compounds can be
 (a) 2-pentanone and 3-pentanone
 (b) 2-pentanone and 3-methyl-2-butanone
 (c) 2-pentanone and pentanal
 (d) 3-pentanone and 3-methyl-2-butanone.
- Identify the main product in the given reaction.




- Which of the following can reduce ester to alcohol?
 (a) NaBH_4 (b) $\text{Na}/\text{alcohol}$
 (c) H_2/Ni (d) NaBH_3CN
- The final product (C) in the given reaction is



8. Which of the following does not give formaldehyde on heating or distillation?

- (a) Formalin (b) Trioxane
(c) Paraldehyde (d) Paraformaldehyde

9. Which of the following is the correct order of pK_a values for the given compounds?







- (a) $A > B > C$ (b) $C > B > A$
(c) $B > A > C$ (d) $C > A > B$






10. $R\text{CH}_2\text{CH}_2\text{OH}$ can be converted to $R\text{CH}_2\text{CH}_2\text{COOH}$ by which of the following sequence of steps?

- (a) $\text{PBr}_3, \text{KCN}, \text{H}_3\text{O}^+$ (b) $\text{PBr}_3, \text{KCN}, \text{H}_2/\text{Ni}$
(c) $\text{KCN}, \text{H}_3\text{O}^+$ (d) $\text{HCN}, \text{PBr}_3, \text{H}_3\text{O}^+$

11. Product (A) formed in the given reaction



12. The suitable reaction steps for the following transformation,



Assertion & Reason Type

Directions : In the following questions, a statement of assertion is followed by a statement of reason. Mark the correct choice as :

- (a) If both assertion and reason are true and reason is the correct explanation of assertion.
(b) If both assertion and reason are true but reason is not the correct explanation of assertion.
(c) If assertion is true but reason is false.
(d) If both assertion and reason are false.

13. **Assertion :** Aromatic aldehydes and formaldehyde undergo Cannizzaro reaction.

Reason : Aromatic aldehydes are almost as reactive as formaldehyde.

14. **Assertion :** *p*-Hydroxybenzoic acid has lower boiling point than *o*-hydroxybenzoic acid.

Reason : There exists intermolecular hydrogen bonding in *o*-hydroxybenzoic acid while intramolecular hydrogen bonding in *p*-hydroxybenzoic acid.

15. **Assertion :** Lower aldehydes and ketones are soluble in water but the solubility decreases as molecular mass increases.

Reason : Aldehydes and ketones can be distinguished by Tollens' test.

NEW LAUNCH

ONLINE TEST SERIES

Practice Part Syllabus/ Full Syllabus
40 Mock Test for

JEE Main


Now on your android & IOS Smart phones
with the same login of web portal.

Log on to <http://pcmbtoday.com>


JEE MAIN / JEE ADVANCED

Only One Option Correct Type

16. Compound 'A' (molecular formula, C_3H_8O) is treated with acidified $K_2Cr_2O_7$ to form a product 'B' (molecular formula, C_3H_6O). 'B' forms a shining silver mirror on heating with ammoniacal silver nitrate. 'B' when treated with an aqueous solution of $H_2NCONHNH_2 \cdot HCl$ and sodium acetate gives a product 'C'. Identify the structure of 'C'.
- (a) $CH_3CH_2CH = NNHCONH_2$
 (b) $(CH_3)_2C = NNHCONH_2$
 (c) $(CH_3)_2C = NCONHNH_2$
 (d) $CH_3CH_2CH = NCONHNH_2$
17. The one which has least iodine value is
 (a) ginger oil (b) ghee
 (c) groundnut oil (d) sunflower oil.
18. The correct set of reagents for the following conversion is




- (a) $P/I_2, Na, \text{conc. } H_2SO_4$
 (b) $P_2O_5, LiAlH_4$
 (c) $P_2O_5, H_2O, P/I_2, Na$
 (d) $P/I_2, Na, P_2O_5$
19. Which of the following has the largest value of dissociation constant (K_a)?



More than One Options Correct Type

20. Which of the following reactions give benzophenone?
- (a) $2C_6H_6 + CCl_4 \xrightarrow[\text{(ii) } H_2O]{\text{(i) } AlCl_3}$
 (b) $C_6H_6 + C_6H_5COCl \xrightarrow{AlCl_3}$
 (c) $o-CH_3C_6H_4COC_6H_5 \xrightarrow{\text{heat}}$
 (d) $o-HOOC_6H_4COC_6H_5 \xrightarrow[260^\circ C]{Cu}$

21. Which of the following are examples of aldol condensation?



22. Which of the following statements are true about $\begin{array}{c} \diagdown \\ C \\ \diagup \end{array} = O$ group?

- (a) The carbon atom of the carbonyl group is sp^2 hybridized.
 (b) The $C = O$ bond length is longer than that of $C = C$ bond length.
 (c) The portion of the molecule immediately surrounding the carbonyl group is planar.
 (d) None of the above.

23. Which of the following statements are not true about $HCOOH$?


- (a) It gives CO and H_2O on heating with conc. H_2SO_4 .
 (b) It reduces Tollens' reagent.
 (c) It is a weaker acid than CH_3COOH .
 (d) It forms formyl chloride with PCl_5 .

Integer Answer Type


24. A compound $A(C_5H_8O_2)$ is reduced to pentane with $Zn-Hg/HCl$. It forms a dioxime with NH_2OH and also gives positive iodoform and Tollens' tests. The number of ketonic groups present in A is
25. Amongst the following, the number of carboxylic acids which easily undergo decarboxylation on heating are : $HCOOH$, $(COOH)_2$, $CH_2(COOH)_2$, $C_6H_5COCH_2COOH$, $C_6H_5COCOOH$,

EXAM CORNER

Exam	Date
JEE Main (Offline)	8 th April, 2018
WB JEE	22 nd April, 2018
JEE Advanced	20 th May, 2018
AIIMS	27 th May, 2018




26. In the given sequence of reactions, the total number of intramolecular aldol condensation products formed from 'Y' is



Comprehension Type

An acyclic hydrocarbon *P*, having molecular formula C_6H_{10} , gave acetone as the only product through the following sequence of reactions, in which *Q* is an intermediate compound.



27. The structure of the compound *P* is
- $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{C}\equiv\text{CH}$
 - $\text{CH}_3\text{CH}_2\text{C}\equiv\text{CCH}_2\text{CH}_3$
 - $(\text{CH}_3)_2\text{CHC}\equiv\text{CCH}_3$
 - $(\text{CH}_3)_3\text{CC}\equiv\text{CH}$
28. The structure of the compound *Q* is

- $(\text{CH}_3)_2\text{CHCH}(\text{OH})\text{CH}_2\text{CH}_3$
- $(\text{CH}_3)_3\text{CCH}(\text{OH})\text{CH}_3$
- $(\text{CH}_3)_2\text{CHCH}_2\text{CH}(\text{OH})\text{CH}_3$
- $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}(\text{OH})\text{CH}_2\text{CH}_3$

Matrix Match Type

29. Match column I with column II and choose the correct option using the codes given below.

Column I	Column II
(A) $\text{CH}_3-\text{CH}=\text{CH}-\overset{\text{O}}{\parallel}{\text{C}}-\text{H} \rightarrow$	(p) $\text{H}_2/\text{Pd}-\text{BaSO}_4$
$\text{CH}_3-\text{CH}=\text{CH}-\text{CH}_2\text{OH}$	
(B) $\text{C}_6\text{H}_5-\text{C}\equiv\text{N} \rightarrow$	(q) LiAlH_4
$\text{C}_6\text{H}_5\text{CHO}$	
(C) $\text{C}_6\text{H}_5-\overset{\text{O}}{\parallel}{\text{C}}-\text{Cl} \rightarrow$	(r) $\text{SnCl}_2/\text{HCl}, \text{H}_2\text{O}/\text{H}^+$
$\text{C}_6\text{H}_5\text{CHO}$	
(D) $\text{C}_6\text{H}_5-\text{CH}=\text{CH}-\overset{\text{O}}{\parallel}{\text{C}}-\text{H} \rightarrow$	(s) NaBH_4
$\text{C}_6\text{H}_5\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$	

A	B	C	D
(a) p	r	s	q
(b) q	s	p	r
(c) s	q	r	p
(d) s	r	p	q

30. Match column I with column II and choose the correct option using the codes given below.

Column I	Column II
(A) $(\text{COOH})_2 \cdot 2\text{H}_2\text{O} \xrightarrow[105^\circ\text{C}]{\Delta} \rightarrow$	(p) $2\text{CO}_2 + \text{H}_2\text{O}$
(B) $\begin{matrix} \text{COOH} \\ \diagdown \\ \text{C} \\ \diagup \\ \text{COOH} \end{matrix} \xrightarrow[\Delta]{\text{P}_2\text{O}_5} \rightarrow$	(q) $(\text{COOH})_2$
(C) $(\text{COOH})_2 \cdot 2\text{H}_2\text{O} \xrightarrow[\text{KMnO}_4]{\Delta} \rightarrow$	(r) Carbon suboxide
(D) $\begin{matrix} \text{COOH} \\ \diagdown \\ \text{C} \\ \diagup \\ \text{COOH} \end{matrix} \xrightarrow[\text{reflux, H}_2\text{SO}_4]{\Delta, 140-150^\circ\text{C} \text{ or}} \rightarrow$	(s) CH_3COOH

A	B	C	D
(a) p	q	s	r
(b) r	s	q	p
(c) q	r	p	s
(d) p	q	r	s



Keys are published in this issue. Search now! ☺

SELF CHECK

No. of questions attempted
No. of questions correct
Marks scored in percentage

Check your score! If your score is

> 90%	EXCELLENT WORK !	You are well prepared to take the challenge of final exam.
90-75%	GOOD WORK !	You can score good in the final exam.
74-60%	SATISFACTORY !	You need to score more next time.
< 60%	NOT SATISFACTORY !	Revise thoroughly and strengthen your concepts.


CHEMISTRY MUSING

PROBLEM SET 53


Chemistry Musing was started from August '13 issue of Chemistry Today. The aim of Chemistry Musing is to augment the chances of bright students preparing for JEE (Main and Advanced) / NEET / AIIMS / JIPMER with additional study material. In every issue of Chemistry Today, 10 challenging problems are proposed in various topics of JEE (Main and Advanced) / NEET. The detailed solutions of these problems will be published in next issue of Chemistry Today. The readers who have solved five or more problems may send their solutions. The names of those who send atleast five correct solutions will be published in the next issue. We hope that our readers will enrich their problem solving skills through "Chemistry Musing" and stand in better stead while facing the competitive exams.

JEE MAIN/NEET




1. Consider the following reaction,




Here, X and Y respectively are




2. Consider the following set of reactions,



Here, X, Y and Z are



3. Observe the given sequence of reactions and choose the incorrect statement.




- (a) 'T' is formed by coupling reaction.
 (b) 'R' on heating with ammonia gives benzamide.

(c) 'S' is a stable compound and can be isolated in solid form.

(d) 'Q' reacts with Hinsberg's reagent to give a sulphonamide which is soluble in alkali.

4. Consider the following reaction,



the correct statement about P is

- (a) the anion present in crystalline P exists as a dimer due to hydrogen bonding
 (b) P cannot be isolated in solid state
 (c) P is generally called potash alum
 (d) P gives thermally unstable carbonate on heating.
5. A gas is expanded from volume V_1 to V_2 through three different processes :

I. Reversible adiabatic

II. Reversible isothermal

III. Irreversible adiabatic (against a constant external pressure, P_{ex})

The correct option is

- (a) $(T_f)_{\text{Reversible isothermal}} > (T_f)_{\text{Reversible adiabatic}} > (T_f)_{\text{Irreversible adiabatic}}$
 (b) $(T_f)_{\text{Reversible isothermal}} > (T_f)_{\text{Irreversible adiabatic}} > (T_f)_{\text{Reversible adiabatic}}$
 (c) $W_{\text{Reversible isothermal}} > W_{\text{Irreversible adiabatic}} > W_{\text{Reversible adiabatic}}$
 (d) $W_{\text{Irreversible adiabatic}} > W_{\text{Reversible isothermal}} > W_{\text{Reversible adiabatic}}$

JEE ADVANCED

6. For a homogenous gaseous reaction, $\text{X}_{(g)} + 2\text{Y}_{(g)} \rightleftharpoons \text{Z}_{(g)}$ at 473 K, the value of $K_c = 0.35$. When 2 moles of Y are mixed with 1 mole of X, what would be the pressure when 60% of X is converted to Z?

- (a) 200 atm (b) 180 atm
 (c) 400 atm (d) 150 atm

Contd. on page no. 89








CONCEPT BOOSTER

Dear Students! I am back again with another mind blowing question-answer series of ORGANIC CHEMISTRY. Questions are of quite standard and conceptual. Try to solve every question by yourself and then look at the solution and don't do the reverse!! Hope you will love to learn. Do take care.


*Arunava Sarkar

VERY IMPORTANT PROBLEMS

1. Glycerol $\xrightarrow{\text{anhy. KHSO}_4}$ A $\xrightarrow{\text{Al(OEt)}_3}$ B;
Identify A and B.













Identify the final product.



SOLUTIONS

1. Glycerol undergoes dehydration when heated with KHSO_4 as follows :





Your favourite MTG Books/Magazines available in **HIMACHAL PRADESH** at

- Shiv Shankar Copy House - Bilaspur(Hp) Mob: 9816255326
- Dadhwal Book Depot - Dharamshala Ph: 01892-224867; Mob: 9418087897
- Aryan Enterprises - Hamirpur Ph: 01972-222180; Mob: 9817036236, 9817034100
- Bhagat Enterprises - Hamirpur Ph: 1972233121; Mob: 9817475121, 9817475121
- Gautam Brothers - Hamirpur Ph: 01972-222450; Mob: 9418016277
- Rajesh General Store - Hamirpur Ph: 01972-223037; Mob: 9418024037
- Narendra Book Stall - Mandi Ph: 01905-225578; Mob: 9805471400
- Raj Book Depot - Mandi Ph: 01905-230086; Mob: 9418926708
- Jain Bros. - Nahan Ph: 01702-224253; Mob: 9418265353
- Goel Book Depot - Palampur Ph: 01894-230853; Mob: 9816630751, 9816030751
- Vickey General Store - Paonta Sahib Ph: 01704-225335; Mob: 9418088230
- Kharbanda General Store - Sundarnagar Ph: 01907-262462; Mob: 9418000462

Visit "MTG IN YOUR CITY" on www.mtg.in to locate nearest book seller OR write to info@mtg.in OR call **0124-6601200** for further assistance.


JEE Advanced

PRACTICE PROBLEMS

SECTION-1 (MAXIMUM MARKS : 40)

This section contains 10 multiple choice questions. Each question has 4 choices (a), (b), (c) and (d), out of which one or more than one answer is correct. For each question you will be awarded 4 marks if you have darkened only the bubble corresponding to the correct answer and zero mark if no bubble is darkened. No negative marking in this section.

- Which of the following statements regarding a peptide linkage in a protein molecule is/are correct?
 - It is an amide linkage.
 - It has partial double bond character.
 - It is hydrophilic in nature.
 - It connects protein molecules through H-bonds.
- Under Wolff-Kishner reduction conditions, the conversions which may be brought about are
 - benzophenone into diphenylmethane
 - benzaldehyde into benzyl alcohol
 - cyclohexanone into cyclohexane
 - cyclohexanone into cyclohexanol.
- Metallurgies of which of the following metals involve(s) leaching?
 - Au
 - Ag
 - Al
 - Fe
- Find the correct statement(s).
 - Schottky defect is also called dislocation defect.
 - Unit cell having crystal parameters, $a = b \neq c$, $\alpha = \beta = 90^\circ$, $\gamma = 120^\circ$ is hexagonal.
 - In ionic compounds having Frenkel defect, the ratio r^+/r^- is high.
 - The coordination number of Na^+ ion in NaCl is 8.
- For cyclohexane, four types of conformations have been found. These are chair, half chair, twist boat and boat form. Energy diagram of different conformers of cyclohexane is given. Choose the correct option(s).




- Point R corresponds to chair form.
 - Point S corresponds to chair form.
 - Point Q corresponds to boat form.
 - None of these.
- During one of his adventures Chacha Chaudhary got trapped in an underground cave which was sealed two hundred years back. The air inside was poisonous and contains CO in addition to O_2 and N_2 . Sabu, being huge, could not enter cave. In order to save Chacha Chaudhary he started sucking the poisonous air out of the cave by mouth. In each cycle he used to fill his lungs with cave air and exhale it out in the surroundings. In the mean time fresh air ($\text{N}_2 + \text{O}_2$) from the surrounding effused into cave till the pressure was 1 atmosphere. Each time Sabu sucked air, the pressure in the cave dropped to $1/2$ atm. An initial sample of air taken from the cave measured 11.2 mL at STP and give 7 J on complete combustion at constant pressure.
 - If the safe level of CO required in cave for life is less than 0.001% by volume, how many times does Sabu need to suck out air in order to save Chacha Chaudhary.
 - Sabu should rescue Chacha Chaudhary within 10 minutes else he will die. Precious 80 s was wasted in thinking of a way to rescue him. At maximum how much time should each cycle of inhaling and exhaling take?

[Given : $\Delta H_{\text{comb}} \text{CO} = -280 \text{ kJ mol}^{-1}$. Neglect Graham's law effect during operations.]


- (a) 8 times (b) 13 times
(c) 40 s (d) 20 s

7. The diagram shows an apparatus to find the transition temperature (18 °C) at which white tin and grey tin are in equilibrium. Below 18 °C, white tin dissolves from W and is deposited on G as grey tin.





Which of the following statements are correct?

- (a) The stable form of tin at 25 °C is grey.
(b) Below 18°C, electrons flow through the external circuit from W to G.
(c) At 18 °C, no current flows.
(d) At 25 °C, current flows from W to G.
8. The gas phase decomposition of dimethyl ether follows first order kinetics,
 $\text{CH}_3\text{—O—CH}_3(\text{g}) \longrightarrow \text{CH}_4(\text{g}) + \text{H}_2(\text{g}) + \text{CO}(\text{g})$
 The reaction is carried out in a constant volume container at 500 °C and has a half life of 0.116 hours. Initially only dimethyl ether is present at a pressure of 10 atm. (Assume ideal gas behaviour.)
 (a) The total pressure of the system after 23.03 min is 9 atm.
 (b) The total pressure of the system after 23.03 min is 28 atm.
 (c) The partial pressure of dimethyl ether after 23.03 min is 1 atm.
 (d) The partial pressure of methane after 23.03 min is 9 atm.
9. Which of the statements regarding following structures is true?




- (a) I and II are epimers.
(b) I and III are epimers.
(c) I and II are enantiomers.
(d) I and III are enantiomers.

10. Choose the correct characteristic test(s) of anions.
 (a) Sulphide : With dilute H_2SO_4 , H_2S is evolved which turns lead acetate paper black and with sodium nitroprusside, sulphide gives purple colour.





- (b) Sulphite : With dilute H_2SO_4 , SO_2 is released which turns potassium dichromate paper green.






- (c) Nitrite : With dilute H_2SO_4 , nitric oxide is released which combines with atmospheric oxygen to give reddish-brown fumes. The nitric oxide turns ferrous sulphate solution brownish-black ($\text{FeSO}_4 \cdot \text{NO}$).

- (d) Chlorides : With concentrated H_2SO_4 , HCl gas is released which gives white fumes with ammonia. Silver nitrate solution gives white precipitate with Cl^- which is soluble in ammonia.



The characteristic test of Cl^- is chromyl chloride test :



SECTION-2 (MAXIMUM MARKS : 40)


This section contains 10 integer type questions. Answers are to be given in between 0 to 9 in the form of nearest integer. Each question carries 4 marks if you darken the correct answer and no negative mark will be awarded for an incorrectly bubbled answer.

11. When $\text{C}_2\text{H}_5\text{COONa}$ is electrolysed, the main organic compound formed is $\text{C}_p\text{H}_q\text{O}_r$. Then integer value of 'r' is
12. How many of the following sulphides salts are black in colour?
 CdS , ZnS , NiS , PbS , CuS , Bi_2S_3
13. A certain gas diffuses from two different vessels P and Q. The vessel P has a circular orifice while vessel Q has square orifice of length equal to the radius

of the orifice of vessel *P*. The rates of diffusion of the gas from vessel *P* to vessel *Q* under similar temperature and pressure condition is


14. 1 g sample of pure organic compound containing chlorine is fused with Na_2O_2 to convert chlorine to NaCl . The sample is then dissolved in water and the chloride precipitated with AgNO_3 , giving 1.96 g of AgCl . If the molecular weight of organic compound is 147, how many chlorine atoms does each molecule contain?

15. Amongst the following, the total number of compounds which are soluble in aqueous NaOH is



16. Read the following statements from 0 to 9 and choose the incorrect statement.

0. Both glucose and fructose give positive test with Tollens' reagent.




gives positive test with Tollens' reagent.

2. Thermosetting polymers are the polymers which undergo permanent change on heating.
 3. A polymer formed by the condensation of two or more than two monomers with the elimination of simple molecules like water, ammonia, hydrogen chloride, alcohol, etc. is called condensation polymer.
 4. Buna-S is obtained by polymerization of

1,3-butadiene and styrene in the ratio of 3 : 1 in presence of sodium.

5. Natural rubber is a polymer of chloroprene.
 6. Glycine gives purple colour with ninhydrin while acetamide does not.
 7. Raffinose is a trisaccharide and on hydrolysis it gives three molecules of monosaccharide units one each of glucose, fructose and galactose.
 8. According to Maxwell distribution law, at high temperatures a greater fraction of the molecules can be expected to have high speed than at low temperatures.
 9. The entropy of a perfectly ordered crystalline substance at 0 K is zero.



In the substitution product, $-\text{NH}_2$ group is present at C-___ position.

18. The vapour pressure of pure liquid *A* is 0.80 atm. When a non-volatile liquid *B* is added to *A* its vapour pressure drops to 0.60 atm. The mole fraction of *B* in the solution is $1/x$. Find the value of *x*.
 19. In antifluorite structure coordination number of anion is ___.
 20. 3 moles of acetone when heated with dry HCl gives phoron. How many sp^2 hybridised carbon atoms are present in phoron?

ANSWER KEY

1. (a,b,c,d) 2. (a,c) 3. (a,b,c) 4. (b)
 5. (d) 6. (b,c) 7. (b,c,d)
 8. (b,c,d) 9. (a,b) 10. (a,b,c,d) 11. (0)
 12. (4) 13. (3) 14. (2) 15. (7)
 16. (5) 17. (3) 18. (4) 19. (8)
 20. (5)

For detailed solution to JEE Advanced Practice Problems, visit our website : www.vidyalankar.org

Solution Senders of Chemistry Musing

Set - 52

- Subhankar Panda, West Bengal
- Raghav Malhotra, Maharashtra

Set - 51

- Anshuman Pan, West Bengal

CHEMISTRY MUSING

SOLUTION SET 52

1. (d): Molar volume from pyknometric density

$$= \frac{\text{Weight or mass}}{\text{Density}} = \frac{\text{Weight } (w)}{2.165 \times 10^3} \text{ m}^3$$


$$\text{Molar volume from X-ray density} = \frac{w}{2.178 \times 10^3} \text{ m}^3$$

$$\begin{aligned} \text{Volume unoccupied} &= \frac{w}{10^3} \left(\frac{1}{2.165} - \frac{1}{2.178} \right) \text{ m}^3 \\ &= \frac{0.013 w \times 10^{-3}}{2.165 \times 2.178} \end{aligned}$$


∴ Fraction unoccupied

$$= \frac{\left(\frac{0.013 w \times 10^{-3}}{2.165 \times 2.178} \right)}{\left(\frac{w \times 10^{-3}}{2.165} \right)} = 5.96 \times 10^{-3}$$

2. (a): $4B + 3O_2 \xrightarrow{700^\circ\text{C}} 2B_2O_3$




'A' (white amorphous powder) 'B'




'C' 'D' (white powder)

'D' turns red litmus blue because it is basic in nature and gives brown ppt. with K_2HgI_4 .




Brown ppt.

Its aqueous solution gives reddish brown ppt. with Nessler reagent, K_2HgI_4 .



'C' 'E'


3. (b): Leaching of Carnotite,



4. (d): Reaction $T_2O_3 \longrightarrow T$ is possible above that temperature where line of T crosses the line of R .

5. (c): $LiCl \cdot 3NH_3(s) \rightleftharpoons LiCl \cdot NH_3(s) + 2NH_3(g)$

$$K_p = 9 \text{ atm}^2$$



$$t = 0 \quad 0.1 \quad a \quad 0$$

$$t = t_{eq} \quad 0 \quad (a - 0.2) \quad 0.1$$

Initially a moles of NH_3 should be added to bring in completion of reaction.

$$K_{p1} = \frac{1}{(p'_{NH_3})^2} \Rightarrow \frac{1}{9} = \frac{1}{(p'_{NH_3})^2}$$

$$\Rightarrow p'_{NH_3} = 3 \text{ atm}$$

$$PV = nRT$$

$$3 \times 5 = n \times 0.0821 \times 313$$

$$n = 0.5837$$


$$(a - 0.2) = 0.5837$$

Mole of $NH_3 = a = 0.5837 + 0.2$

$= 0.7837$ mole of NH_3 should be added for completion of reaction.


6. (c): In case of octahedral complexes of d^4 ions when $\Delta_o > P$, low spin complexes are formed and when $\Delta_o < P$, high spin complexes are formed. In case of octahedral complexes of d^9 ions in both cases the number of unpaired electrons is same.

(i) For d^9 complex



Distribution of d^9 electrons in t_{2g} and e_g orbitals

(ii) For d^4 complex,




$\Delta_o > P$ (LS complex) $(n = 2)$

$\Delta_o < P$ (HS complex) $(n = 4)$

Distribution of d^4 electrons in t_{2g} and e_g orbitals

7. (b): The structure given in option (b) contributes heavily to the dipole moment as it shows greater aromatic character than others.



8. (d): The all *cis*-isomer (a), a fully convex decagon, would have angles of 144° which creates large amount of angle strain.


The all *cis*-isomer can adopt boat like conformation.

(c) to relieve the angle strain but it is still unstable as compared to *trans, cis, trans, cis, cis* isomer (b). (b) is unstable due to steric repulsion between two internal hydrogens.

The non-planar *trans, cis, cis, cis, cis* isomer (d) is the most stable of all the possible isomers.

9. (4): If the isoelectric point (pI) of a polypeptide is more than 7, then it will exist as positively charged at pH = 7. This is possible only if the polypeptide is basic in nature, *i.e.*, it has two or more amino groups. In other words, side chain R_1 and R_2 must contain $-\text{NH}_2$ groups but not $-\text{CONH}_2$ groups which are almost neutral. So, polypeptides (IV), (VI), (VIII) and (IX) will be positively charged at pH = 7, *i.e.*, the correct integer is 4.

10. (2): The equations required are



Contd. from page no. 82

CHEMISTRY MUSING

PROBLEM SET 53

COMPREHENSION

A mixture of two aromatic compounds (A) and (B) was separated by dissolving in chloroform followed by extraction with aqueous KOH solution. The organic layer containing (A), when heated with alcoholic solution of KOH produce $\text{C}_7\text{H}_5\text{N}$ (C) associated with unpleasant odour.

7. What is A?
- (a) $\text{C}_6\text{H}_5\text{NH}_2$ (b) $\text{C}_6\text{H}_5\text{CH}_3$
 (c) $\text{C}_6\text{H}_5\text{CHO}$ (d) None of these
8. The alkaline aqueous layer (B) when heated with chloroform and then acidified give a mixture of isomeric compounds of molecular formula $\text{C}_7\text{H}_6\text{O}_2$. (B) is
- (a) $\text{C}_6\text{H}_5\text{CHO}$ (b) $\text{C}_6\text{H}_5\text{COOH}$
 (c) $\text{C}_6\text{H}_5\text{CH}_3$ (d) none of these.

INTEGER VALUE

9. When a certain metal was irradiated with light of frequency $4.0 \times 10^{16} \text{ s}^{-1}$, the photoelectrons emitted has three times the kinetic energy as the kinetic energy of photoelectrons emitted when the metal was irradiated with light of frequency $2.0 \times 10^{16} \text{ s}^{-1}$.

Milliequivalents of iodine = Milliequivalents of KI
 = Milliequivalents of O_3 reacted = $2 \times$ Milliequivalents of hypo

Milliequivalents of $\text{Na}_2\text{S}_2\text{O}_3 = 1.5 \times 0.01 = 1.5 \times 10^{-2}$

Millimoles of iodine = $\frac{1.5 \times 10^{-2}}{2} = 7.5 \times 10^{-3}$

[\because 'n' factor for iodine = 2]

Millimoles of ozone = 7.5×10^{-3}

Volume of ozone = $\frac{nRT}{P} = \frac{7.5 \times 10^{-6} \times 0.0821 \times 300}{1}$
 = $184.725 \times 10^{-6} \text{ L}$

Volume percentage of ozone = $\frac{184.725 \times 10^{-6}}{9} \times 100$
 = 2.05×10^{-3}

Thus, x is 2.




The critical frequency (ν_0) of the metal calculated is $x \times 10^{16} \text{ s}^{-1}$. Then, the value of x is

10. The resistance of N/10 solution is found to be 2.5×10^3 ohms. The equivalent conductance of the solution is (cell constant = 1.25 cm^{-1})



NEW LAUNCH




ONLINE TEST SERIES

5 Full Syllabus Mock Tests





AIMS RANK INDICATOR

Now on your android & IOS Smart phones with the same login of web portal.



Log on to <http://pcmbtoday.com>

Now, save up to Rs 2,020*



Subscribe to MTG magazines today.

Our 2017 offers are here. Pick the combo best suited for your needs. Fill-in the Subscription Form at the bottom and mail it to us today. If in a rush, log on to www.mtg.in now to subscribe online.

*On cover price of ₹ 30/- each.



About MTG's Magazines

Perfect for students who like to prepare at a steady pace, MTG's magazines-Physics For You, Chemistry Today, Mathematics Today & Biology Today-ensure you practice bit by bit, month by month, to build all-round command over key subjects. Did you know these magazines are the only source for solved test papers of all national and state level engineering and medical college entrance exams?

Trust of over 1 Crore readers since 1982.

- Practice steadily, paced month by month, with very-similar & model test papers
- Self-assessment tests for you to evaluate your readiness and confidence for the big exams
- Content put together by a team

- comprising experts and members from MTG's well-experienced Editorial Board
- Stay up-to-date with important information such as examination dates, trends & changes in syllabi
- All-round skill enhancement –

- confidence-building exercises, new studying techniques, time management, even advice from past JEE/NEET toppers
- **Bonus:** Exposure to competition at a global level, with questions from Intl. Olympiads & Contests

SUBSCRIPTION FORM

Please accept my subscription to:

Note: Magazines are despatched by Book-Post on 4th of every month (each magazine separately).

Tick the appropriate box.



PCMB combo

- 1 yr: ₹ 1,000 (save ₹ 440) 2 yr: ₹ 1,800 (save ₹ 1,080) 3 yr: ₹ 2,300 (save ₹ 2,020)

PCM combo

- 1 yr: ₹ 900 (save ₹ 180) 2 yr: ₹ 1,500 (save ₹ 660) 3 yr: ₹ 1,900 (save ₹ 1,340)

PCB combo

- 1 yr: ₹ 900 (save ₹ 180) 2 yr: ₹ 1,500 (save ₹ 660) 3 yr: ₹ 1,900 (save ₹ 1,340)

Individual magazines

Physics Chemistry Mathematics Biology

- 1 yr: ₹ 330 (save ₹ 30) 2 yr: ₹ 600 (save ₹ 120) 3 yr: ₹ 775 (save ₹ 305)

Enclose Demand Draft favouring **MTG Learning Media (P) Ltd.** payable at New Delhi. You can also pay via Money Orders. Mail this Subscription Form to Subscription Dept., **MTG Learning Media (P) Ltd.** Plot 99, Sector 44, Gurgaon – 122 003 (HR).

Want the magazines by courier; add the courier charges given below:

- 1 yr: ₹ 240 2 yr: ₹ 450 3 yr: ₹ 600

Tick the appropriate box.

Student Class XI XII Teacher Library Coaching

Name: _____

Complete Postal Address: _____

Pin Code Mobile #

Other Phone # 0

Email: _____



E-mail subscription@mtg.in. Visit www.mtg.in to subscribe online. Call (0)8800255334/5 for more info.


To be a NEET champion, you need help from a CHAMPION



mtg

Skill. Passion. Hard work and determination. As a student sitting for the highly competitive NEET, you need all that. However, only a few will win, very likely with the help of a champion coach.


MTG's Champion Series for NEET is just the coach you need. It will guide you in identifying what's important for success and what's not. And then help you check your readiness with its most comprehensive question bank. So you know your strengths and weaknesses right from the word go and course-correct accordingly. Put simply, MTG's Champion Series will help you manage your preparation effort for NEET for maximum outcome. The best part is you study at a pace you're comfortable with. Because it's all chapterwise, topicwise.



NCERT-based • Chapterwise • Topicwise • 10 years' solved previous test papers (all major medical entrance exams) • Concise summary at the start of each chapter for quick revision of key concepts • Analysis of importance of topics basis historical examination pattern • Test papers for self-assessment

Visit www.MTG.in to buy online.
Or visit a leading bookseller near you.
For more information, call 1800 300 23355 (toll-free) or 0124-6601200 today.
Email info@mtg.in

ABRACADABRA



More than magic words, you need help from a magician to pass JEE with flying colours. Which is why MTG has collaborated with Dr Ramaniah Gundala, the popular Maths professor from Chennai, to bring you the all-new Objective Mathematics for JEE Main & Advanced.

Includes
6 assignments for self-assessment
Previous years' JEE questions (2017-13)

Introductory price
₹ 750/-



Dr Ramaniah Gundala

After securing his DIIT and Ph.D from IIT Kharagpur, Dr Gundala was elected Fellow of National Academy of Sciences (FNASc). His 50+ years of teaching experience includes distinguished tenures at IIT Kharagpur and Anna University, Chennai. He has authored 7 books and published an astonishing 85 research papers. He's now retired and prepares students for success in IIT-JEE at two leading coaching institutes in Chennai and Warangal.

Highlights of MTG's Objective Mathematics for JEE Main & Advanced

- Well-structured theory covering summary of important concepts and easy-to-understand illustrations
- 5800+ questions
- Unique and brain-stimulating exercises including questions of the following types:
 - Single Correct Option
 - More Than One Correct Option
 - Assertion and Reason, Comprehension Type
 - Matrix Match
 - Integer Answer Type

Visit www.MTG.in to buy online or visit a leading bookseller near you.
For more information, e-mail info@mtg.in or call 1800 300 23355 (toll-free) today.

mtg

TRUST OF MILLIONS, SINCE 1982

Test Series & Question Bank JEE Advanced 2018

- Test Series for JEE Advanced (10 Mock Papers) Fees : ₹ 1700/-
 - Question Bank (2400 challenging problems) Fees : ₹ 2000/-
- A Compendium with all key formulae in Physics, Chemistry & Maths and memory tips for quick revision and Special Module on short cut methods will also be provided with the Test Series / Question Bank.

Online Test Series JEE Main 2018 / BITSAT 2018

- 10 Online Mock Papers with performance evaluation for JEE Main Fees : ₹ 1700/-
 - 10 Online Mock Papers with performance evaluation for BITSAT Fees : ₹ 1700/-
- Note: Paper based Test Series is also available for JEE Main 2018.

Test Series NEET 2018

- Test Series for NEET (10 Mock Papers) Fees : ₹ 1700/-
- A Compendium with all key formulae in Physics, Chemistry & Biology and memory tips for quick revision and Special Module on short cut methods will also be provided with the Test Series.

Self Study Series IIT JEE 2019

- Highlights of Self Study Series
- Comprehensive Theory illustrating core concepts & short-cut methods
 - Topic-wise & Full length Tests with performance evaluation
 - 15,000 challenging problems with solutions, Test-Series and Question Bank
 - Fees : ₹ 17,500/- (all inclusive)

Vidyalankar's Merit Concession Scheme for above Courses*

Merit Concession Category	Fees Payable (₹)		
	Test Series	Question Bank	Self Study Series
• Olympiad qualified students [Phy (NSEP)/Chem (NSEC)/Astronomy (NSEA)] • KVPY Scholarship holders	Nil	Nil	Nil
• Advanced 2017 Qualified students (Ranks upto Top 10000)	Nil	Nil	NA
• KVPY Level I qualified / NTSE Scholarship holders / Maths RMO qualified	Nil	Nil	5250/-
• NTSE Level I qualified students	850/-	1000/-	8750/-
• Students Scoring 98% and above in Maths and Science in Std. X Board Exam			
• Students Scoring 95% and above in Maths and Science in Std. X Board Exam			
• Students Scoring 90% and above in Maths and Science in Std. X Board Exam			









*Conditions Apply

To Enroll:

Send your application on the below address, stating your name, complete postal address, contact number, Email ID, course of your interest & year of appearing for IIT JEE along with necessary documents, one Passport size colour photograph & Money Order / Bank Draft payable at Mumbai, in the name of "Vidyalankar Classes and Publications"

For online payment visit : www.epay.vidyalankar.org

Some of Previous Years' IIT JEE Toppers

ARCHISMAN D.	ANIMESH B.	DHRUV SHAH	VILLY GOHIL	RANVIR RANA	MOHD. H.	NAVNEET M.	PRAVEEN S.
							
AIR-16* 2017	AIR-19 2016	AIR-111 2015	AIR-20* 2014	AIR-19 2013	AIR-20 2012	AIR-24 2011	AIR-10 2010

* Category Rank & many more...

Corporate Office : Vidyalankar (51), Pearl Center, Senapati Bapat Marg, Dadar (W), Mumbai - 400 028.

• Tel.: (022) 4232 42 32 / 2430 63 67 • Email : iit@vidyalankar.org / Web site : www.vidyalankar.org



IMPARTING QUALITY EDUCATION SINCE 1986

YOU HAVE A DREAM

WE WILL WALK WITH YOU TO REALISE IT

VIDYAMANDIR CLASSES (VMC) has been the pioneers in IIT JEE Preparation in India. Over the last three decades, we have helped thousands of students realize their IIT dream. Our philosophy of keeping the student's interests at the forefront and our unmatched passion for excellence is what separates VMC from the rest. It has been our constant endeavor to provide students with the best possible academic system including Best-in-Class

content and Testing system.

After a good research into the Pre Medical Entrance examination space in India, Vidyamandir Classes is now pleased to announce the launch of the Medical Test Preparation division from the upcoming session. The Medical division will help students prepare for PMT exams like AIIMS, NEET, AFMC and the like.

NATIONAL  **25TH & 31ST**
ADMISSION CUM
SCHOLARSHIP TEST **DECEMBER**

COURSES FOR MEDICAL PREPARATION Offered For Students Currently in Class 8TH, 9TH & 10TH

COURSES FOR IITJEE PREPARATION Offered For Students Currently in Class 8TH, 9TH, 10TH & 11TH

VMC DELHI NCR CENTRES (Centre for both JEE & MEDICAL preparation)

www.vidyamandir.com

+ H.O.: Pitampura
(011) 45221191-93

+ Anand Vihar
(011) 43068924/26


Dwarka
(011) 45221178

+ Faridabad
(0129) 4163119/20

Ghaziabad
9650495709

+ Greater Noida
7290045422

Gurugram
(0124) 4148541/42

 /vidyamandirclasses

+ Janakpuri
9643407598

+ Noida
7878480490

Punjabi Bagh
Information Centre
7290045419

+ Safdarjung Enclave
(011) 41046279/80

Kalu Sarai
Correspondence Wing
(011) 42156441

Vikasपुरी
9350902825

Rohtak
9999335300

 /Vidyamandir_in

ALSO OFFERING, INTENSIFIED CORRESPONDENCE COURSE & VMC ALL INDIA TEST SERIES (VATS). FOR MORE INFO, CALL (011) 42156441



READY. STEADY. VITEEE

EXPERIENCE EDUCATION
THAT'S AT PAR WITH TOP
UNIVERSITIES ABROAD



VITEEE 2018

4th April to 15th April (Computer Based Test)

Applications are invited for B.Tech Programmes

Biotech., Bioengg., Chemical, Civil, CSE, CSE (Specialisation in Digital Forensics & Cyber security), CSE (Bioinfo.), CSE (Data Analytics), CSE (Gaming Tech.), CSE (Information Security), CSE (Networks & Security), ECE, ECE (Biomedical), ECE (Embedded Systems), ECE (IOT), ECE (Sensors & Wearable Tech.), ECE (VLSI), EEE, EIE, Electronics & Computer Engg., IT, Mech., Mech (Auto), Mech. (Energy), Production & Industrial Engg., B.Des.Ind.Design

To apply online and to get details like eligibility and fee, please visit www.vit.ac.in

LAST DATE FOR SUBMISSION: 28th FEBRUARY 2018

ONLINE APPLICATION FEE: ₹1150

International Transfer Programme (ITP)



VIT offers students the choice to study the first two years at VIT University and the next two in the following universities abroad & get their degree

United States of America: • Purdue University • State University of New York (SUNY)
• Rochester Institute of Technology (RIT) • University of Massachusetts (UMass)
United Kingdom: • Queen Mary University of London **Australia:** • Australian National University • Deakin University • Queensland University of Technology • Royal Melbourne Institute of Technology (RMIT) • Curtin University

Opportunities Unlimited

VIT offers students, opportunity to get placed in companies with regular, dream (5L CTC & above), Superdream (10L CTC & above) offers. In 2016-17, 427 companies came for placement to VIT, which itself is a record, of which 45+ were Super Dream Companies. Recruiting companies include:

- | | | | | |
|------------|-----------|-------------|-------------------|-----------------|
| •Microsoft | •Amazon | •Maruti | •ABB | •Deloitte |
| •Deshaw | •Paypal | •Hyundai | •L & T | •Morgan Stanley |
| •Oracle | •Visa | •Mahindra | •ITC | •KPMG |
| •SAP | •Cisco | •Honda | •Shapoorji | •JP Morgan |
| •Vmware | •Barclays | •TVS Motors | •Johnson Controls | •PWC |



Ranked as 13th best engineering institution in India by NIRF, MHRD, GOI, for 2 consecutive years in 2016 & 2017

FFCSTM
FULLY FLEXIBLE CREDIT SYSTEM
Option to Change Branch
Opportunity to graduate with a double major
Choice of major and minor programmes
Degree with honours

GET THE VIT EDGE

- National Record in Slot 1 companies recruitment for 9 years in a row
- Record 7943 job offers from Slot 1 companies (Accenture, Cognizant, Infosys, TCS, Wipro) at VIT University for the year 2016
- Excellent international internship and placement opportunities

Advantage VIT

- Curriculum for Applied Learning (CAL)TM
- Project Based Learning (PBL)
- Opportunity for Semester Abroad Programme
- One module of every subject handled by Industry experts
- 35,000 students from 60 countries
- 110 student clubs and chapters
- More than 170 Adjunct Professors from reputed universities abroad

International Accreditations

- ABET, US, <http://www.abet.org> has accredited 14 B.Tech Programs at VIT Vellore & Chennai Campus
- 3 cycles of NAAC accreditation completed
- Awarded 'A' grade for last two consecutive cycles
- First University in India to get QS 4 Stars Rating



Offline application forms can be obtained: • From the designated branches of Post Offices on cash payment of ₹1200/- (list available on our website) • By handing over a DD for ₹1200/- drawn in favour of VIT University, payable at Vellore. Applications also available in VIT Chennai, VIT Bhopal & VIT AP.



For enquiries contact us on our helpline number +91-416-330 5555

For more details visit www.vit.ac.in

f VITUniversity | VIT_Univ



VIT[®]
VELLORE
CHENNAI
www.vit.ac.in



VIT[®]
BHOPAL
www.vitbhopal.ac.in



VIT[®]
AP
www.vitap.ac.in

A SCHOLARSHIP OPPORTUNITY

THAT YOU CAN'T AFFORD TO MISS



A UNIQUE OPPORTUNITY TO HELP YOU FULFIL YOUR DREAMS

For Class XII Studying / Passed Students

GET UP TO 100% SCHOLARSHIP*

Every day is a new opportunity and every opportunity a new chance. Aakash brings ATQ, a fantastic opportunity for those students who believe they can & they will. It is a chance to prove their potential and build a better tomorrow. With ATQ, students can win scholarships up to 100% in Short Term Courses for 2018 such as Crash Courses / Test Series Courses & cash awards. It provides a platform to students nationwide who have been waiting for the right opportunity to embark on a successful journey ahead.

*Terms & Conditions apply.

Date of Examination: **24th December, 2017 (Sunday)**

Exam Fee: ₹ 300/-

Test Centres: **All Aakash Centres**

paytm
Pay Fees
on Paytm!



Last Date of receiving duly filled enrollment form by post / by hand / through online - **17th December, 2017** (Sunday) and with late fee - **19th December, 2017** (Tuesday)

Download Enrollment Form from our website: www.aakash.ac.in or www.myatq.in The result shall be displayed on our website.



Aakash

Medical | IIT-JEE | Foundations

(Divisions of Aakash Educational Services Pvt. Ltd.)

www.myatq.in | www.aakash.ac.in

TOLL FREE: **1800-180-2727**

GIVE A MISSED CALL: **9599280605**

SMS Aakash to **53030**

Registered Office : Aakash Tower, 8, Pusa Road, New Delhi-110005. Ph.: (011) 47623456 | E-mail: atq@aeslin | iitjee@aeslin | medical@aeslin

[instagram.com/aakasheducation](https://www.instagram.com/aakasheducation)

[facebook.com/aakasheducation](https://www.facebook.com/aakasheducation)

[youtube.com/AakashEducation](https://www.youtube.com/AakashEducation)

twitter.com/aakash_twitted

aakashinstitute.blogspot.com