

ACE YOUR WAY CBSE Class XI-XII

NEET | JEE Class XI-XII
ESSENTIALS

CHEMISTRY

India's #1
CHEMISTRY MONTHLY FOR
JEE (Main & Advanced) & NEET

 **today**

**EXAMINER'S
MIND**

**MONTHLY
PRACTICE
PROBLEMS
(XI & XII)**

**CONCEPT
BOOSTER**

**CHEMISTRY
MUSING**

mtg

Trust of more than
1 Crore Readers
Since 1982



2017100008707

CONCEPT MAP



What Makes ALLEN

INDIA's Most Trusted Career Coaching Institute...

- ✓ 29 Years of Leadership
- ✓ Trust of 6.50 Lac+ Students & their Parents
- ✓ 4500+ Team members including 350+ IITians and 70+ Doctors
- ✓ Unmatched Education System with Indian Values
- ✓ With Maximum Talented Students joining ALLEN, it becomes a National Level Competition provider
- ✓ Only Institute giving Quality Results in both Pre-Medical & Pre-Engineering year after year
- ✓ All India Ranks 1, 2, 3 in both IIT-JEE (Adv.) & NEET 2016 from Classroom, Breaking all Records
- ✓ Limca Book of Records (2014) recognized ALLEN as Largest Educational (coaching) Institute of India by student strength (66,504) at a single location KOTA

AIIMS 2017

FIRST TIME IN THE HISTORY

All Top 10 AIR Secured by Students of ALLEN
34 in Top 50 AIR



NISHITA PUROHIT
Classroom

NEET 2017

6 in Top 10 AIR
54 in Top 100 NEET OVERALL RANKS



ARCHIT GUPTA
Classroom

MANISH MULCHANDANI
Classroom

ABHISHEK DOGRA
Classroom

IIT-JEE 2017

25 in Top 100 AIR
130 in Top 500 AIR



ONKAR M.D.
Distance

RACHIT BANSAL
Classroom

LAKSHAY SHARMA
Classroom



KOTA | AHMEDABAD | BENGALURU | BHILWARA | CHANDIGARH | INDORE
JAIPUR | MUMBAI | RAJKOT | RANCHI | SURAT | VADODARA

Corporate Office : "SANKALP", CP-6, Indra Vihar, Kota (Rajasthan)-324005, India

WANT TO KNOW MORE,
COME TO **ALLEN** AND EXPERIENCE THE LEADERSHIP

allen.ac.in | dlp.allen.ac.in | tab.allen.ac.in

info@allen.ac.in

+91-744-5156100

JEE (Advanced) | JEE (Main) | NEET-UG/AIIMS | KVPY, NTSE & Olympiads | Class 6th to 12th

To be a NEET champion, you need help from a CHAMPION



mtg

Skill. Passion. Hard work and determination. As a student sitting for the highly competitive NEET, you need all that. However, only a few will win, very likely with the help of a champion coach.

MTG's Champion Series for NEET is just the coach you need. It will guide you in identifying what's important for success and what's not. And then help you check your readiness with its most comprehensive question bank. So you know your strengths and weaknesses right from the word go and course-correct accordingly. Put simply, MTG's Champion Series will help you manage your preparation effort for NEET for maximum outcome. The best part is you study at a pace you're comfortable with. Because it's all chapterwise, topicwise.



NCERT-based • Chapterwise • Topicwise • 10 years' solved previous test papers (all major medical entrance exams) • Concise summary at the start of each chapter for quick revision of key concepts • Analysis of importance of topics basis historical examination pattern • Test papers for self-assessment

Visit www.MTG.in to buy online.
Or visit a leading bookseller near you.
For more information, call 1800 300 23355 (toll-free) or 0124-6601200 today.
Email info@mtg.in

A Premier Institute in INDIA
Integrated PUC, +1 & +2 | NEET | IIT-JEE | AIIMS | JIPMER | EAMCET

EXCEL ACADAMICS®

Bengaluru

IIT-Main & Advanced | NEET | EAMCET | BITS | AIIMS | JIPMER | CET | COMED-K

REGISTRATION OPEN for PUC /12th /+2 Passed Students



ONE YEAR LONG TERM MEDICAL COURSE

(PCB) (Repeaters Course)

NEET - 2018

The only Institute in INDIA successfully conducting "Daily Test" for NEET (Medical) aspirants.

Features of One Year Long Term Medical Programme

- Substantial Medical Entrance Coaching at EXCEL ACADAMICS done by the eminent faculty those who are Medical Guru's in India.
- Starting with very basics and Strengthening them at Fundamentals during coaching.
- Maintaining high end competent atmosphere.
- Timely completion of Syllabus and perfect revision.
- Special focus on Physics Numerical questions.
- Exclusive A.C. Campus with LCD projectors for One Year Long Term Medical Programme.
- Daily 6 (Six) hours PCB teaching followed by counselling hours and doubt clarification session.
- Targeting for practicing 3 (Three) Lakh MCQ's Physics, Chemistry & Biology in this programme.
- Separate Weekend Examination for NEET, JIPMER and AIIMS.
- Updating to the parents through SMS/email regarding student's attendance, rank and their performance on regular basis.
- Coaching at Excel Academics will be helpful to attend All National Medical Entrance Examinations like NEET, AIIMS, JIPMER etc... along with CET, EAMCET, Kerala PMT, MHCET and COMED-K.
- 24 x 7 Library facility with PCB Medical Entrance Study Materials and Magazines.
- Brand new high security hostel facility for Boys and Girls.
- Periodical Parent, Teacher, Student Meeting.
- Semi-residential facility.
- Lowest fee structure including all taxes.

Targeting **3 Lakhs MCQ's** for **NEET 2018**

Help Line No.
76769 17777
76764 16666

Registration Started for Integrated PUC, CET, NEET and IIT-JEE coaching



LAUNCHING E-40 for sure success in **NEET - 2018**

EXCEL ACADAMICS, Opp: Peoples Tree Hospital, Sector A, Yalahanka New Town, Bangalore - 560064, KARNATAKA

★★ **Contact : 9535656277, 9880155284, 9036357499** ★★

Separate Deluxe Hostel for Boys and Girls

CBSE CHAMPION Chapterwise -Topicwise Solved Papers



CBSE CHAMPION Chapterwise -Topicwise Solved Papers Series contains topicwise questions and solutions asked over last decade in CBSE-Board examination.

Questions are supported with topicwise graphical analysis of previous years CBSE Board questions as well as comprehensive and lucid theory. The questions in each topic have been arranged in descending order as per their marking scheme. Questions from Delhi, All India, Foreign and Compartment papers are included. This ensures that all types of questions that are necessary for Board exam preparation have been covered.


Important feature of these books is that the solutions to all the questions have been given according to CBSE marking scheme. CBSE sample paper and practice papers are also supplemented.

Examination papers for Class-10 and 12 Boards are based on a certain pattern. To excel, studying right is therefore more important than studying hard, which is why we created this series.

Visit
www.mtg.in
 for latest offers
 and to buy
 online!



Available at all leading book shops throughout India.
 For more information or for help in placing your order:
 Call 0124-6601200 or email info@mtg.in



**YOUR SUCCESS JOURNEY
TO KEEP AHEAD.**

AIR-1

Cash Prize ₹1,50,000

11 Prizes
(Gold Medal+Trophy)

AIR-2

Cash Prize ₹1,00,000

11 Prizes
(Silver Medal+Trophy)

AIR-3

Cash Prize ₹75,000

11 Prizes
(Bronze Medal+Trophy)

For Ambitious Students of Class 6th to 12th

LAST DATE OF REGISTRATION

**Sunday
10 September 2017**

DATE OF EXAMINATION

**Sunday
01 October 2017**

Registration Fee: ₹ 150 | **4200 prizes worth ₹2 crore**

Total prizes for each Category 150 | Total Categories 11

For more Details & Registration visit: www.resostart.in

RESONANCE EDUVENTURES LTD.

Corporate Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Rajasthan)- 324005

Registered Office : J-2, Jawahar Nagar Main Road, Kota (Rajasthan)- 324005 | Tel. No.: 0744-3012100, 3012222, 6635555 | CIN: U80302RJ2007PLC024029

To know more: sms RESO <space> START at 56677 | E-mail: start@resonance.ac.in

Toll Free: 1800 258 5555



facebook.com/ResoStArT/



twitter.com/ResonanceEdu



www.youtube.com/c/ResoStArTVideo



blog.resonance.ac.in

CHEMISTRY today

Volume 26 No. 9 September 2017

Managing Editor

Mahabir Singh

Editor

Anil Ahlawat

Corporate Office:

Plot 99, Sector 44 Institutional area, Gurgaon -122 003 (HR).

Tel : 0124-6601200 e-mail : info@mtg.in website : www.mtg.in

Regd. Office:

406, Taj Apartment, Near Safdarjung Hospital, New Delhi - 110029.

CONTENTS

Class 11

NEET JEE Essentials	8
Examiner's Mind	20
Ace Your Way CBSE	29
MPP-5	36
Concept Map	46

Class 12

NEET JEE Essentials	40
Concept Map	47
Examiner's Mind	54
Ace Your Way CBSE	63
MPP-5	72

Competition Edge

Chemistry Musing Problem Set 50	76
JEE Advanced Practice Problems	77
Concept Booster	79
Advanced Chemistry Bloc	81
Chemistry Musing Solution Set 49	83
Crossword	85

Subscribe online at www.mtg.in

	Individual Subscription Rates			Combined Subscription Rates			
	1 yr.	2 yrs.	3 yrs.		1 yr.	2 yrs.	3 yrs.
Mathematics Today	330	600	775	PCM	900	1500	1900
Chemistry Today	330	600	775	PCB	900	1500	1900
Physics For You	330	600	775	PCMB	1000	1800	2300
Biology Today	330	600	775				

Send D.D./M.O in favour of MTG Learning Media (P) Ltd.

Payments should be made directly to : MTG Learning Media (P) Ltd,

Plot No. 99, Sector 44, Gurgaon - 122003 (Haryana)

We have not appointed any subscription agent.

Owned, Printed and Published by MTG Learning Media Pvt. Ltd. 406, Taj Apartment, New Delhi - 29 and printed by HT Media Ltd., B-2, Sector-63, Noida, UP-201307. Readers are advised to make appropriate thorough enquiries before acting upon any advertisements published in this magazine. Focus/Infoocus features are marketing incentives. MTG does not vouch or subscribe to the claims and representations made by advertisers. All disputes are subject to Delhi jurisdiction only.

Editor : Anil Ahlawat

Copyright © MTG Learning Media (P) Ltd.

All rights reserved. Reproduction in any form is prohibited.

NEET | JEE

ESSENTIALS

Class
XI


Maximise your chance of success, and high rank in NEET, JEE (Main and Advanced) by reading this column. This specially designed column is updated year after year by a panel of highly qualified teaching experts well-tuned to the requirements of these Entrance Tests.

Unit 3

States of Matter : Gases & Liquids | Thermodynamics


STATES OF MATTER : GASES & LIQUIDS

GASEOUS STATE OF MATTER



Study right. Dig deep.

Build a solid foundation for success
in JEE (Main)



Are you a do-it-yourself type of a student? Then for success in JEE Main, choose MTG's JEE (Main) combo, comprising coursebooks for Physics, Chemistry & Mathematics. This combo is all class 11 and 12 students need for a solid and deep understanding of concepts in these three key subjects.

FEATURES:

- Based on latest pattern of JEE (Main)
- Full of graphic illustrations & MCQs for deep understanding of concepts
- Covers the entire syllabus
- NCERT Drill MCQs framed from NCERT Books
- 11 Years (2017-2007) Previous Years MCQs of JEE Main / AIEEE
- 2017 JEE Main (Offline & Online) Solved Paper included




Scan now with your
smartphone or tablet
Application to read
QR codes required

Note: Coursebooks are also available separately.

Available at all leading book shops throughout India. To buy online visit www.mtg.in.

For more information or for help in placing your order, call 0124-6601200 or e-mail: info@mtg.in



DALTON'S LAW OF PARTIAL PRESSURES

- The total pressure exerted by a mixture of two or more non-reacting gases enclosed in a vessel is equal to the sum of the individual partial pressures which each gas would exert when present alone in the same vessel at the same temperature.

$$P_{\text{total}} = P_1 + P_2 + P_3 + \dots + P_n$$

$$p_1 = x_1 P_{\text{total}}$$

x_1 = Mole fraction of the gas 1.

GRAHAM'S LAW OF DIFFUSION / EFFUSION

- Under similar conditions of temperature and pressure, the rates of diffusion/effusion of different gases are inversely proportional to the square root of their densities.

$$\frac{r_1}{r_2} = \sqrt{\frac{d_2}{d_1}}$$

where d_1, d_2 are densities of two gases.

$$\frac{r_1}{r_2} = \frac{P_1}{P_2} \sqrt{\frac{d_2}{d_1}} = \frac{P_1}{P_2} \sqrt{\frac{M_2}{M_1}}$$

$$\frac{r_1}{r_2} = \sqrt{\frac{T_1 d_2}{T_2 d_1}} = \sqrt{\frac{T_1 M_2}{T_2 M_1}}$$

KINETIC MOLECULAR THEORY OF GASES

- The molecules are separated from one another by large distances.
- Molecular collisions are perfectly elastic.
- There are no forces of interactions between the molecules.

- Kinetic gas equation, $PV = \frac{1}{3} mn\bar{u}^2$

where, P = Pressure exerted by the gas,

m = Mass of each molecule of gas,

n = Total number of molecules of the gas present in volume V ,

\bar{u} = Root mean square speed of the gas

$$\text{K.E., } E_k = \frac{3}{2} nRT$$

$$\text{Average kinetic energy, } \bar{E}_k = \frac{1}{2} m\bar{u}^2 = \frac{3}{2} \frac{RT}{N_A}$$

where, N_A = Avogadro's constant

- The average kinetic energy of gas molecules is directly proportional to the absolute temperature of the gas.
- All gases are made up of a very large number of molecules.



How to choose the right answer, fast?



The answer is practice...

Our team has seen that in NEET, AIIMS, JIPMER and JEE, Multiple Choice Questions (MCQs) are based on the NCERT syllabus. Largely!! With Objective NCERT at your FINGERTIPS, you can become a pro at handling MCQs. Practice to increase your accuracy and improve timing with a bank of over 15,000 questions, all framed from NCERT course books. Don't take our word, have a look what some of our readers have to say...

Features:

- Chapterwise student-friendly synopses for quick-and-easy revision
- Topicwise MCQs to check your progress
- NCERT Exemplar MCQs
- Assertion & Reason questions for an edge in your AIIMS/JEE preparation
- HOTS MCQs to boost your concepts
- 6 Practice papers for self-assessment

Sanjay Shankar says, "Awesome book!! Everything is just perfect and the collaboration of the 11th and 12th std. just made it easier for us and with this less price. I will definitely recommend this book for every NEET preparing student."

Shweta says, "Must read for good score in NEET. Many questions in NEET are from this book in last 3 years. It also covers outside NCERT topics. Nice book."

Vijay says, "This book is ideal for practising MCQs (chapterwise). It appreciably covers all the important as well as less important questions. HOTS and sample question papers are provided as well. No demerits of the book can be listed. Though, it is not light weighted and thus cannot be carried, you wouldn't get bored revising each chapter from the revision section and then answering the questions. The language is appropriate and lucid as well as easy to understand."

S.J. Uday says, "It is an awesome book. Firstly I was scared how it will be, but after having it, I was amazed. One must have this book who is interested in going for the NEET examination."

Sonal Singh says, "Book is very good. As it contains all the topicwise questions from every topic of NCERT, one can develop a question solving ability and also understand the basic concepts."

Sunehri says, "This book contains over 150 MCQs in each chapter, has categories like MCQs, NCERT, HOTS based questions, AIIMS assertion reasoning questions. Every chapter gives a short summary of chapter. Great book for entrance exams like NEET, AIIMS etc."

Prashant says, "The book is really awesome. It makes you cover up whole NCERT in a simple way. Solving the problems can increase your performance in exam. I would suggest each & every NEET candidate to solve the book. The book is also error free; not like other publications books which are full of errors."

Arka says, "It is a nice question bank of NCERT. I think it is the best of its kind. The book is a must to prepare for NEET."



Scan now with your smartphone or tablet

Application to read QR codes required




MTG Learning Media (P) Ltd.
Plot #99, Sector 44, Gurgaon – 122 003 (HR)

Available at all leading book shops throughout India.

For more information or for help in placing your order,
Call 0124-6601200 or e-mail: info@mtg.in

MAXWELL-BOLTZMANN DISTRIBUTION OF MOLECULAR SPEEDS

- The plot between the fraction of molecules ($\Delta N/N$) possessing particular speeds against their corresponding speeds at a particular temperature is known as Maxwell's distribution curve.



$$u_{mp} : u_{av} : u_{rms} = 1 : 1.128 : 1.224$$

DEVIATION FROM IDEAL GAS BEHAVIOUR

- Real gases do not follow ideal gas equation perfectly at all conditions of temperatures and pressures.

- $\frac{PV}{nRT} = Z = \text{Compressibility factor}$

→ For an ideal gas, $Z = 1$

→ For a real gas :

$Z > 1$ (Positive deviation)

$Z < 1$ (Negative deviation)

- Equation of state for real gases (van der Waals' equation) :**

$$\left(P + \frac{an^2}{V^2} \right) (V - nb) = nRT$$

where, a and b are van der Waals' constants.

LIQUEFACTION OF GASES

- Gases can be liquefied by decreasing temperature or increasing pressure.
- Critical pressure :** The minimum pressure required to liquefy a gas at its critical temperature.

$$P_c = \frac{a}{27b^2}$$

- Critical volume :** Volume occupied by one mole of a gas at its critical temperature and pressure.

$$V_c = 3b$$

- Critical temperature :** A temperature above which a gas cannot be liquefied however high pressure may be applied on the gas.

$$T_c = \frac{8a}{27Rb}$$

LIQUID STATE OF MATTER

- Liquids have definite volumes but not definite shapes.

Properties of Liquid State

- Boiling point :** Temperature at which the vapour pressure of a liquid becomes equal to the atmospheric pressure.
- Surface tension :** Force per unit length acting perpendicular to the tangential line drawn at the surface of liquid is known as surface tension.

Surface Tension

Surface tension causes

Rise of a liquid in a capillary tube.

Spherical shape of a drop

Liquid wet things.

Unit : N m^{-1} or dyne cm^{-1}

Surface tension decreases with increase in temperature.

Greater the molecular forces, higher is the surface tension.

- Viscosity :** Internal resistance to the flow in liquids.

$$F = \eta A \frac{dv}{dx}, \text{ where, } A = \text{Area,}$$

$$\frac{dv}{dx} = \text{Velocity gradient,}$$

$\eta = \text{Coefficient of viscosity,}$

INFOSHOTS

New state of matter !

A new form of matter, a supersolid, which combines the properties of solids with those of superfluids. By using lasers to manipulate a superfluid gas known as Bose-Einstein condensate was coaxed into a quantum phase of matter that has a rigid structure-like a solid and can flow without viscosity (like superfluids).

THERMODYNAMICS

- A specified part of the universe which is under thermodynamic consideration is called the system.


THEMODYNAMICS TERMS

- **Extensive properties** : Properties of the system whose value depends upon the amount or size of the substance present in the system *e.g.*, Gibbs free energy, enthalpy, entropy, internal energy, etc.
- **Intensive properties** : Properties of the system whose value is independent of the amount or size of the substance present in the system but depends only upon the nature of the substance *e.g.*, temperature, pressure, viscosity, etc.
- **State function** : A physical quantity is said to be a state function if its value depends only upon the


state of the system and is independent of the path by which the state is reached *e.g.*, volume, temperature, internal energy, enthalpy.

- **Path function** : A physical quantity is a path function if it depends upon the path by which the change is brought about *e.g.*, work and heat.

Different Types of Systems



Types of Processes



FIRST LAW OF THERMODYNAMICS

Energy can neither be created nor destroyed, although it can be converted to one form to another form. The total energy of the universe remains constant.

$$\Delta U = q + w$$

where, ΔU = Change in internal energy;
 q = heat change; w = work done

Sign Conventions

$q = +ve$ (when heat is absorbed by the system)
 $q = -ve$ (heat is evolved by the system)
 $w = +ve$ (work is done on the system)
 $w = -ve$ (work is done by the system)

Work

- For isothermal reversible expansion,

$$\begin{aligned} q = -w &= p_{ext}(V_f - V_i) \\ &= nRT \ln \frac{V_f}{V_i} \\ &= 2.303 nRT \log \frac{V_f}{V_i} \\ &= 2.303 nRT \log \frac{P_i}{P_f} \end{aligned}$$

- For adiabatic reversible process, $q = 0$


$$\Delta U = w_{ad} = nC_v \Delta T = \frac{nR}{\gamma - 1} \Delta T$$

Enthalpy change (ΔH)

Heat absorbed or evolved (ΔH) during a reaction at constant pressure.

Internal energy change (ΔU)

The change in internal energy (ΔU) is the heat absorbed or evolved at constant volume.



HEAT CAPACITY

It is the amount of heat required to raise the temperature of the system through 1°C

$$C = q/\Delta T$$

Heat capacity at constant volume (C_v)

$C_v = 3R/2$ (For monoatomic gas)
 $C_v = 5R/2$ (For diatomic gas)

Heat capacity at constant pressure (C_p)


$C_p = 5R/2$ (For monoatomic gas)
 $C_p = 7R/2$ (For diatomic gas)

$$C_p/C_v = \gamma$$

$$C_p = C_v + nR$$

Hess's Law

- The enthalpy change in a particular reaction is the same whether the reaction takes place in one step or in a number of steps.



- Enthalpy of reaction = (Sum of bond enthalpies of reactants) – (Sum of bond enthalpies of products)
- Enthalpy of reaction = (Sum of enthalpies of products) – (Sum of enthalpies of reactants)

ENTROPY

- The property of a system which measures the degree of disorder or randomness in the system.

$$\Delta S = \frac{q_{rev}}{T}$$

$$\Delta S = nC_v \ln \frac{T_2}{T_1} + nR \ln \frac{V_2}{V_1} = nC_p \ln \frac{T_2}{T_1} + nR \ln \frac{P_1}{P_2}$$

- For isothermal process, $\Delta S_f = nC_v \ln \frac{V_2}{V_1} = nC_p \ln \frac{P_1}{P_2}$
- For isochoric process, $\Delta S_v = nC_v \ln \frac{T_2}{T_1}$
- For isobaric process, $\Delta S_p = nC_p \ln \frac{T_2}{T_1}$
- $\Delta_{fusion}S^\circ = \frac{\Delta_{fusion}H^\circ}{T_f}$; $\Delta_{vap}S^\circ = \frac{\Delta_{vap}H^\circ}{T_b}$
- Second law of thermodynamics** : The entropy of the universe is continuously increasing in the course of every spontaneous change.

GIBBS FREE ENERGY

- $G = H - TS$, where, G = Gibbs free energy
- $\Delta G = \Delta H - T\Delta S$
- Units** : kJ/mol, J/mol
- $\Delta G^\circ = \sum \Delta_f G^\circ_{(products)} - \sum \Delta_f G^\circ_{(reactants)}$
- $\Delta G^\circ = -2.303 RT \log K_{eq}$
- $\Delta G = RT \ln \frac{V_1}{V_2} = RT \ln \frac{P_2}{P_1}$

Gibbs Free Energy

$\Delta G = -ve$ Spontaneous	$\Delta G = 0$ At equilibrium	$\Delta G = +ve$ Non-spontaneous
---------------------------------	----------------------------------	-------------------------------------

- If $\Delta H = +ve$; $\Delta S = +ve \Rightarrow \Delta G = -ve$ if $T\Delta S > \Delta H$ (At high temperature)
- If $\Delta H = -ve$, $\Delta S = +ve \Rightarrow \Delta G = -ve$ (at all temperatures)
- If $\Delta H = -ve$, $\Delta S = -ve \Rightarrow \Delta G = -ve$ if $\Delta H > T\Delta S$ (at low temperature)

THIRD LAW OF THERMODYNAMICS

- At absolute zero, the entropy of a perfectly crystalline substance is taken as zero. $S_0 = 0$
- The most important application of the third law of thermodynamics is that it helps in the calculation of the absolute entropies of the substances at room temperature (or at any temperature T). These determinations are based upon the heat capacity measurements.
- Residual entropy** : The entropy possessed by a substance at absolute zero is called its residual entropy.

SPEED PRACTICE

- The heat of combustion of benzene in a bomb calorimeter (*i.e.* at constant volume) was found to be $3263.9 \text{ kJ mol}^{-1}$ at 25°C . Calculate the heat of combustion of benzene (in kJ mol^{-1}) at constant pressure.
 - -3263.9
 - -3267.6
 - -3260.2
 - $+436.1$
- Equal moles of hydrogen and oxygen gases are placed in a container with a pin-hole through which both can escape. What fraction of the oxygen escapes in the time required for one-half of the hydrogen to escape?
 - $3/8$
 - $1/2$
 - $1/8$
 - $1/4$

(NEET Phase-I 2016)
- Calculate the heat required to raise the temperature of 60.0 g of aluminium from 35 to 55°C . (Molar heat capacity of Al is $24 \text{ J mol}^{-1} \text{ K}^{-1}$.)
 - 1.07 kJ
 - 1.57 kJ
 - 1.07 J
 - 1.57 J
- A gaseous mixture of 2 moles of A, 3 moles of B, 5 moles of C and 10 moles of D is contained in a vessel. Assuming that gases are ideal and the partial pressure of C is 1.5 atm , the total pressure is
 - 3 atm
 - 6 atm
 - 9 atm
 - 15 atm
- The heat of reaction, $\text{N}_2 + 3\text{H}_2 \longrightarrow 2\text{NH}_3$, at 27°C was found to be -21.976 kcal . What will be the heat of the reaction at 50°C ? (The molar heat capacities at constant pressure for nitrogen, hydrogen and ammonia are 6.8 , 6.77 and $8.86 \text{ cal mol}^{-1} \text{ degree}^{-1}$ respectively.)
 - -22.192 kcal
 - -24.132 kcal
 - -21.760 kcal
 - -21.976 kcal
- What will be the pressure of the gaseous mixture when 0.5 L of H_2 at 0.8 bar and 2.0 L of oxygen at 0.7 bar are introduced in 1 L vessel at 27°C ?
 - 1.8 bar
 - 0.4 bar
 - 1.4 bar
 - 1.5 bar
- Calculate ΔH at 85°C for the reaction, $\text{Fe}_2\text{O}_3(\text{s}) + 3\text{H}_2(\text{g}) \longrightarrow 2\text{Fe}(\text{s}) + 3\text{H}_2\text{O}(\text{l})$.
Given : $\Delta H^\circ_{(298 \text{ K})} = -33.29 \text{ kJ/mol}$ and

Substance	$\text{Fe}_2\text{O}_3(\text{s})$	$\text{Fe}(\text{s})$	$\text{H}_2\text{O}(\text{l})$	$\text{H}_2(\text{g})$
$C_p^\circ (\text{J/K-mol})$	103.8	25.1	75.3	28.8

 - -28.14 kJ/mol
 - 35.9 kJ/mol
 - -303.29 kJ/mol
 - -25.1 kJ/mol
- The average energy per molecule of a gas at a given temperature T , is given by
 - $\frac{3}{2} RT$
 - $\sqrt{\frac{3RT}{M}}$
 - $\sqrt{\frac{8(R/N_A)T}{\pi M}}$
 - $\frac{3}{2} \left(\frac{R}{N_A} \right) T$
- What will be the entropy change for the reaction, $2\text{H}_2(\text{g}) + \text{O}_2(\text{g}) \longrightarrow 2\text{H}_2\text{O}(\text{l})$ at 300 K ? (Standard entropies of $\text{H}_2(\text{g})$, $\text{O}_2(\text{g})$ and $\text{H}_2\text{O}(\text{l})$ are 126.6 , 201.20 and $68.0 \text{ J K}^{-1} \text{ mol}^{-1}$ respectively.)
 - $-318.4 \text{ J K}^{-1} \text{ mol}^{-1}$
 - $318.4 \text{ J K}^{-1} \text{ mol}^{-1}$
 - $31.84 \text{ J K}^{-1} \text{ mol}^{-1}$
 - None of these
- For a given reaction, $\Delta H = 35.5 \text{ kJ mol}^{-1}$ and $\Delta S = 83.6 \text{ J K}^{-1} \text{ mol}^{-1}$. The reaction is spontaneous at (Assume that ΔH and ΔS do not vary with temperature.)
 - $T > 425 \text{ K}$
 - all temperatures
 - $T > 298 \text{ K}$
 - $T < 425 \text{ K}$

(NEET 2017)
- Given : $\text{C} + 2\text{S} \longrightarrow \text{CS}_2$; $\Delta H = 117 \text{ kJ mol}^{-1}$
 $\text{C} + \text{O}_2 \longrightarrow \text{CO}_2$; $\Delta H = -393 \text{ kJ mol}^{-1}$
 $\text{S} + \text{O}_2 \longrightarrow \text{SO}_2$; $\Delta H = -297 \text{ kJ mol}^{-1}$
 The heat of combustion of CS_2 to form CO_2 and SO_2 is
 - $-1104 \text{ kJ mol}^{-1}$
 - 1104 kJ mol^{-1}
 - $+807 \text{ kJ mol}^{-1}$
 - -807 kJ mol^{-1}
- Calculate the root mean square speeds of nitrogen at N.T.P.
 - $493 \times 10^4 \text{ m/s}$
 - 493.13 m/s
 - $493 \times 10^2 \text{ m/s}$
 - $493 \times 10^3 \text{ m/s}$

13. An irreversible process taking place at constant P and T , in which only pressure-volume work is being done by the system, the change in Gibbs energy (dG) and change in entropy (dS), satisfy the criterion
- $(dS)_{V,U} = 0, (dG)_{T,P} = 0$
 - $(dS)_{V,U} = 0, (dG)_{T,P} = +ve$
 - $(dS)_{V,U} = -ve, (dG)_{T,P} = -ve$
 - $(dS)_{V,U} = +ve, (dG)_{T,P} = -ve$
14. At 27 °C, 500 mL of helium diffuses in 30 minutes. What is the time (in hours) taken for 1000 mL of SO_2 to diffuse under the same experimental conditions?
- 240
 - 3
 - 2
 - 4
15. 4 g of graphite is burnt in a bomb calorimeter of heat capacity of 30 kJ K^{-1} in excess of oxygen at 1 atmospheric pressure. The temperature rises from 300 K to 304 K. What is the enthalpy of combustion of graphite (in kJ mol^{-1})?
- 360
 - 1440
 - 360
 - 1440
16. The r.m.s. velocity of hydrogen is $\sqrt{7}$ times the r.m.s. velocity of nitrogen. If T is temperature of the gas then
- $T_{(H_2)} = T_{(N_2)}$
 - $T_{(H_2)} > T_{(N_2)}$
 - $T_{(H_2)} < T_{(N_2)}$
 - $T_{(H_2)} = \sqrt{7} T_{(N_2)}$
17. Among the following, the incorrect statement is
- at very large volume, real gases show ideal behaviour
 - at very low temperature, real gases show ideal behaviour
 - at Boyle's temperature, real gases show ideal behaviour
 - at low pressure, real gases show ideal behaviour
- (JEE Main Online 2017)**
18. The volume of 0.0168 mol of O_2 obtained by decomposition of $KClO_3$ and collected by displacement of water is 428 mL at pressure 754 mmHg at 25 °C. The pressure of water vapour at 25 °C is
- 18.5 mm Hg
 - 20.6 mm Hg
 - 22.3 mm Hg
 - 24.6 mm Hg
19. In which reaction, ΔS is positive?
- $H_2O_{(l)} \longrightarrow H_2O_{(s)}$
 - $3O_{2(g)} \longrightarrow 2O_{3(g)}$
 - $H_2O_{(l)} \longrightarrow H_2O_{(g)}$
 - $N_{2(g)} + 3H_{2(g)} \longrightarrow 2NH_{3(g)}$
20. The pressure of a fixed amount of an ideal gas is 0.75 $N\ m^{-2}$. What will be the pressure after the volume of the gas is tripled and the absolute temperature is doubled?
- 0.75 $N\ m^{-2}$
 - 0.25 $N\ m^{-2}$
 - 0.50 $N\ m^{-2}$
 - 1.00 $N\ m^{-2}$
21. The following data are available for the melting of KCl: $\Delta H_{fus} = 7.25\ kJ\ mol^{-1}$ and $\Delta S_{fus} = 0.007\ kJ\ K^{-1}\ mol^{-1}$. Calculate the melting point of KCl.
- 1035.7 K
 - 762.7 K
 - 897.5 K
 - 1308.7 K
22. For gaseous state, if most probable speed is denoted by c^* , average speed by \bar{c} and root mean square speed by c , then for a large number of molecules the ratio of these speeds are
- $c^* : \bar{c} : c = 1 : 1.128 : 1.224$
 - $c^* : \bar{c} : c = 1 : 1.224 : 1.128$
 - $c^* : \bar{c} : c = 1.224 : 1.228 : 1$
 - $c^* : \bar{c} : c = 1.128 : 1.224 : 1$
- (JEE Main 2013)**
23. A reaction, $A + B \longrightarrow C + D + q$ is found to have a positive entropy change. The reaction will be
- possible at high temperature
 - possible only at low temperature
 - not possible at any temperature
 - possible at any temperature.
24. Capillary action of the liquid can be explained on the basis of its
- resistance to flow
 - surface tension
 - heat of vapourisation
 - refractive index.
25. For the reaction, $Ag_2O_{(s)} \longrightarrow 2Ag_{(s)} + \frac{1}{2}O_{2(g)}$ which one of the following is true?
- $\Delta H = \Delta U$
 - $\Delta H < \Delta U$
 - $\Delta H > \Delta U$
 - $\Delta H = \frac{1}{2}\Delta U$
26. For a sample of perfect gas when its pressure is changed isothermally from p_i to p_f , the entropy change is given by
- $\Delta S = nR \ln \left(\frac{p_f}{p_i} \right)$
 - $\Delta S = nR \ln \left(\frac{p_i}{p_f} \right)$


$$(c) \Delta S = nRT \ln \left(\frac{p_f}{p_i} \right)$$

$$(d) \Delta S = RT \ln \left(\frac{p_i}{p_f} \right) \quad (\text{NEET Phase-II 2016})$$

27. The specific heat of a gas at constant volume is 0.075 cal/g-K. Predict the atomicity of the gas. (Molar mass of gas is 40 g mol⁻¹.)
 (a) 1 (b) 2
 (c) 3 (d) None of these
28. When the temperature is increased, surface tension of water
 (a) increases
 (b) decreases
 (c) remains constant
 (d) shows irregular behaviour.
29. If an endothermic reaction occurs spontaneously at constant temperature T and pressure P , then which of the following is true?
 (a) $\Delta G > 0$ (b) $\Delta H < 0$
 (c) $\Delta S > 0$ (d) $\Delta S < 0$
30. 22 g solid CO₂ or dry ice is enclosed in a bottle of one litre properly closed. If temperature of bottle is raised to 25 °C to sublime all the CO₂, the pressure in bottle is
 (a) 13.23 atm (b) 12.23 atm
 (c) 11.23 atm (d) 14.23 atm

SOLUTIONS

1. (b) : The reaction is



In this reaction, O₂ is the only gaseous reactant and CO₂ is the only gaseous product.

$$\therefore \Delta n_g = n_p - n_r = 6 - 7\frac{1}{2} = -\frac{3}{2}$$

$$\text{Given : } \Delta U(\text{or } q_v) = -3263.9 \text{ kJ mol}^{-1}$$

$$T = 25^\circ\text{C} = 298 \text{ K}$$

$$R = 8.314 \text{ J K}^{-1} \text{ mol}^{-1} = \frac{8.314}{1000} \text{ kJ K}^{-1} \text{ mol}^{-1}$$

$$\therefore \Delta H(\text{or } q_p) = \Delta U + \Delta n_g RT$$

$$= -3263.9 - \frac{3}{2} \times \frac{8.314}{1000} \times 298$$

$$= -3263.9 - 3.7 = -3267.6 \text{ kJ mol}^{-1}$$

2. (c) : Let the number of moles of each gas = x

$$\text{Fraction of hydrogen escaped} = \frac{1}{2}x$$

$$\frac{r_{O_2}}{r_{H_2}} = \sqrt{\frac{M_{H_2}}{M_{O_2}}} \Rightarrow \frac{n_{O_2}/t}{\frac{x}{2}/t} = \sqrt{\frac{2}{32}} = \sqrt{\frac{1}{16}} = \frac{1}{4}$$

$$\Rightarrow \frac{n_{O_2}/t}{\frac{x}{2}/t} = \frac{1}{4} \Rightarrow n_{O_2} = \frac{1}{8}x$$

$$\text{Hence, fraction of oxygen escaped} = \frac{1}{8}$$

3. (a) : Mass of aluminium, $m = 60.0 \text{ g}$
 Rise in temperature, $\Delta t = (328 \text{ K} - 308 \text{ K}) = 20 \text{ K}$
 Molar heat capacity, $C_m = 24 \text{ J K}^{-1} \text{ mol}^{-1}$

$$Q = \frac{C_m \times m \times \Delta t}{M}$$

$$\text{Molar mass of Al} = 27 \text{ g mol}^{-1}$$

$$\text{Heat required} = \frac{60 \times 24 \times 20}{27} \text{ J} = 1066.7 \text{ J} \approx 1.07 \text{ kJ}$$

4. (b) : Mole fraction of C (x_C)

$$= \frac{\text{Moles of C}}{\text{Total moles in the mixture}}$$

$$= \frac{5}{2+3+5+10} = \frac{5}{20} = \frac{1}{4}$$

$$\therefore p_C = P_{\text{total}} \times x_C$$

where, p_C = Partial pressure of C

$$\therefore P_{\text{total}} = 1.5 \times 4 = 6 \text{ atm}$$

5. (a) : $\Delta H = -21.976 \text{ kcal}$

$$\text{Here, } T_2 = 273 + 50 = 323 \text{ K}$$

$$T_1 = 273 + 27 = 300 \text{ K}$$

$$\Delta T = (T_2 - T_1) = (323 - 300) \text{ K} = 23 \text{ K}$$

$$\Delta C_p = \text{Sum of heat capacities of products}$$

$$- \text{Sum of heat capacities of reactants}$$

$$= (2 \times 8.86) - [6.8 + (3 \times 6.77)]$$

$$= 17.72 - (6.8 + 20.31) = -9.39 \text{ cal degree}^{-1}$$

$$= -9.39 \times 10^{-3} \text{ kcal degree}^{-1}$$

Substituting these values in Kirchoff's equation, we get

$$\Delta H_2 = \Delta H_1 + (T_2 - T_1)\Delta C_p$$

$$= -21.976 + [23 \times (-9.39 \times 10^{-3})]$$

$$= -21.976 + (-0.216)$$

$$= -22.192 \text{ kcal}$$

6. (a) : Applying ideal gas equation : $pV = nRT$

For H₂ gas,

$$0.8 \times 0.5 = n_{H_2} \cdot RT \Rightarrow n_{H_2} = \frac{0.8 \times 0.5}{RT} = \frac{0.4}{RT}$$

For O₂ gas,


$$0.7 \times 2.0 = n_{O_2} \cdot RT \Rightarrow n_{O_2} = \frac{0.7 \times 2.0}{RT} = \frac{1.4}{RT}$$

When gas mixture is introduced in 1 L vessel, then

$$p.V = (n_{\text{H}_2} + n_{\text{O}_2}) RT$$

$$p \times 1 = \left(\frac{0.4}{RT} + \frac{1.4}{RT} \right) RT$$

$$\therefore p = 0.4 + 1.4 = 1.8 \text{ bar}$$



$$\frac{\Delta H_{T_2} - \Delta H_{T_1}}{T_2 - T_1} = \Delta C_p$$

$$\begin{aligned} \therefore \Delta C_p &= \Sigma C_p (\text{products}) - \Sigma C_p (\text{reactants}) \\ &= (2 \times 25.1 + 3 \times 75.3) - (103.8 + 3 \times 28.8) \\ &= 276.1 - 190.2 = 85.9 \text{ J/K} = 85.9 \times 10^{-3} \text{ kJ/K} \end{aligned}$$

$$\therefore \frac{\Delta H_{(358 \text{ K})} - (-33.29)}{358 - 298} = 85.9 \times 10^{-3}$$

$$\Delta H_{358 \text{ K}} = -28.14 \text{ kJ/mol}$$

8. (d) : Average K.E. for 1 mol of gas = $\frac{3}{2} RT$

$$\text{Average K.E. for 1 molecule} = \frac{3 RT}{2 N_A}$$

9. (a) : $\Delta S_{\text{reaction}} = \Sigma S_{\text{products}} - \Sigma S_{\text{reactants}}$
 $= 2 \times S_{\text{H}_2\text{O}} - [2 \times S_{\text{H}_2} + S_{\text{O}_2}]$
 $= 2 \times 68 - [2 \times 126.6 + 201.20]$
 $= -318.4 \text{ J K}^{-1} \text{ mol}^{-1}$


10. (a) : For a spontaneous reaction,
 $\Delta G < 0$ i.e., $\Delta H - T\Delta S < 0$

$$T > \frac{\Delta H}{\Delta S}$$


$$T > \left(\frac{35.5 \times 1000}{83.6} = 424.6 \approx 425 \text{ K} \right)$$

$$\therefore T > 425 \text{ K}$$


11. (a) : The required equation is




Given :



Multiplying eq. (iii) by 2 and subtract eq. (i) from it, we get



Now, adding eq. (ii) and eq. (iv), we get



12. (b) : For N_2 at N.T.P.,

$$P = 760 \text{ mmHg} = 101325 \text{ N m}^{-2}$$

$$V = 22.4 \text{ L} = 0.0224 \text{ m}^3$$

$$M = 28 \text{ g/mol or } 0.028 \text{ kg/mol}$$

Putting these values in the equation,

$$u = \sqrt{\frac{3PV}{M}}$$

$$u = \sqrt{\frac{3 \times 101325 \times 0.0224}{0.028}} = 493.13 \text{ m s}^{-1}$$

13. (d) : For an irreversible process (work is being done by the system), $(dG)_{T,P} = -ve$ and $(dS)_{V,U} = +ve$.

14. (d) : $\therefore \frac{r_1}{r_2} = \frac{v/t_1}{v/t_2} = \sqrt{\frac{M_2}{M_1}}$

$$\frac{500 \text{ mL}}{0.5 \text{ h}} = \sqrt{\frac{M_{\text{SO}_2}}{M_{\text{He}}}} = \sqrt{\frac{64}{4}} = 4$$

$$\frac{1000 \text{ mL}}{t_2} = 4$$

Therefore, $t_2 = 4 \text{ h}$

15. (c) : $q = C \times \Delta t = 30 \times 4 = 120 \text{ kJ}$

Enthalpy of combustion of 4 g of graphite = -120 kJ

Enthalpy of combustion of 1 mole of graphite

$$= -\frac{12}{4} \times 120 = -360 \text{ kJ mol}^{-1}$$

16. (c) : $c_{\text{rms}} = \sqrt{\frac{3RT}{M}}$

$$\frac{c_{\text{rms}}(\text{H}_2)}{c_{\text{rms}}(\text{N}_2)} = \sqrt{\frac{T(\text{H}_2)}{M(\text{H}_2)} \times \frac{M(\text{N}_2)}{T(\text{N}_2)}}$$

$$\sqrt{7} = \sqrt{\frac{T(\text{H}_2)}{T(\text{N}_2)} \times \frac{28}{2}} \Rightarrow \frac{T(\text{H}_2)}{T(\text{N}_2)} = \frac{1}{2}$$

$$2T(\text{H}_2) = T(\text{N}_2)$$

$$\therefore T(\text{H}_2) < T(\text{N}_2)$$

17. (b) : Real gases show ideal behaviour at high temperature and low pressure.

18. (d) : Volume of 0.0168 mol of O_2 at STP

$$= 0.0168 \times 22400 \text{ mL} = 376.3 \text{ mL}$$

$$V_1 = 376.3 \text{ mL}, P_1 = 760 \text{ mmHg}, T_1 = 273 \text{ K}$$

$$V_2 = 428 \text{ mL}, P_2 = ?, T_2 = 298 \text{ K}$$

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2} \Rightarrow \frac{760 \times 376.3}{273} = \frac{P_2 \times 428}{298}$$

$$\Rightarrow P_2 = 729.4 \text{ mmHg}$$

$$\therefore \text{Pressure of water vapour} = 754 - 729.4$$

$$= 24.6 \text{ mmHg}$$

19. (c)

20. (c) : $\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2} \Rightarrow P_2 = \frac{P_1 V_1 T_2}{T_1 V_2}$... (i)

Substituting values in eq. (i),

$$p_2 = (0.75 \text{ N m}^{-2}) \frac{V_1}{(3V_1)} \frac{(2T_1)}{T_1}$$

$$p_2 = 0.5 \text{ N m}^{-2}$$

21. (a) : The entropy change at the melting point of a substance is given by

$$\Delta S_{fus} = \frac{\Delta H_{fus}}{T_{melting}}$$

$$\begin{aligned} \therefore T_{melting} &= \frac{\Delta H_{fus}}{\Delta S_{fus}} \\ &= \frac{7.25 \text{ kJ mol}^{-1}}{0.007 \text{ kJ K}^{-1} \text{ mol}^{-1}} = 1035.7 \text{ K} \end{aligned}$$

Hence, the melting point of KCl is 1035.7 K

$$\begin{aligned} 22. (a) : \bar{c}^* : \bar{c} : c &= \sqrt{\frac{2RT}{M}} : \sqrt{\frac{8RT}{\pi M}} : \sqrt{\frac{3RT}{M}} \\ &= \sqrt{2} : \sqrt{\frac{8}{\pi}} : \sqrt{3} = 1 : 1.128 : 1.224 \end{aligned}$$

23. (d) 24. (b)

25. (c) : $\Delta H = \Delta U + \Delta n_g RT$

$$\therefore \Delta n_g = \frac{1}{2} - 0 = \frac{1}{2}$$

$$\therefore \Delta H = \Delta U + \frac{1}{2} RT$$

$$\therefore \Delta H > \Delta U$$

26. (b) : For an ideal gas undergoing reversible expansion, when temperature changes from T_i to T_f and pressure changes from p_i to p_f ,

$$\Delta S = nC_p \ln \frac{T_f}{T_i} + nR \ln \frac{p_i}{p_f}$$

For an isothermal process, $T_i = T_f$ so, $\ln 1 = 0$

$$\therefore \Delta S = nR \ln \frac{p_i}{p_f}$$

27. (a) : $C_v = 0.075 \times 40 = 3 \text{ cal K}^{-1} \text{ mol}^{-1}$

$$C_p - C_v = R$$

$$\Rightarrow C_p - 3 = 2 \Rightarrow C_p = 5 \text{ cal K}^{-1} \text{ mol}^{-1}$$

$$\gamma = \frac{C_p}{C_v} = \frac{5}{3} = 1.66$$

Thus, the gas is monoatomic.

28. (b) : The decrease in surface tension with increase in temperature is due to the fact that with the

increase in temperature, the kinetic energy of the molecules increases and hence, the intermolecular attraction between the molecule decreases.

29. (c) : $\Delta S = \Delta S_{\text{system}} + \Delta S_{\text{surroundings}}$
 $\Delta S > 0$, for spontaneous process.

30. (a) : $w = 22 \text{ g}$; $V = 1 \text{ litre}$, $T = 298 \text{ K}$

$$PV = \frac{w}{M} RT \quad (\text{Molar mass of } \text{CO}_2 = 44 \text{ g mol}^{-1})$$

$$P_{\text{CO}_2} \times 1 = \frac{22}{44} \times 0.0821 \times 298$$

$$\therefore P_{\text{CO}_2} = 12.23 \text{ atm}$$


$$\begin{aligned} \therefore P_{\text{in bottle}} &= P_{\text{CO}_2} + P_{\text{atm}} \\ &= 12.23 + 1 = 13.23 \text{ atm} \end{aligned}$$

Your favourite MTG Books/Magazines available in
UTTAR PRADESH at

- Anwar Book Depot - Aligarh Mob: 9368487670, 9319326795
- Vimal Book House - Aligarh Ph: 0571-2704991; Mob: 9837192947, 8979371485
- New Vimal Book - Aligarh Ph: 2704992; Mob: 9456404096, 9319397725
- Friends Book Depot - Allahabad Ph: 0532-2461221; Mob: 9451743891
- Allahabad Book Centre - Allahabad
Ph: 0532-2623468, 2427636; Mob: 9415279039
- Natraj News Agency - Allahabad
Ph: 0532-2461291; Mob: 9335151613, 8400900917
- Vidya Kendra - Ballia Mob: 9415281234
- Competition Book Centre - Bareilly Mob: 9917233119, 8958555537
- Book World - Ghaziabad Ph: 0120-4113783; Mob: 9810933635, 9810636816
- Discount Wholesale Book Store - Gorakhpur
Ph: 0551-2347847; Mob: 9336403332, 9307021482
- Atul Pustak Bhandar - Gorakhpur Mob: 9935210622
- Modern Book Depot - Jhansi Mob: 7619845201
- Raj Book Distributors - Kanpur
Mob: 9235616506, 9235616515, 9415031904, 9336336918
- Sakshi Book Seller - Kanpur Ph: 0512-3938874; Mob: 9935939510, 9369505152
- Aashirwad Book Depot - Lucknow
Ph: 0522-4004346; Mob: 9235501197, 9335527368
- Balaji Book Centre - Lucknow Ph: 0522-2202082; Mob: 9415418393
- Book Sadan - Lucknow Ph: 0522-4002509; Mob: 9839487327
- Books N Books - Lucknow Ph: 00522-4005415; Mob: 9415026661, 7897055107
- Rama Book Depot - Lucknow Ph: 0522-4080133; Mob: 7388948811
- Ravi Stationery And Books Shope - Lucknow Mob: 8127206914, 7619840391
- Universal Booksellers - Lucknow
Ph: 0522-2625894, 2624135, 3919708; Mob: 9838204366
- Student Book Store - Mathura Ph: 0565-6450961; Mob: 9359518693
- Asha Book Agency - Meerut Ph: 0121-2640540, 4006978; Mob: 9927009672
- Ideal Book Depot - Meerut Ph: 0121-2660648, 2668279; Mob: 9837894819
- Mahi Book Palace - Meerut Ph: 0121-2641791; Mob: 9927012549
- Delta Stationers - Noida Ph: 0120-2550643; Mob: 9818189817, 9999048591
- Suvidha Stationers - Noida Mob: 9810987964
- Gaurav Book Agency - Varanasi Mob: 8173997477
- Shri Krishna Book Agency & Stationers - Varanasi Mob: 9415820103, 9369396321

Visit "MTG IN YOUR CITY" on www.mtg.in to locate nearest book seller OR write to info@mtg.in OR call **0124-6601200** for further assistance.

EXAMINER'S MIND CLASS XI



The questions given in this column have been prepared strictly on the basis of NCERT Chemistry for Class XI. This year JEE (Main & Advanced)/NEET/AIIMS have drawn their papers heavily from NCERT books.

Section - I	Q. 1 to 10 Only One Option Correct Type MCQs.
Section - II	Q. 11 to 13 More than One Options Correct Type MCQs.
Section - III	Q. 14 to 17 Paragraph Type MCQs having Only One Option Correct.
Section - IV	Q. 18 & 19 Matching List Type MCQs having Only One Option Correct.
Section - V	Q. 20 to 22 Assertion Reason Type MCQs having Only One Option Correct. Mark the correct choice as : (a) If both assertion and reason are true and reason is the correct explanation of assertion. (b) If both assertion and reason are true but reason is not the correct explanation of assertion. (c) If assertion is true but reason is false. (d) If both assertion and reason are false.
Section - VI	Q. 23 to 25 Integer Value Correct Type Questions having Single Digit Integer Answer, ranging from 0 to 9 (both inclusive).

CHEMICAL BONDING AND MOLECULAR STRUCTURE

SECTION - I

Only One Option Correct Type

- Which of the following has highest bond angle?
(a) NO_2^+ (b) NO_2 (c) NO_2^- (d) NO_3^-
- In PO_4^{3-} ion, the formal charge on the oxygen atom of P—O bond is
(a) +1 (b) -1 (c) -0.75 (d) +0.75
- The ionic bond $A^+ B^-$ is formed when
(1) I.E. of A is low (2) E.A. of B is high
(3) lattice energy of AB is high
(4) lattice energy of AB is low.
(a) 1 and 2 (b) 1, 2 and 3
(c) 3 only (d) 4 only
- Which of the following compounds/ions has V-shape?
(a) SF_5^- (b) SF_4 (c) OSF_4 (d) SF_2
- In which one of the following species the central atom has the type of hybridisation which is not the same as in the other three?
(a) SbCl_5^{2-} (b) PCl_5 (c) SF_4 (d) I_3^-
- The correct increasing order of *s*-character (in percentage) in the hybrid orbitals in the given molecules/ions is (assume all hybrid orbitals are exactly equivalent) :

CO_3^{2-}	XeF_4	I_3^-	NCl_3	BeCl_2
I	II	III	IV	V

 (a) $\text{II} < \text{III} < \text{IV} < \text{I} < \text{V}$ (b) $\text{II} < \text{IV} < \text{III} < \text{V} < \text{I}$
 (c) $\text{III} < \text{II} < \text{I} < \text{V} < \text{IV}$ (d) $\text{II} < \text{IV} < \text{III} < \text{I} < \text{V}$
- In ICl_2^+ , ICl_2^- , and ICl_4^- , sum of the bond pairs and lone pairs on each iodine atom are respectively
(a) 2, 2 and 4 (b) 2, 3 and 2
(c) 4, 5 and 4 (d) 4, 5 and 6
- Which of the following molecules is formed by *p-p* overlapping?
(a) F_2 (b) H_2O (c) HCl (d) NH_3
- In XeF_2 molecule, the angle between two lone pair orbitals is α , the angle between lone pair orbital and bond pair orbital is β and the angle between two bond pair orbitals is γ then, which one is correct order of angle?
(a) $\alpha = \beta = \gamma$ (b) $\alpha > \beta > \gamma$
(c) $\gamma > \beta > \alpha$ (d) $\gamma > \alpha > \beta$
- Which one of the following properties is not shown by NO ?
(a) Its bond order is 2.5.
(b) It is diamagnetic in gaseous state.
(c) It is a neutral oxide.
(d) It combines with oxygen to form nitrogen dioxide.

SECTION - II

More than One Options Correct Type

11. According to VBT in XeF_2 , Xe uses five sp^3d hybridised orbital for molecule formation. Select correct statement for XeF_2 .
- Three sp^3d orbitals are used for covalent bonding with F.
 - Three sp^3d orbitals occupy lone pairs of Xe.
 - Two sp^3d orbitals are used for covalent bonding with F.
 - Two sp^3d orbitals are occupied by lone pairs of Xe.
12. Which statements are correct for AB_x type molecule?
- If the electronegativity of central atom decreases, the bond angle decreases.
 - If the size of central atom increases, the bond angle decreases.
 - If the electronegativity of atom B decreases, the bond angle increases.
 - If the electronegativity of atom B decreases, the bond angle decreases.
13. Which of the following statements are not correct?
- Hybrid orbitals form stronger bonds than p -orbitals.
 - Excitation of electron is essential for hybridisation.
 - Boiling point of H_2O is more than H_2S .
 - Resonance plays an important role in molecular orbital theory.

SECTION - III

Paragraph Type

Paragraph for Questions 14 and 15

Hydrogen bonding originates from the dipole-dipole interaction between H-atom and any of the other atom like F, O, N and in some cases with Cl atom also. There are two types of H-bonding *i.e.*; intermolecular and intramolecular H-bonding.

14. Which of the following molecules does not consist of intramolecular H-bonding?
- Chloral
 - Chloral hydrate
 - o*-Hydroxybenzaldehyde
 - o*-Chlorophenol
15. Which of the following statements is incorrect?
- Boiling point of H_2O_2 is greater than that of H_2O .
 - Ethylene glycol is less viscous than glycerol.

- o*-Nitrophenol can be separated from its *m*- and *p*-isomers using its steam volatile property.
- In ice, each 'O' atom is tetrahedrally arranged by four H-atoms which are all equidistant.

Paragraph for Questions 16 and 17

The VSEPR model considers double and triple bonds to have slightly greater repulsive effects than single bonds because of the repulsive effect of π -electrons. Consider molecule $(\text{CH}_3)_2\text{C}=\text{CH}_2$ for given questions :

16. Which type of overlapping is not observed in the given molecule?
- sp^3-s
 - sp^2-s
 - $sp-s$
 - sp^2-sp^2
17. Select which has the largest bond angle in the given molecule.
- $\text{CH}_3-\text{C}(\text{H})_2-\text{CH}_3$
 - $\text{H}_3\text{C}-\text{C}(\text{H})=\text{C}$
 - $\text{H}-\text{C}(\text{H})_2-\text{H}$
 - All are same.

SECTION - IV

Matching List Type

18. Match the List I with List II and select the correct answer using the codes given below the lists :

List I (Species)		List II (Properties)	
A.	O_2^{2-}	P.	Isoelectronic with N_2
B.	CO	Q.	Fractional bond order
C.	NO^+	R.	Paramagnetic
D.	He_2^+	S.	Diamagnetic

	A	B	C	D
(a)	P	R, S	Q, P	P, S
(b)	Q, P	S	P, S	R, S
(c)	S	P, S	P, S	Q, R
(d)	R	R, S	S	P, Q

19. Match the List I with List II and select the correct answer using the codes given below the lists :


List I		List II	
(P)	CH_4	1.	Linear molecule
(Q)	BeCl_2	2.	Tetrahedral molecule
(R)	H_2O	3.	Pyramidal molecule
(S)	NH_3	4.	V-shaped molecule

	P	Q	R	S
(a)	2	1	4	3
(b)	2	3	4	1
(c)	1	2	3	4
(d)	2	3	1	4

SECTION - V

Assertion Reason Type

20. **Assertion :** C_2H_2 molecule is linear.
Reason : In C_2H_2 , carbon atoms remain unhybridised.
21. **Assertion :** H_2 molecule is more stable than HeH molecule.
Reason : The antibonding electron in the molecule destabilises HeH.

22. **Assertion :** In ClF_3 molecule,
- 
- $P = Q \neq 90^\circ$

Reason : The molecule is slightly bent T-shaped and there is repulsion between lone pairs of electrons.

SECTION - VI

Integer Value Correct Type

23. How many of the following compounds violate octet rule?
 BrF_5 , SF_6 , IF_7 , $XeOF_4$, ClF_2^- , PCl_4^+
24. A diatomic molecule has a dipole moment of 1.2 D. If the bond distance is 1.0 Å, $1/x$ of an electronic charge 'e' exists on each atom. The value of x is
25. The number of dative bonds in sulphuric acid molecule is

HYDROGEN

SECTION - I


Only One Option Correct Type

1. Which one of the following processes will produce permanent hard water?
 (a) Addition of Na_2SO_4 to water
 (b) Saturation of water with $CaCO_3$
 (c) Saturation of water with $MgCO_3$
 (d) Saturation of water with $CaSO_4$
2. Reaction between following pairs will produce hydrogen except
 (a) $Cu + HCl$ (b) $Fe + H_2O_{(steam)}$
 (c) $Mg + H_2O (hot)$ (d) $Na + Alcohol$.
3. Protium, deuterium, tritium differ in
 (a) number of protons and physical properties
 (b) atomic number and chemical properties
 (c) number of neutrons and physical properties
 (d) number of neutrons and chemical properties.
4. Moist hydrogen peroxide cannot be dried over conc. H_2SO_4 because
 (a) it is oxidised by H_2SO_4
 (b) it is reduced by H_2SO_4
 (c) it can catch fire
 (d) all of these.
5. Which of the following equations depicts reducing nature of H_2O_2 ?
 (a) $2[Fe(CN)_6]^{4-} + 2H^+ + H_2O_2 \longrightarrow 2[Fe(CN)_6]^{3-} + 2H_2O$
 (b) $I_2 + H_2O_2 + 2OH^- \longrightarrow 2I^- + 2H_2O + O_2$
 (c) $Mn^{2+} + H_2O_2 \longrightarrow Mn^{4+} + 2OH^-$
 (d) $PbS + 4H_2O_2 \longrightarrow PbSO_4 + 4H_2O$
6. Heavy water is manufactured
 (a) by repeated electrolysis of water with 3% aqueous NaOH
 (b) by electrolysis of water containing heavy hydrogen dissolved in it
 (c) by combination of hydrogen and heavier isotope of oxygen
 (d) none of the above.
7. Which of the following explanations justifies for not placing hydrogen in either the group of alkali metals or halogens?
 (a) Hydrogen is much lighter than alkali metals or halogens.
 (b) Hydrogen atom does not contain any neutron.
 (c) The ionization energy of hydrogen is too high for group of alkali metals and too low for halogen group.
 (d) Hydrogen can form compounds with almost all other elements.
8. The freezing point of heavy water is
 (a) $-3.82^\circ C$ (b) $3.82^\circ C$
 (c) $0^\circ C$ (d) $-10^\circ C$
9. Water softening by Clark's process involves use of
 (a) calcium carbonate (b) sodium carbonate
 (c) potash alum (d) calcium hydroxide.
10. Which is false about H_2O_2 ?
 (a) It can act as both oxidising and reducing agent.
 (b) Two $-OH$ bonds lie in the same plane.
 (c) It is pale blue liquid.
 (d) It can be oxidised by O_3 .

FULLY LOADED & COMPLETELY UPDATED

mtg

MTG's BITSAT Explorer is not only the most exhaustive prep-tool, but also the only book available at present, updated as per the latest BITSAT syllabus for students aspiring for top rank in BITSAT 2018.



₹ 750



Free Interactive CD
with MTG's BITSAT Explorer.
Simulate the online testing
experience with this unique
interactive CD.

Runs on both PCs and Macs.

Get MTG's BITSAT Explorer today for a real-world feel of BITSAT. Find out what's different about the BITSAT test, including its pattern of examination and key success factors. Be it with chapter-wise MCQs or model test papers, check how good your chances are for glory in BITSAT 2018.

FEATURES:

- Covers all 5 subjects - Physics, Chemistry, Mathematics, English & Logical Reasoning
- Chapterwise 1,000+ MCQs in each section for practice
- 5 Model Test Papers with detailed solutions
- Free interactive CD

Visit www.MTG.in to buy online. Or visit a leading bookseller near you.
For more information, email info@mtg.in or call 1800 300 23355 (toll-free).

SECTION - II

More than One Options Correct Type

11. Which of the following statements are correct?
 (a) Metallic hydrides are deficient of hydrogen.
 (b) Metallic hydrides conduct heat and electricity.
 (c) Ionic hydrides do not conduct electricity in solid state.
 (d) Ionic hydrides are very good conductors of electricity in solid state.
12. Which of the following statements are correct?
 (a) Dissociation of H_2 molecules is an endothermic process.
 (b) Hydrogen at the moment of liberation is more active.
 (c) Atomic hydrogen is powerful reducing agent.
 (d) Ordinary hydrogen is an oxidising agent.
13. Which of the following statements are correct about 6.8% strength of H_2O_2 ?
 (a) Its normality is 4 N.
 (b) Its molarity is 2 M.
 (c) Its volume strength is 22.4 V.
 (d) Volume strength = $11.2 \times M$.

SECTION - III

Paragraph Type

Paragraph for Questions 14 and 15

Hydrogen peroxide is a powerful oxidising agent, both in the acidic and alkaline medium.


In acidic medium; $H_2O_2 + 2H^+ + 2e^- \longrightarrow 2H_2O$

In alkaline medium; $H_2O_2 + 2e^- \longrightarrow 2OH^-$

Hydrogen peroxide acts as a reducing agent towards powerful oxidising agents.

In acidic medium; $H_2O_2 \longrightarrow 2H^+ + O_2 + 2e^-$


In alkaline medium, however, its reducing nature is more effective.



14. On addition of H_2O_2 to acidified $KMnO_4$, $KMnO_4$ gets decolourised due to
 (a) oxidation of $KMnO_4$
 (b) reduction of $KMnO_4$
 (c) both oxidation and reduction
 (d) none of these.
15. In which of the following reactions, H_2O_2 acts as an oxidising agent?
 (a) $2I^- + H_2O_2 + 2H^+ \longrightarrow I_2 + 2H_2O$
 (b) $IO_4^- + H_2O_2 \longrightarrow IO_3^- + H_2O + O_2$
 (c) $Ag_2O + H_2O_2 \longrightarrow 2Ag + H_2O + O_2$
 (d) $2MnO_4^- + 6H^+ + 5H_2O_2 \longrightarrow 2Mn^{2+} + 8H_2O + 5O_2$

Paragraph for Questions 16 and 17

Formation of methane from syn gas is represented by the following equations :



16. X in reaction (ii) refers to
 (a) liquid water (b) steam
 (c) oxygen (d) carbon monoxide.
17. Hydrogen prepared by above method is passed over Ni catalyst
 (a) to remove traces of CO
 (b) to prepare H_2O (c) to prepare CH_4
 (d) to separate H_2 from water gas.

SECTION - IV

Matching List Type

18. Match the List I with List II and select the correct answer using the codes given below the lists :

List I	List II
(P) Synthesis gas	1. Forms stoichiometric compounds with s-block elements
(Q) Dihydrogen	2. Prolonged electrolysis of water
(R) Heavy water	3. Mixture of CO and H_2
(S) Hydrogen peroxide	4. Zn + NaOH
	5. Oxidising agent and/or Reducing agent

P	Q	R	S
(a) 1, 2	3	4, 5	1, 2
(b) 3	1, 4, 5	2	5
(c) 1, 2	3, 4	1, 5	4
(d) 1, 3	2, 3	4	5

19. Match the List I with List II and select the correct answer using the codes given below the lists :

List I	List II
(P) Ionic hydride	1. LiH
(Q) Electron deficient hydride	2. CaH_2
(R) Hydrolith	3. AlH_3
(S) Covalent hydride	4. SiH_4

	P	Q	R	S
(a)	1	2,3	3	1, 4
(b)	2,3	1	2	1, 4
(c)	1, 2	3	2	3, 4
(d)	3,4	3	2	1, 2

SECTION - V

Assertion Reason Type

20. **Assertion :** *Para*-hydrogen with lower energy is favoured at high temperature.
Reason : The thermal conductivity of *para*-hydrogen is lesser than that of *ortho*-hydrogen.
21. **Assertion :** Permanent hardness of water is removed by treatment with washing soda.
Reason : Washing soda reacts with soluble magnesium and calcium sulphates to form insoluble carbonates.

SECTION - VI

Integer Value Correct Type

23. Half litre each of three samples of H_2O_2 labelled 10 vol, 15 vol, 20 vol are mixed and then diluted with 1700 mL of water. Calculate relative strength of resultant H_2O_2 solution.
24. Calculate the degree of hardness of a sample of water containing 6 mg of $MgSO_4$ per kg of water.
25. What is the sum of protons, electrons and neutrons in the heaviest isotope of hydrogen?

SOLUTIONS

CHEMICAL BONDING AND MOLECULAR STRUCTURE


1. (a): NO_2^+ has linear geometry and therefore, bond angle is highest (180°).
2. (c): In PO_4^{3-} ion, formal charge on each O-atom of P—O bond

$$= \frac{\text{Total charge}}{\text{Number of O-atoms}} = -\frac{3}{4} = -0.75$$
3. (b)
4. (d): SF_2 has sp^3 hybridisation and 'V' shape due to the presence of two lone pairs on sulphur.
5. (a): $SbCl_5^{2-}$ has sp^3d^2 hybridisation while all other species involve sp^3d hybridisation.
6. (a):

Species	Hybridisation
CO_3^{2-}	sp^2
XeF_4	sp^3d^2
I_3^-	sp^3d
NCl_3	sp^3
$BeCl_2$	sp

Therefore, the correct increasing order of percentage of s-character is :


$$II < III < IV < I < V$$



So, total number of lone pairs and bond pairs in ICl_2^+ = 2 + 2 = 4


in ICl_2^- = 3 + 2 = 5 and ICl_4^- = 2 + 4 = 6

8. (a): F_2 is formed by *p-p* overlapping whereas, H_2O , HCl and NH_3 are formed by *s-p* overlapping.



10. (b): NO is paramagnetic in gaseous state due to the presence of one unpaired electron.





11. (b,c):






12. (a, b, c)

13. (b, d): Excitation of electron is not essential for hybridisation, vacant orbitals can also participate in hybridisation. Resonance has no role in M.O. theory.

14. (a)





15. (d)

There is no $sp-s$ overlapping.


Diamagnetic

$$B.O. = \frac{N_b - N_a}{2} = \frac{10 - 8}{2} = 1$$



Diamagnetic

$$B.O. = \frac{10 - 4}{2} = 3$$



Paramagnetic

$$B.O. = \frac{2 - 1}{2} = \frac{1}{2}$$

19. (a): (P) $\rightarrow 2: CH_4$ is sp^3 hybridised so, it is tetrahedral molecule.




(Q) $\rightarrow 1: BeCl_2$ is sp hybridised, hence linear in shape.

(R) $\rightarrow 4: H_2O$ is sp^3 hybridised, but due to the presence of the lone pairs, it is V-shaped.

(S) $\rightarrow 3: NH_3$ is sp^3 hybridised, but due to the presence of one lone pair of electrons, its shape is pyramidal.

20. (c): In C_2H_2 , C atom is sp -hybridised. Thus, linear in shape.

21. (a): Bond order of $H_2 = 1$ (2 electrons; $\sigma 1s^2$). Bond order of $HeH = 0.5$ (3 electrons, $\sigma 1s^2, \sigma^* 1s^1$). Greater the bond order of molecule, more stable the molecule. Hence, H_2 is more stable than HeH .

23. (5): (i) $BrF_5 = 12$ electrons(ii) $SF_6 = 12$ electrons(iii) $IF_7 = 14$ electrons(v) ClF_2^- ; $[F-\ddot{Cl}-F]^- = 10$ electronsHence, except PCl_4^+ all five molecules violate octet rule.

$$24. (4): \text{Partial charge} = \frac{\text{Dipole moment}}{\text{Bond distance}} = \frac{1.2 \times 10^{-18} \text{ esu.cm}}{1.0 \times 10^{-8} \text{ cm}} = 1.2 \times 10^{-10} \text{ esu}$$

The fraction of an electronic charge is

$$\frac{1.2 \times 10^{-10}}{4.8 \times 10^{-10}} = \frac{1}{4} \quad \therefore \text{Value of } x = 4$$

25. (2)





HYDROGEN

1. (d): Permanent hardness of water is due to the presence of soluble chlorides and sulphates of calcium and magnesium.


2. (a): (a) $Cu + HCl \rightarrow$ No reaction
(b) $3Fe + 4H_2O \rightarrow Fe_3O_4 + 4H_2$
 Steam Ferroso
 ferric oxide

mtg

Master Resource Books in Chemistry



These books comprehensively cover all topics, concepts and techniques to prepare students for all the competitive exams. i.e. JEE (Main & Advanced), PET, NEET, AIIMS and JIPMER.



HIGHLIGHTS:


- Concise theory for competitive exams
- Illustrations with detailed solutions
- Twists and turns to learn important formulae
- Elaborate solutions to mysterious NCERT problems
- Practice assignments with pinch of hints
- Solved MCQs single and multiple option correct type, Assertion & Reason, Fill in the blanks, True or False, Comprehension, Integer & Matching types with Miscellaneous Questions

Success in **CET 2018 Now Made Easy**



Available at all leading book shops throughout India.
For more information or for help in placing your order:
Call 0124-6601200 or email: info@mtg.in

Visit
www.mtg.in
for latest offers
and to buy
online!




3. (c)
 4. (a): H_2SO_4 acts as an oxidising agent and decomposes H_2O_2 .
 $\text{H}_2\text{O}_2 + [\text{O}] \longrightarrow \text{H}_2\text{O} + \text{O}_2$
 5. (b): When H_2O_2 behaves as a reducing agent, it evolves molecular oxygen during the reaction. Therefore, only in option (b), it behaves as a reducing agent.
 6. (a): Heavy water is manufactured by repeated electrolysis of water (containing a little NaOH).
 7. (c): Hydrogen has ionization energy value which is too high as compared to alkali metals and too low as compared to halogens and thus, cannot be placed in any of these two groups.

8. (b) 9. (d)

10. (b): The two O–H bonds are in different planes due to repulsion between different bonding and antibonding orbitals.

11. (a, b, c)

12. (a, b, c): Compounds which undergo reduction on heating with H_2 are reduced by atomic H at ordinary temperature.



The H so produced is very reactive and its life is only one-third of a second.

13. (a, b, c, d):

$$\text{Volume strength} = 5.6 \times \frac{\text{Percentage strength}}{\text{eq. wt. of } \text{H}_2\text{O}_2} \times 10$$


$$= 5.6 \times \frac{6.8}{17} \times 10 = 22.4 \text{ V}$$

(a) Normality = $\frac{\text{Volume strength}}{5.6} = 4 \text{ N}$


(b) Molarity = $\frac{\text{Volume strength}}{11.2} = 2 \text{ M}$

(c) Volume strength = 22.4 V

(d) Volume strength = $11.2 \times \text{molarity} = 22.4 \text{ V}$



On the other hand, we can write it as,



15. (a): $2\text{I}^- \longrightarrow \text{I}_2$, hence H_2O_2 acts as an oxidising agent.

16. (b): Water gas is mixed with steam, producing CO_2 .

17. (a): The hydrogen manufactured by the given method is utilised for synthesis of NH_3 in which CO acts as a poison for catalyst. Thus, Ni is used to remove traces of CO.

18. (b) 19. (c)

20. (d): *Para*-hydrogen with lower energy is favoured at low temperature.

The thermal conductivity of *para*-hydrogen is 50% greater than that of *ortho*-hydrogen.

21. (a) 22. (a)

23. (7): Volume strength of $\text{H}_2\text{O}_2 = 5.6 \times N$

$$10 \text{ vol. } \text{H}_2\text{O}_2 = \frac{10}{5.6} N \text{ H}_2\text{O}_2$$

$$15 \text{ vol. } \text{H}_2\text{O}_2 = \frac{15}{5.6} N \text{ H}_2\text{O}_2$$

$$20 \text{ vol. } \text{H}_2\text{O}_2 = \frac{20}{5.6} N \text{ H}_2\text{O}_2$$

As 500 mL of each is mixed then total volume of mixture = 1500 mL

Also this is diluted with 1700 mL, so total volume becomes 3200 mL

$$N \times 3200 = \frac{10 \times 500}{5.6} + \frac{15 \times 500}{5.6} + \frac{20 \times 500}{5.6}$$

$$\text{or } N_{\text{H}_2\text{O}_2} = \frac{500 \times 45}{5.6 \times 3200} = 1.255 \text{ N}$$

$$\therefore \text{Volume strength of } \text{H}_2\text{O}_2 = 1.255 \times 5.6 = 7.03 \approx 7$$

24. (5): 1 g-mole or 120 g $\text{MgSO}_4 \equiv 1$ g-mole or 100 g CaCO_3

$$6 \times 10^{-3} \text{ g } \text{MgSO}_4 = \frac{100 \times 6 \times 10^{-3}}{120}$$

$$= 5 \times 10^{-3} \text{ g of } \text{CaCO}_3$$

Thus, 1000 g of water contains MgSO_4 equivalent to

$$5 \times 10^{-3} \text{ g of } \text{CaCO}_3 = \frac{5 \times 10^{-3}}{1000} \times 10^6$$

$$10^6 \text{ g of water will contain} = 5 \text{ g of } \text{CaCO}_3$$

\therefore Hardness of given water sample = 5 ppm

25. (4): Tritium T or ${}^3_1\text{H}$, the heaviest isotope of hydrogen has 1 proton, 2 neutrons and 1 electron. Hence, sum of protons, neutrons and electrons is 4.



ACE

YOUR WAY CBSE XI



Series 3

CHAPTERWISE PRACTICE PAPER : STATES OF MATTER | THERMODYNAMICS

Time Allowed : 3 hours

Maximum Marks : 70

GENERAL INSTRUCTIONS

- (i) All questions are compulsory.
- (ii) Q. no. 1 to 5 are very short answer questions and carry 1 mark each.
- (iii) Q. no. 6 to 10 are short answer questions and carry 2 marks each.
- (iv) Q. no. 11 to 22 are also short answer questions and carry 3 marks each.
- (v) Q. no. 23 is a value based question and carries 4 marks.
- (vi) Q. no. 24 to 26 are long answer questions and carry 5 marks each.
- (vii) Use log tables if necessary, use of calculators is not allowed.


1. Under what conditions q and w become state functions?
 2. What would have happened to the gas if the molecular collisions were not elastic?
 3. The C_p and C_v of a gas are 20.834 and $12.520 \text{ J K}^{-1} \text{ mol}^{-1}$ respectively. What is the atomicity of the gas?
 4. How is the mole fraction of a gas component related to its partial pressure and the total pressure?
 5. What is residual entropy?
 6. Acetic acid (ethanoic acid) and hydrochloric acid react with KOH solution. The enthalpy of neutralisation of ethanoic acid is $-55.8 \text{ kJ mol}^{-1}$ while that of hydrochloric acid is $-57.1 \text{ kJ mol}^{-1}$. Why are these two values different?
 7. (a) Why does sharp glass edge become smooth on heating it upto its melting point in a flame? Explain which property of liquids is responsible for this phenomenon.
(b) Which two other properties of liquids can be explained on the basis of the above property?
 8. Distinguish between the total kinetic energy of a molecule and its translational kinetic energy. For what type of gas molecules these two are same?
 9. (a) Define Charles' law.
(b) What is the coefficient of volume expansion of a gas?
 10. What do you understand by the term 'absolute zero temperature'? What is its significance?
- OR**
- How does the magnitude of the enthalpy change depend on the strength of the intermolecular interaction for the substances undergoing phase changes? Explain with example.
11. Which of the following processes are accompanied by an increase of entropy?
 - (a) Dissolution of iodine in a solvent.
 - (b) HCl is added to AgNO_3 solution and precipitate of AgCl is obtained.
 - (c) A partition is removed to allow the gases to mix.

12. 20% of N_2O_4 molecules are dissociated in a sample of gas at 27°C and 760 torr. Calculate the density of the equilibrium mixture.
13. Calculate the standard Gibbs energy change for the formation of propane at 298 K.
 $3\text{C}(\text{graphite}) + 4\text{H}_2(\text{g}) \rightarrow \text{C}_3\text{H}_8(\text{g})$
 $\Delta_f H^\circ$ for propane, $\text{C}_3\text{H}_8(\text{g}) = -103.8 \text{ kJ mol}^{-1}$
 (Given : $S_m^\circ [\text{C}_3\text{H}_8(\text{g})] = 270.2 \text{ J K}^{-1} \text{ mol}^{-1}$
 $S_m^\circ (\text{graphite}) = 5.70 \text{ J K}^{-1} \text{ mol}^{-1}$
 and $S_m^\circ [\text{H}_2(\text{g})] = 130.7 \text{ J K}^{-1} \text{ mol}^{-1}$)
14. Although heat is a path function but heat absorbed by the system under certain specific conditions is independent of path. What are those conditions? Explain.
15. 1 Litre flask containing $\text{NH}_3(\text{g})$ at 2.0 atm at 200 K is connected by a narrow tube of negligible volume to another 800 mL flask containing $\text{HCl}(\text{g})$ at 8.0 atm at 200 K. The two gases react according to equation :
 $\text{NH}_3(\text{g}) + \text{HCl}(\text{g}) \rightarrow \text{NH}_4\text{Cl}(\text{s}) ; \Delta H = -43 \text{ kJ/mol}$
 If heat capacity of $\text{HCl}(\text{g})$ is $20 \text{ J K}^{-1} \text{ mol}^{-1}$, determine the heat produced, final temperature and final pressure inside the flask. (The heat capacity of flask and volume of solid NH_4Cl in flask is negligible.)
16. (a) (i) How many calories are required to heat 100 g of copper ($s = 0.092 \text{ cal/g/K}$) from 10 to 100°C ?
 (ii) The same quantity of heat as in (i) is added to 100 g of aluminium ($s = 0.217 \text{ cal/g/K}$) at 10°C . Which gets hotter, the copper or aluminium?
 (b) How much heat is required to change 10 g ice at 0°C to steam at 100°C ? Latent heat of fusion and vaporization for H_2O are 80 cal/g and 540 cal/g respectively. Specific heat of water is 1 cal/g.
17. A student forgot to add the reaction mixture to the round bottom flask at 27°C but put it on the flame. After a lapse of time, he realised his mistake, using a pyrometer he found the temperature of the flask was 477°C . What fraction of air would have been expelled out?

OR

The drain cleaner, Drainex contains small bits of aluminium which react with caustic soda to produce dihydrogen. What volume of dihydrogen at 20°C and one bar will be released when 0.15 g of aluminium reacts?

18. (a) 1.0 mol of a monoatomic ideal gas is expanded from state (1) to state (2) as shown in figure. Calculate the work done for the expansion of gas from state (1) to state (2) at 298 K.
 (b) A certain volume of dry air at NTP is expanded reversibly to four times its volume isothermally. Calculate the final pressure.



19. A sample of 0.16 g CH_4 was subjected to combustion at 27°C in a bomb calorimeter. The temperature of the calorimeter system (including water) was found to rise by 0.5°C . Calculate the heat of combustion of methane at (i) constant volume and (ii) constant pressure. The thermal capacity of calorimeter system is 17.7 kJ K^{-1} and R is $8.314 \text{ J K}^{-1} \text{ mol}^{-1}$.
20. Give reasons for the following :
 (a) The size of balloon becomes larger and larger as it ascends into higher altitudes.
 (b) Tyres of automobiles are inflated to lesser pressure in summer than in winter.
21. Calculate the rise in temperature when a gas, for which $\gamma = 1.5$, is compressed to 27 times its original pressure, assuming the initial temperature to be 27°C .
22. (a) Calculate the maximum efficiency of an engine operating between 100°C and 25°C .
 (b) Same mass of diamond and graphite (both being carbon) are burnt in oxygen. Will the heat produced be same or different? Why?
23. Sejal asked Saloni that why people are undergoing medical check up before going on Amarnath Yatra. Saloni explained the reason to Sejal and Sejal got satisfied, meanwhile Rishabh told that liquid boils at lower temperature at a hill station than in a plane area.
 (a) What reason was explained by Saloni to satisfy Sejal?
 (b) Why does liquid boil at lower temperature at hills stations?
 (c) What values are associated with Saloni?
 (d) What is critical temperature?
24. Nitrogen molecule (N_2) has radius of about 0.2 nm. Assuming that nitrogen molecule is spherical in shape, calculate

- (a) volume of a single molecule of N_2 .
 (b) the percentage of empty space in one mole of N_2 gas at S.T.P.


OR

- (a) When an ideal gas expands in vacuum, there is neither absorption nor evolution of heat. Why?
 (b) Determine the value of ΔH and ΔU for the reversible isothermal evaporation of 90.0 g of water at 100°C . Assume that water behaves as an ideal gas and heat of evaporation of water is 540 cal g^{-1} .
25. The density of the vapours of a substance at 1 atm pressure and 500 K is 0.36 kg m^{-3} . The vapours effuse through a small hole at a rate of 1.33 times faster than oxygen under the same conditions :
- (a) Determine : (i) molecular weight, (ii) molar volume, (iii) compressibility factor (Z) of the vapours, (iv) Which forces among gas molecules are dominating, the attractive or the repulsive?
 (b) If the vapours behave ideally at 1000 K, determine the average translational kinetic energy of a molecule.

OR

Two moles of a perfect gas undergo the following process :

- (a) A reversible isobaric expansion from (1 atm, 20 L) to (1 atm, 40 L).
 (b) A reversible isochoric change of state from (1 atm, 40 L) to (0.5 atm, 40 L).
 (c) A reversible isothermal compression from (0.5 atm, 40 L) to (1 atm, 20 L).
 (i) Sketch with labels each of the process on the same P - V diagram.
 (ii) Calculate the total work (w) and the total heat change (q) involved in the above process.
 (iii) What will be the values of ΔU , ΔH and ΔS for the overall process?
26. (a) The figure given below represents P - V diagrams of different stages of a thermodynamic process. Calculate the work done in each stage and also the net work done in the complete cyclic process.



- (b) Give a statement which includes the main ideas of the first law and second law of thermodynamics.

OR

- (a) The composition of the equilibrium mixture ($\text{Cl}_2 \rightleftharpoons 2\text{Cl}$), which is attained at 1200°C , is determined by measuring the rate of effusion through a pin hole. It is observed that at 1.80 mm Hg pressure, the mixture effuses 1.16 times as fast as krypton effuses under the same conditions. Calculate the fraction of chlorine molecules dissociated into atoms (atomic weight of Kr is 84).
 (b) The compression factor (compressibility factor) for 1 mol of a van der Waals' gas at 0°C and 100 atm pressure is found to be 0.5. Assuming that the volume of a gas molecule is negligible, calculate the van der Waals' constant a .

SOLUTIONS

- In a thermodynamically reversible process.
- On every collision, there would have been loss of energy. As a result, the molecules would have slowed down and ultimately settle down. The gas pressure would have gradually become zero.
- $\gamma = \frac{C_p}{C_v} = \frac{20.834}{12.520} = 1.66$
Therefore, the gas is monoatomic.
- Mole fraction, is equal to the partial pressure divided by the total pressure.
$$x_i = \frac{p_i}{P_{\text{total}}}$$
- The entropy possessed by a substance at absolute zero is called residual entropy.
- Ethanoic acid is a weak acid while hydrochloric acid is strong acid (completely ionized). The enthalpy of neutralisation of weak acid (ethanoic acid) and strong base (KOH) is numerically less than $-57.1 \text{ kJ mol}^{-1}$ (heat of neutralisation of strong acid and strong base) because the ionization of ethanoic acid is not complete in solution and some energy is utilised for dissociating acid molecules.
- (a) On heating the glass, it melts and takes up rounded shape at the edges which has minimum surface area. This is due to the property of surface tension of liquids.

(b) The following two properties of liquids can be explained on the basis of surface tension :

- Capillary action of water.
- Spherical shape of small droplets.

8. The total kinetic energy of a molecule includes the translational kinetic energy, rotational kinetic energy and vibrational kinetic energy.

For monoatomic gases, both rotational and vibrational kinetic energies are zero. Therefore, for these gases, total kinetic energy is equal to translational kinetic energy.

9. (a) Charles' law states that "The volume of a given amount of a gas at constant pressure varies directly with its absolute temperature.

$V \propto T$ (pressure is constant)

(b) According to Charles' law,

$$V_t = V_0 \left(1 + \frac{t}{273} \right) = V_0 (1 + \alpha t)$$

$\alpha = \frac{1}{273}$ = Coefficient of volume expansion

here, V_0 = Volume at 0°C and V_t = Volume at $t^\circ\text{C}$

10. According to Charles' law, if a gas is cooled to -273°C , its volume becomes zero as

$$\text{Volume at } -273^\circ\text{C} = V_0 \left(1 - \frac{273}{273} \right) = 0$$


where, V_0 = Volume at 0°C

This means that -273°C should be the lowest temperature because any further cooling would lead to a volume less than zero or negative volume which is meaningless. Therefore, this temperature is termed as absolute zero temperature. But volume never approaches zero at -273°C because all gases condense to liquid or solid before the attainment of this temperature.

OR


The magnitude of the enthalpy change depends directly on the strength of the intermolecular interaction.

For example, intermolecular hydrogen bonding between water molecules lead to large attractive intermolecular energy holding water molecules tightly in liquid phase. For an organic liquid such as acetone, the intermolecular dipole-dipole interactions are significantly weaker. Thus, it requires less heat to vaporise 1 mole of acetone than it does to vaporise one mole of water.




11. (a) Entropy is increased because solid iodine is converted into liquid phase (solution).

Order of entropy : Gases > liquids > solids



In this reaction, liquid HCl is mixed with the solution of AgNO_3 and the formation of $\text{AgCl}(\text{ppt.})$ occurs, hence entropy is decreased.

(c) Entropy is increased because after removing the partition both the gases will mix and move upto larger space, thus the randomness of the molecules will increase.



At $t = 0$ 1 mol 0

At equilibrium (1 - 0.2) mol 0.4

Total moles = 0.8 + 0.4 = 1.2

$PV = nRT$


$1 \times V = 1.2 \times 0.0821 \times 300$ ($\because P = 760 \text{ torr} = 1 \text{ atm}$)

$V = 29.556 \text{ L}$

$$d_{\text{N}_2\text{O}_4} = \frac{m}{V} = \frac{0.8 \times 92}{29.556} = 2.490$$

$$d_{\text{NO}_2} = \frac{m}{V} = \frac{0.4 \times 46}{29.556} = 0.6225$$

$$d_{\text{mix}} = 2.490 + 0.6225 = 3.1125 \text{ g/L}$$



$$\Delta_r H^\circ = \Delta_f H^\circ (\text{C}_3\text{H}_8) - \{3\Delta_f H^\circ (\text{C}) + 4\Delta_f H^\circ (\text{H}_2)\}$$

$$= -103.8 - 0 - 0 = -103.8 \text{ kJ mol}^{-1}$$

$$\Delta_r S^\circ = S_m^\circ (\text{C}_3\text{H}_8) - \{3S_m^\circ (\text{C}) + 4S_m^\circ (\text{H}_2)\}$$

$$= 270.2 - \{3 \times 5.70 + 4 \times 130.7\}$$

$$= 270.2 - 539.9 = -269.7 \text{ J K}^{-1} \text{ mol}^{-1}$$

$$\Delta_r G^\circ = \Delta_r H^\circ - T\Delta_r S^\circ$$

$$= -103.8 - 298 \times (-269.7 \times 10^{-3}) = -23.43 \text{ kJ mol}^{-1}$$

14. The two conditions under which heat becomes independent of path are

(i) when volume remains constant

(ii) when pressure remains constant.

(i) At constant volume : By first law of thermodynamics,

$$\Delta U = q + w \quad \text{or} \quad q = \Delta U - w$$

$$w = -P\Delta V \quad \therefore \quad q = \Delta U + P\Delta V$$

But as volume remains constant, $\Delta V = 0$

$$\therefore q_v = \Delta U$$


As ΔU is a state function. Hence, q_v is a state function.

(ii) At constant pressure : $q_p = \Delta U + P\Delta V$

$$\text{But } \Delta U + P\Delta V = \Delta H$$

$$\therefore q_p = \Delta H$$

As ΔH is a state function, therefore, q_p is a state function.



Thus, heat produced during reaction,

$$Q = n_{\text{NH}_4\text{Cl}} \times \Delta H = 0.125 \times 43 \times 10^3 = 5375 \text{ J}$$

\therefore During the formation of 1 mole $\text{NH}_4\text{Cl}(s)$, heat produced is ΔH . Because this heat is used to increase the temperature of HCl in both the flasks.

Thus, $Q = n \times C_v \times \Delta T$

$$5375 = 0.275 \times 20 \times \Delta T \Rightarrow \Delta T = 977.27 \text{ K}$$

$$\therefore T_f = 200 + 977.27 = 1177.27 \text{ K}$$

$$\text{and } P_f = \frac{nRT_f}{V} = \frac{0.275 \times 0.0821 \times 1177.27}{1.8}$$

$$= 14.76 \text{ atm} \quad (\text{Here, } V = 1 + 0.8 = 1.8 \text{ L})$$

16. (a) (i) $\Delta H = ms \Delta T$

$$= (0.092 \text{ cal/g/K}) (100 \text{ g}) [(373 - 283) \text{ K}] = 828 \text{ cal}$$

(ii) Since the specific heat capacity of copper is less than that of aluminium, less heat is required to raise the temperature of a mass of copper by 1 K than is required for an equal mass of aluminium. Hence, the copper gets hotter.

(b) Total heat absorbed

$$= \Delta H_{\text{fusion}} + \Delta H_{\text{temp. rise}} + \Delta H_{\text{vap.}}$$

$$= 10 \times 80 + 10 \times 1 \times 100 + 10 \times 540 = 7200 \text{ cal}$$

17. Suppose the number of moles of gas present at 27°C in flask of volume V at pressure P is n_1 , then assuming ideal gas behaviour,

$$PV = n_1 R \times 300 \quad \dots(i)$$

Suppose, $n_2 =$ number of moles at 477°C , then

$$PV = n_2 R \times 750 \quad \dots(ii)$$


From eq. (i) and eq. (ii), we get

$$n_2 = \frac{300}{750} \times n_1 = 0.4 n_1$$

\therefore 0.6 moles have been expelled out.

OR

The reaction between aluminium and caustic soda is



\therefore 54 g of Al produces H_2 at S.T.P. = $3 \times 22.4 \text{ L}$

0.15 g of Al will produce H_2 at S.T.P.

$$= \frac{3 \times 22.4}{54} \times 0.15 = 0.186 \text{ L}$$

Applying ideal gas equation,

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2} \Rightarrow \frac{1 \times 0.186}{273} = \frac{0.987 \times V_2}{293}$$

$$V_2 = \frac{293}{0.987} \times \frac{1 \times 0.186}{273} = 0.2030 \text{ L} = 203 \text{ mL}$$

18. (a) It is clear from the figure that the process has been carried out in infinite steps and therefore, it is an isothermal reversible expansion process.

$$w = -2.303 nRT \log \frac{V_2}{V_1} = -2.303 nRT \log \frac{P_1}{P_2}$$

$$= -2.303 \times 1 \text{ mol} \times 8.314 \text{ J K}^{-1} \text{ mol}^{-1} \times 298 \text{ K} \times \log 2 = -1717.6 \text{ J}$$

(b) At constant temperature,

$$P_1 V_1 = P_2 V_2$$

$$P_2 = \frac{P_1 V_1}{V_2} = \frac{1 \times V_1}{4V_1} = 0.25 \text{ atm}$$


19. (i) Heat of combustion at constant volume,

$\Delta U =$ Heat capacity of calorimeter system \times

$$\text{rise in temperature} \times \frac{\text{Mol. mass of compound}}{\text{Mass of compound}}$$

$$= 17.7 \times 0.5 \times \frac{6}{0.6} = 8$$

$$\Delta U = -885 \text{ kJ mol}^{-1}$$



$$\Delta n_g = 1 - 3 = -2, T = 300 \text{ K,}$$

$$R = 8.314 \times 10^{-3} \text{ kJ K}^{-1} \text{ mol}^{-1}$$

Heat of combustion at constant pressure,

$$\Delta H = \Delta U + \Delta n_g RT$$

$$= -885 + (-2) \times 8.314 \times 10^{-3} \times 300$$

$$= -889.988 \text{ kJ mol}^{-1}$$

20. (a) At higher altitudes, the atmospheric pressure decreases. Thus, the pressure outside the balloon decreases. To regain equilibrium with the external pressure, the gas inside, expands to decrease its pressure. Hence, the size of the balloon increases.

(b) In summer, due to higher temperature, the average kinetic energy of the air molecules inside the tyres increases *i.e.*, molecules start moving faster. Hence, the pressure on the walls of the tube increases. If pressure inside is not kept low at the time of inflation, at higher temperature, the pressure may become so high that the tyre may burst.

21. From adiabatic gas equation, $\left(\frac{T_1}{T_2}\right)^\gamma = \left(\frac{P_1}{P_2}\right)^{\gamma-1}$

Given that, $\gamma = 1.5$, $P_2/P_1 = 27$, $T_1 = 300$ K
Substituting the values,

$$\left(\frac{300}{T_2}\right)^{1.5} = \left(\frac{1}{27}\right)^{1.5-1}$$

After taking log on both sides,

$$1.5(\log 300 - \log T_2) = 0.5(\log 1 - \log 27)$$

$$\log 300 - \log T_2 = -\frac{0.5 \log 27}{1.5}$$

$$\log T_2 = \log 300 + 0.33 \log 27$$

$$= 2.477 + 0.4723 = 2.9493$$

$$\therefore T_2 = \text{antilog } 2.9493 = 889.81$$

$$= (889.81 - 273)^\circ\text{C} = 616.81^\circ\text{C}$$

Hence, the rise in temperature
 $= 616.81 - 27 = 589.81^\circ\text{C}$

22. (a) Efficiency = $\frac{T_2 - T_1}{T_2}$

$$T_2 = 100 + 273 = 373 \text{ K}$$

$$T_1 = 25 + 273 = 298 \text{ K}$$

$$\text{Efficiency} = \frac{373 - 298}{373} = \frac{75}{373} = 0.20 = 20 \%$$

(b) Heat evolved will be different. This is because they have different crystal structures.

23. (a) At higher altitude, atmospheric pressure is low, so generally people have difficulty in breathing therefore, people must need to undergo for medical check-up before going on Amarnath Yatra.

(b) At higher altitude, atmospheric pressure is low. Therefore, the liquid boils at lower temperature.

(c) Knowledge, health concern.

(d) When density of liquid and vapours become the same, the clear boundary between liquid and vapours disappears. This temperature is called critical temperature.

24. (a) The volume of a sphere = $\frac{4}{3}\pi r^3$

where, r is the radius of the sphere.

For N_2 molecule,

$$r = 0.2 \text{ nm} = 0.2 \times 10^{-9} \text{ m} = 2 \times 10^{-8} \text{ cm}$$

$$\text{Volume of a molecule of } \text{N}_2 = \frac{4}{3} \times \frac{2}{7} \times (2 \times 10^{-8})^3 \text{ cm}^3$$

$$= 3.35 \times 10^{-23} \text{ cm}^3$$

(b) To calculate the empty space, let us first find the total volume of 1 mole (6.02×10^{23} molecules) of N_2 .

$$\text{Volume of } 6.02 \times 10^{23} \text{ molecules of } \text{N}_2$$

$$= 3.35 \times 10^{-23} \times 6.02 \times 10^{23} = 20.17 \text{ cm}^3$$

Now, volume occupied by 1 mole of gas at S.T.P.
 $= 22.4 \text{ litre} = 22400 \text{ cm}^3$

$$\text{Empty volume} = \text{Total volume of gas} -$$

$$\text{Volume occupied by molecules}$$

$$= (22400 - 20.17) \text{ cm}^3$$

$$= 22379.83 \text{ cm}^3$$

$$\therefore \text{Percentage empty space} = \frac{\text{Empty space}}{\text{Total volume}} \times 100$$

$$= \frac{22379.83}{22400} \times 100 = 99.9 \%$$

Thus, 99.9% of space of 1 mole of N_2 at S.T.P is empty.

OR

(a) In an ideal gas, there are no intermolecular forces of attraction and therefore, no force opposes the expansion in vacuum. Hence, no energy is required to overcome these forces. Moreover, when a gas expands against vacuum, work done is zero because $p_{\text{ext}} = 0$. Therefore, internal energy of the system does not change, i.e., there is no absorption or evolution of heat.

(b) Total heat change,

$$\Delta H = 90.0 \times 540 = 48600 \text{ cal}$$

$$\text{Now, } \Delta H = \Delta U + P\Delta V$$

$$\text{Here, } \Delta V = (V_{\text{vapour}} - V_{\text{liquid}}) = V_{\text{vapour}}$$

(Vol. of liquid is negligible as compared to volume of vapour)

$$\Delta H = \Delta U + PV_{\text{vapour}} = \Delta U + nRT$$

$$n = \frac{90}{18} = 5 \text{ mol, } R = 2 \text{ cal K}^{-1} \text{ mol}^{-1}, T = 373 \text{ K}$$

$$\therefore \Delta H = \Delta U + 5 \times 2 \times 373$$

$$\Delta H = \Delta U + 3730; \quad \Delta U = \Delta H - 3730$$

$$= 48600 - 3730 = 44870 \text{ cal}$$

25. (a) (i) $\frac{r_{\text{vapour}}}{r_{\text{O}_2}} = \sqrt{\frac{M_{\text{O}_2}}{M_{\text{vapour}}}}$; $1.33 = \sqrt{\frac{32}{M_{\text{vapour}}}}$

$$M_{\text{vapour}} = 18.1$$

(ii) Molar volume = $\frac{\text{Molar mass}}{\text{Density}}$

$$= \frac{18.1}{0.36 \times 10^3} = 50.27 \times 10^{-3} \text{ m}^3 / \text{mol}$$

(iii) Compressibility factor,

$$Z = \frac{PV}{RT} = \frac{101325 \times 50.27 \times 10^{-3}}{8.314 \times 500} = 1.225$$


(iv) $Z > 1$, shows that repulsive forces are dominant.

(b) Translational K.E. per molecule = $\frac{3}{2} \times \frac{R}{N_A} \times T$

$$= \frac{3}{2} \times \frac{8.314}{6.023 \times 10^{23}} \times 1000 = 2.07 \times 10^{-20} \text{ J}$$

OR

(i) $AB \rightarrow$ Isobaric process
 $BC \rightarrow$ Isochoric process
 $CA \rightarrow$ Isothermal compression



(ii) Total work = $w_{AB} + w_{BC} + w_{CA}$

$$w = -P \times (V_2 - V_1) + 0 + 2.303 nRT \log(V_2/V_1)$$

$$= -1 \times (40 - 20) + 0 + 2.303 P_1 V_1 \times \log(V_2/V_1)$$

$[nRT = P_1 V_1]$

$$= -20 + 2.303 \times (20) \log \frac{40}{20}$$

$$= -20 + (2.303)(20)(0.301029)$$

$$= -20 + 13.86539574 = -6.13 \text{ L atm}$$

$$= \frac{6.13 \times 8.314 \text{ J}}{0.0821} = -620.76 \text{ J}$$

$$w = q = -620.76 \text{ J}$$

(iii) In cyclic process :

$$\Delta U = 0, \Delta H = 0 \text{ and } \Delta S = 0$$

26. (a) Process $A \rightarrow B$ (expansion), $P = 12 \times 10^5 \text{ N m}^{-2}$,
 $\Delta V = 8 - 2 = 6 \text{ L} = 6 \times 10^{-3} \text{ m}^3$
 \therefore Work done = $-P\Delta V = -(12 \times 10^5) \times (6 \times 10^{-3}) \text{ J}$
 $= -7200 \text{ J}$

Process $B \rightarrow C$, No change in volume, i.e., $\Delta V = 0$

$$\therefore \text{Work done} = 0$$

Process $C \rightarrow D$ (contraction), $\Delta V = 8 - 2 = 6 \text{ L}$
 $= 6 \times 10^{-3} \text{ m}^3$, $P = 4 \times 10^5 \text{ N m}^{-2}$

$$\therefore \text{Work done} = P\Delta V = (4 \times 10^5) (6 \times 10^{-3}) = 2400 \text{ J}$$

Process $D \rightarrow A$, No change in volume, i.e., $\Delta V = 0$

$$\therefore \text{Work done} = 0$$

$$\therefore \text{Net work done in the complete cyclic process}$$

$$= -7200 + 2400 \text{ J} = -4800 \text{ J}$$

-ve sign shows that net work has been done by the gas.


(b) The energy of the universe is constant, whereas the entropy of the universe is continuously increasing and tends to maximum.

OR

(a) According to Graham's law of diffusion

$$\frac{r_{\text{mix}}}{r_{\text{Kr}}} = \sqrt{\frac{M_{\text{Kr}}}{M_{\text{mix}}}} \Rightarrow 1.16 = \sqrt{\frac{84}{M_{\text{mix}}}}$$

$$\therefore M_{\text{mix}} = 62.425$$



Initial mole 1 0

Final mole 1 - x 2x

$$\text{Total moles} = 1 - x + 2x = 1 + x$$

$$\therefore M_{\text{mix}} = \frac{2x(35.5) + (1-x) \times 71}{1+x} = 62.425$$

$$\frac{71}{1+x} = 62.425$$

$$x = 0.1373 = 13.7\% \text{ dissociated}$$

(b) For 1 mol of a gas, the van der Waals' equation is

$$\left(p + \frac{a}{V_m^2} \right) (V_m - b) = RT$$

Ignoring b , we get (given volume of gas molecule is negligible)

$$\left(p + \frac{a}{V_m^2} \right) V_m = RT \Rightarrow pV_m + \frac{a}{V_m} = RT$$

$$\frac{pV_m}{RT} + \frac{a}{V_m RT} = 1 \Rightarrow Z = \frac{pV_m}{RT} = 1 - \frac{a}{V_m RT} \quad \dots(i)$$

$$Z = \frac{pV_m}{RT} = 0.5 \Rightarrow V_m = \frac{0.5RT}{p}$$

Now, from eq. (i), $0.5 = 1 - \frac{a}{(0.5RT/p)RT}$

$$a = (0.5) \left(\frac{0.5RT}{p} \right) RT = 0.25 \frac{R^2 T^2}{p}$$

Substituting the given values, we get

$$a = (0.25) \left[\frac{(0.082 \text{ L atm K}^{-1} \text{ mol}^{-1})^2 (273 \text{ K})^2}{100 \text{ atm}} \right]$$

$$= 1.2528 \text{ L}^2 \text{ atm mol}^{-2}$$



MPP-5 MONTHLY Practice Problems

Class XI

This specially designed column enables students to self analyse their extent of understanding of specified chapter. Give yourself four marks for correct answer and deduct one mark for wrong answer. Self check table given at the end will help you to check your readiness.

Equilibrium

Total Marks : 120

Time Taken : 60 Min.


NEET / AIIMS

Only One Option Correct Type

- Vapour density of PCl_5 at 200°C is 70. Determine the degree of dissociation of PCl_5 at this temperature.
(a) 48.9% (b) 57.9% (c) 38.8% (d) 46.1%
- Starting with 1 mol of $\text{O}_2(\text{g})$ and 2 mol of $\text{SO}_2(\text{g})$ the equilibrium for the formation of $\text{SO}_3(\text{g})$ was established at a certain temperature. If V is the volume of the vessel and $2x$ is the number of moles of $\text{SO}_3(\text{g})$ present, the equilibrium constant will be
(a) $\frac{x^2V}{(1-x)^3}$ (b) $\frac{4x^2V}{(2-x)(1-x)}$
(c) $\frac{(1-x)^3}{2V}$ (d) $\frac{x^2V}{(2-x)(1-x)V}$
- Which one of the following solutions will have pH close to unity?
(a) 100 mL of M/10 HCl + 100 mL of M/10 NaOH
(b) 55 mL of M/10 HCl + 45 mL of M/10 NaOH
(c) 10 mL of M/10 HCl + 90 mL of M/10 NaOH
(d) 75 mL of M/5 HCl + 25 mL of M/5 NaOH
- When different types of salts have nearly same solubility product constant (K_{sp}) which is lesser than K_{sp} value of one salt, then the most soluble salt is the one
(a) which produces maximum number of ions
(b) which produces minimum number of ions
(c) which produces more charge on ion
(d) none of these.
- Consider the following reactions in which all the reactants and the products are in gaseous state :
 $2\text{PQ} \rightleftharpoons \text{P}_2 + \text{Q}_2 ; K_1 = 2.5 \times 10^5$
 $\text{PQ} + 1/2\text{R}_2 \rightleftharpoons \text{PQR} ; K_2 = 5 \times 10^{-3}$
The value of K_3 for the equilibrium :
 $1/2\text{P}_2 + 1/2\text{Q}_2 + 1/2\text{R}_2 \rightleftharpoons \text{PQR}$, is
(a) 2.5×10^{-3} (b) 2.5×10^3
(c) 1.0×10^{-5} (d) 5.0×10^3
- The following equilibrium is established when hydrogen chloride is dissolved in acetic acid,
 $\text{HCl} + \text{CH}_3\text{COOH} \rightleftharpoons \text{Cl}^- + \text{CH}_3\text{COOH}_2^+$
The set that characterises the conjugate acid-base pairs is
(a) (HCl, CH_3COOH) and ($\text{CH}_3\text{COOH}_2^+$, Cl^-)
(b) (HCl, $\text{CH}_3\text{COOH}_2^+$) and (CH_3COOH , Cl^-)
(c) ($\text{CH}_3\text{COOH}_2^+$, HCl) and (Cl^- , CH_3COOH)
(d) (HCl, Cl^-) and ($\text{CH}_3\text{COOH}_2^+$, CH_3COOH)
- One mole of $\text{N}_2\text{O}_4(\text{g})$ at 300 K is kept in a closed container under one atmosphere. It is heated to 600 K when 20% by mass of $\text{N}_2\text{O}_4(\text{g})$ decomposes to $\text{NO}_2(\text{g})$. The resultant pressure is
(a) 1.2 atm (b) 2.4 atm (c) 2.0 atm (d) 1.0 atm
- A 0.02 M solution of pyridinium hydrochloride has pH = 3.44. Calculate the ionisation constant of pyridine.
(a) 1.8×10^{-9} (b) 2.1×10^{-8}
(c) 1.5×10^{-9} (d) 1.0×10^{-9}
- For a reaction $2\text{A} + \text{B} \rightleftharpoons \text{C} + \text{D}$, the value of K_p will be :
(a) $K_p = \frac{n_C n_D}{n_A^2 n_B} \frac{V_P}{RT^2}$ (b) $K_p = \frac{n_C n_D}{n_A^2 n_B} \frac{P}{RT}$
(c) $K_p = \frac{n_C n_D}{n_A^2 n_B} \frac{RT^2}{V_P}$ (d) $K_p = \frac{n_C n_D}{n_A^2 n_B} \frac{V_P}{RT}$
- The equilibrium constant for the following reaction,
 $\text{N}_2\text{O}_4(\text{g}) \rightleftharpoons 2\text{NO}_2(\text{g})$ is $K = \frac{[\text{NO}_2]^2}{[\text{N}_2\text{O}_4]}$

mtg

Last-minute check on your NEET readiness



₹ 550



MTG's NEET Explorer helps students self-assess their readiness for success in NEET. Attempting the tests put together by MTG's experienced team of editors and experts strictly on the NEET pattern and matching difficulty levels, students can easily measure their preparedness for success. **Order now!**



Scan now with your smartphone or tablet*

HIGHLIGHTS:

- 10 Model Test Papers based on latest NEET syllabus
- Last 15 years' solved test papers of AIPMT / NEET
- Includes NEET 2017 solved paper
- Detailed solutions for self-assessment and to practice time management




Available at all leading book shops throughout India. For more information or for help in placing your order: Call 0124-6601200 or email: info@mtg.in

*Application to read QR codes required

Visit www.mtg.in for latest offers and to buy online!


Which of the following graphs is correct?



11. For the following isomerisation reaction :
Cyclohexane \rightleftharpoons 1-Hexene, $K = 1.732$

Which of the following statements holds good at point 'A'?

- (a) $Q > K$
(b) $Q < K$
(c) $Q = K = 1$
(d) $Q = K = 1.732$



12. The decreasing base strength of OH^- , NH_2^- , $\text{HC}\equiv\text{C}^-$ and CH_3CH_2^- is
- (a) $\text{CH}_3\text{CH}_2^- > \text{NH}_2^- > \text{HC}\equiv\text{C}^- > \text{OH}^-$
(b) $\text{HC}\equiv\text{C}^- > \text{CH}_3\text{CH}_2^- > \text{NH}_2^- > \text{OH}^-$
(c) $\text{OH}^- > \text{NH}_2^- > \text{HC}\equiv\text{C}^- > \text{CH}_3\text{CH}_2^-$
(d) $\text{NH}_2^- > \text{HC}\equiv\text{C}^- > \text{OH}^- > \text{CH}_3\text{CH}_2^-$

Assertion & Reason Type


Directions : In the following questions, a statement of assertion is followed by a statement of reason. Mark the correct choice as :

- (a) If both assertion and reason are true and reason is the correct explanation of assertion.
(b) If both assertion and reason are true but reason is not the correct explanation of assertion.
(c) If assertion is true but reason is false.
(d) If both assertion and reason are false.

13. **Assertion :** K_p can be equal to or less than or even greater than the value of K_c .

Reason : $K_p = K_c (RT)^{\Delta n}$

14. **Assertion :** On cooling a freezing mixture, colour of the mixture turns to pink from deep blue for a reaction,



Reason : Reaction is endothermic so on cooling, the reaction moves to backward direction.

15. **Assertion :** Common salt is added during manufacturing of soap.

Reason : Common salt helps in the formation of soap.

JEE MAIN / JEE ADVANCED

Only One Option Correct Type

16. The equilibrium constants K_{p1} and K_{p2} for the reactions : $X \rightleftharpoons 2Y$ and $Z \rightleftharpoons P + Q$, respectively are in the ratio of 1 : 9. If the degree of dissociation of X and Z be equal then the ratio of total pressures at these equilibria is
(a) 1 : 9 (b) 1 : 36 (c) 1 : 1 (d) 1 : 3
17. Which of the following is not correct about the percentage ionisation of BOH ?

- (a) $\frac{K_w[\text{H}^+]}{K_b + K_w}$ (b) $100 \times \sqrt{\frac{K_b}{C}}$
(c) $\frac{100}{1 + 10^{(pK_b - pOH)}}$ (d) $\frac{K_b \times 100}{K_b + \text{OH}^-}$

18. The pH of pure water at 25°C and 35°C are 7 and 6 respectively. The heat of formation of water from H^+ and OH^- is

- (a) 84.551 kcal mol⁻¹ (b) -84.551 kcal mol⁻¹
(c) 74.551 kcal mol⁻¹ (d) -74.551 kcal mol⁻¹

19. Ammonia under a pressure of 15 atm at 27°C is heated to 347°C in a closed vessel in the presence of a catalyst. Under the conditions, NH_3 is partially decomposed according to the equation, $2\text{NH}_3 \rightleftharpoons \text{N}_2 + 3\text{H}_2$. The vessel is such that the volume remains effectively constant whereas pressure increases to 50 atm. Calculate the percentage of NH_3 actually decomposed.
(a) 65% (b) 61.3% (c) 62.5% (d) 64%

More than One Options Correct Type

20. When two reactants, A and B are mixed to give two products, C and D , the reaction quotient, (Q) at the initial stages of the reaction
(a) is zero (b) decreases with time
(c) is independent of time
(d) increases with time.

21. Which of the following are heterogeneous systems?

- (a) $\text{Ice} \rightleftharpoons \text{Water}$ (b) $\text{Water} \rightleftharpoons \text{Liquid}$
(c) $\text{S}_{\text{Rhombic}} \rightleftharpoons \text{S}_{\text{Monoclinic}}$
(d) $\text{C}_{\text{Diamond}} \rightleftharpoons \text{C}_{\text{Amorphous}}$


22. A reaction $\text{S}_{8(g)} \rightleftharpoons 4\text{S}_{2(g)}$ is carried out by taking 2 moles of $\text{S}_{8(g)}$ and 0.2 mole of $\text{S}_{2(g)}$ in a reaction vessel of 1 litre at 627°C. Which of the following are correct if $K_c = 6.30 \times 10^{-6}$?

- (a) Reaction quotient is 8×10^{-4}
(b) Reaction proceeds in backward direction
(c) $K_p = 2.55 \text{ atm}^3$
(d) Reaction proceeds in forward direction

23. The solubility of a sparingly soluble salt A_xB_y in water at $25^\circ\text{C} = 1.4 \times 10^{-4}$ M. The solubility product is 1.1×10^{-11} . The possibilities are
 (a) $x = 1, y = 2$ (b) $x = 2, y = 1$
 (c) $x = 1, y = 3$ (d) $x = 3, y = 1$

Integer Answer Type

24. In the study of reaction,




A and B were mixed in a reaction vessel at 300 K. The initial concentration of B was 1.5 times the initial concentration of A. After the equilibrium has been attained, the equilibrium concentrations of A and D were equal. The value of equilibrium constant at 300 K is

25. In 1 L saturated solution of AgCl [$K_{sp}(\text{AgCl}) = 1.6 \times 10^{-10}$], 0.1 mol of CuCl [$K_{sp}(\text{CuCl}) = 1 \times 10^{-6}$] is added. The resultant concentration of Ag^+ in the solution is 1.6×10^{-x} . The value of x is
26. A mixture of N_2 and H_2 in the molar ratio 1 : 3 attains equilibrium when 50% of mixture has reacted. If P is the total pressure of the mixture, the partial pressure of NH_3 formed is P/y . The value of y is

Comprehension Type

The concentration of hydrogen ion is a measure of acidity or alkalinity of a solution.

For monobasic acids :



where, K_a = Dissociation constant of acid

C = Molarity of acid

$$[\text{H}^+] = C\alpha, [\text{H}^+] = \sqrt{K_a C}$$

pH of a weak acid can be calculated using $[\text{H}^+]$ by any of above methods.

$$[\text{H}^+]_{\text{Total}} = \sqrt{C_1 K_{a_1} + C_2 K_{a_2}} \quad (\text{when two weak acids are mixed})$$

$[\text{H}^+]$ of polyprotic acid (Weak, H_3A) (has dissociation constants $K_{a_1} \gg K_{a_2} \gg K_{a_3}$) ; will be contributed by first dissociation at the most.

27. What will be the sulphide ion concentration of a dilute solution that has been saturated with 0.1 M

H_2S if the pH of the solution is 3?

(Given : $K_{a_1} = 1 \times 10^{-7}$; $K_{a_2} = 1.3 \times 10^{-13}$)

- (a) 1.25×10^{-11} (b) 1.3×10^{-13}
 (c) 1.45×10^{-9} (d) 1.3×10^{-15}

28. The pH of 0.01 M HCOOH ($K_a = 1.4 \times 10^{-4}$) will be
 (a) 2.928 (b) 3.296 (c) 4.962 (d) 5.926

Matrix Match Type

29. Match the Column I with Column II and choose the correct option using the codes given below.

Column I (Salt)		Column II (Solubility product)	
(A) Zirconium phosphate		(p) $27 S^4$	
(B) Aluminium phosphate		(q) $108 S^5$	
(C) Calcium phosphate		(r) $6912 S^7$	
(D) Sodium phosphate		(s) S^2	
A	B	C	D
(a) s	r	q	p
(b) r	s	q	p
(c) p	r	s	q
(d) q	r	s	p

30. Match the Column I with Column II and choose the correct option using the codes given below.

Column I (Salt)		Column II (Degree of hydrolysis)	
(A) NaCl		(p) $h = \sqrt{\frac{K_w}{C \cdot K_b}}$	
(B) CH_3COONa		(q) $h = \sqrt{\frac{K_w}{K_a \times K_b}}$	
(C) NH_4Cl		(r) No hydrolysis	
(D) $\text{CH}_3\text{COONH}_4$		(s) $h = \sqrt{\frac{K_w}{C \cdot K_a}}$	
A	B	C	D
(a) p	q	s	r
(b) r	s	p	q
(c) q	r	s	p
(d) p	r	s	q



Keys are published in this issue. Search now! ☺

SELF CHECK



Check your score! If your score is

No. of questions attempted	> 90%	EXCELLENT WORK !	You are well prepared to take the challenge of final exam.
No. of questions correct	90-75%	GOOD WORK !	You can score good in the final exam.
Marks scored in percentage	74-60%	SATISFACTORY !	You need to score more next time.
		< 60%	NOT SATISFACTORY !	Revise thoroughly and strengthen your concepts.

NEET | JEE

ESSENTIALS

Class
XII

Maximise your chance of success, and high rank in NEET, JEE (Main and Advanced) by reading this column. This specially designed column is updated year after year by a panel of highly qualified teaching experts well-tuned to the requirements of these Entrance Tests.


Unit 3

General Principles and Processes of Isolation of Elements | *p*-Block Elements (Group 15 to 18)

GENERAL PRINCIPLES AND PROCESSES OF ISOLATION OF ELEMENTS

GENERAL INTRODUCTION

The percentage of different elements in the earth crust are :
O - 49%, Si - 26%, Al - 7.5%, Fe - 4.2%, Ca - 3.2%,
Na - 2.4%, K - 2.3%, Mg - 2.5%, H - 1%




ORES / MINERALS

Combined state	Element	Ore/mineral
Oxides	Fe	Haematite (Fe_2O_3) Magnetite (Fe_3O_4) Limonite ($\text{Fe}_2\text{O}_3 \cdot 3\text{H}_2\text{O}$) Chromite ($\text{FeO} \cdot \text{Cr}_2\text{O}_3$)
	Al	Bauxite ($\text{Al}_2\text{O}_3 \cdot 2\text{H}_2\text{O}$) Diaspore ($\text{Al}_2\text{O}_3 \cdot \text{H}_2\text{O}$) Corundum (Al_2O_3)
	Zn	Zincite (ZnO)

	Cu	Cuprite (Cu_2O)
Carbonates	Ca	Calcite (CaCO_3)
	Mg	Magnesite (MgCO_3)
	Cu	Malachite [$\text{CuCO}_3 \cdot \text{Cu}(\text{OH})_2$]
	Zn	Calamine (ZnCO_3)
	Fe	Siderite (FeCO_3)
Sulphides	Fe	Iron pyrite (FeS_2)
	Cu	Copper glance (Cu_2S)
	Hg	Cinnabar (HgS)
	Zn	Zinc blende (ZnS)
	Pb	Galena (PbS)
	Ag	Argentite or Silver glance (Ag_2S)
Halides	Na	Common salt or Rock salt (NaCl)
	Al	Cryolite (Na_3AlF_6)
	K, Mg	Carnallite ($\text{KCl} \cdot \text{MgCl}_2 \cdot 6\text{H}_2\text{O}$)
	Ag	Horn silver (AgCl)

EXTRACTION OF METALS



Conversion of ore into metal oxide

Calcination

- Heating of ore below its fusion temperature in absence of air.
- For carbonate and hydroxide ores.
- In reverberatory furnace. e.g., $\text{ZnCO}_3 \xrightarrow{\Delta} \text{ZnO} + \text{CO}_2 \uparrow$
Calamine Zinc oxide

Roasting

- Ore is heated in presence of air.
- For sulphide ores.
- In reverberatory or blast furnace. e.g., $2\text{ZnS} + 3\text{O}_2 \xrightarrow{\Delta} 2\text{ZnO} + 2\text{SO}_2 \uparrow$


Reduction of metallic oxide to free metal

Pyrometallurgy

Heating of metal oxide with a suitable reducing agent.

Smelting

Reduction of oxides with carbon.



Self reduction (Auto-reduction)

Sulphide ores of less electropositive metals like Hg, Pb, Cu, etc. are heated in air.

By aluminium (Goldschmidt aluminothermite process)

Reduction of metal oxide to metal by aluminium powder. $\text{Cr}_2\text{O}_3 + 2\text{Al} \longrightarrow 2\text{Cr} + \text{Al}_2\text{O}_3$

Electrolytic reduction

The oxides, hydroxides or chlorides of highly electropositive metals like Na, K, Mg, Ca and Al are extracted by electrolytic method, in their fused state. Electrolysis of this method is based on equation $\Delta G^\circ = -nFE^\circ$.

CONCENTRATION OF ORE

Froth floatation

- Based on preferential wetting of ore particles by oil (pine oil).
- Ore particles become light and form froth.
- Froth can be stabilised by stabilisers (aniline or cresols).


Electromagnetic separation

- Conveyor belt moving over two rollers, one of which is magnetic in nature.
- Magnetic ores are attracted by the magnetic roller.

Leaching


- It involves the treatment of the powdered ore with a suitable reagent so as to make it soluble while impurities remain insoluble.

Leaching of bauxite (Baeyer's process) :




Leaching of silver or gold

(Mac-Arthur Forest cyanide process) :




THERMODYNAMIC PRINCIPLES IN EXTRACTION OF METALS

- The free energy change (ΔG), occurring during the reduction processes help in deciding the suitable method for reduction, is given by $\Delta G = \Delta H - T\Delta S$; where, ΔH = enthalpy change; ΔG = Gibbs free energy change; T = temperature; ΔS = entropy change.
- Greater the negative value of ΔG , higher is the reducing power of an element.
- For the reduction of a metal oxide with a reducing agent, the plot of ΔG° vs temperature is studied, which is called Ellingham diagram.




CHARACTERISTICS OF ELLINGHAM DIAGRAM

- ΔG° becomes more positive when temperature increases, *i.e.*, stability of oxides decreases.
- A metal will reduce the oxide of other metals which lie above it in Ellingham diagram, *i.e.*, the metals for which the free energy of formation (ΔG°) of their oxides is more negative can reduce those metal oxides which has less negative ΔG° . Thus, Al reduces FeO, CrO and NiO in thermite reduction but it will not reduce MgO at temperature below 1773 K.
- CO is more effective reducing agent below 1073 K and above 1073 K coke is more effective reducing agent, *e.g.*, CO reduces Fe_2O_3 below 1073 K but above it, coke reduces Fe_2O_3 .



Extraction of Al and Zn



INFOSHOTS

Perovskite solar cells !

A perovskite is an unique crystal structure, consisting of formamidinium with multiple cations and mixed halide anions. A perovskite solar cell (PSC) is a type of solar cell, which includes the perovskite structured compound, most commonly a hybrid organic-inorganic lead or tin halide based material, as the light harvesting active layer. These solar cells not only show relatively high photovoltaic energy conversion efficiencies (above 22%), but can be also easily fabricated using cheap inorganic-organic perovskite compounds.

p-BLOCK ELEMENTS (GROUP 15 TO 18)

Group-15

Electronic configuration : ns^2np^3 ; ($n = 2$ to 6)

Metallic nature : Increases with increase in atomic number.

Ionic/Atomic radii : Increase down the group.

Ionisation energy : Decreases down the group. For an element; $I.E._1 < I.E._2 < I.E._3$

Allotropy : All elements except N and Bi, show allotropy.

Oxidation states : +3 and +5; N and P also show (-3) oxidation state.

Electronegativity : Decreases down the group.

Group-16

Electronic configuration : ns^2np^4 ; ($n = 2$ to 6)

Metallic nature : Increases with increase in atomic number.

Ionic/Atomic radii : Smaller than group-15 elements and increase down the group.

Ionisation energy : Lower than group-15 elements (due to half filled configuration of group-15 elements) and increases down the group.

Allotropy : All elements exhibit allotropy.

Electronegativity : Higher than group-15 elements and decreases down the group. O is second most electronegative element.

Oxidation states : +2, +4

Oxygen commonly shows -2 oxidation state.

Group-17

Electronic configuration : ns^2np^5 ; ($n = 2$ to 6)

Oxidising nature : Standard reduction potential of halogens are positive and decreases from fluorine to iodine.

F_2 is strongest oxidising agent while I_2 is the weakest.

Ionic/Atomic radii : Halogens have lowest atomic radii in their respective period which increase down the group.

Ionisation energy : They have very high ionisation energy and decreases down the group.

Electronegativity : Halogens are highly electronegative.

Oxidation states : F (-1 only)

Cl, Br, I show -1, +1, +3, +5, +7 oxidation states.

Group-18

Electronic configuration : ns^2np^6 ; ($n = 2$ to 6), (stable, fully filled, very less reactive).


Atomic radii : Exceptionally high as atomic radii correspond to van der Waals' radii.

Ionisation enthalpy : High value due to stable fully filled configuration.

Adsorption over charcoal : Except helium, these gases can be adsorbed by coconut charcoal at suitable temperatures.

PHYSICAL PROPERTIES

CHEMICAL PROPERTIES



Oxyacids	Group-15					
	Oxyacids of N	$H_2N_2O_2$ (+1) (Hyponitrous acid)	HNO_2 (+3) (Nitrous acid)	HNO_3 (+5) (Nitric acid)		
	Oxyacids of P	H_3PO_2 (+1) (Monobasic) Phosphinic acid	H_3PO_3 (+3) (Dibasic) Orthophosphorous acid	$H_4P_2O_6$ (+4) (Tetrabasic) Hypophosphoric acid		
		H_3PO_4 (+5) (Tribasic) Orthophosphoric acid	HPO_3 (+5) (Monobasic) Metaphosphoric acid	$H_4P_2O_7$ (+5) (Tetrabasic) Pyrophosphoric acid		
Oxyacids of As and Bi	H_3AsO_3 (+3) Arsenious acid	H_3AsO_4 (+5) Arsenic acid	$HBiO_3$ (+5) Metabismuthic acid			
Group-16						
Oxyacids of S	H_2SO_3 (+4) (Dibasic) Sulphurous acid	H_2SO_4 (+6) (Dibasic) Sulphuric acid	$H_2S_2O_3$ (+6, -2) (Dibasic) Thiosulphuric acid			
$H_2S_2O_7$ (+6) (Dibasic) Pyrosulphuric acid	$H_2S_2O_6$ (+6) (Dibasic) Dithionic acid	$H_2S_2O_8$ (+6) (Dibasic) Peroxydisulphuric acid	H_2SO_5 (+6) (Dibasic) Peroxomonosulphuric acid			
Group-17						
Oxyacids of F	HO_2F (+1) (Hypofluorous acid)	—	—	—		
Oxyacids of Cl	$HOCl$ (+1) (Hypochlorous acid)	$HClO_2$ (+3) Chlorous acid	$HClO_3$ (+5) Chloric acid	$HClO_4$ (+7) Perchloric acid		
Oxyacids of Br	$HOBr$ (+1) (Hypobromous acid)	—	$HBrO_3$ (+5) Bromic acid	$HBrO_4$ (+7) Perbromic acid		
Oxyacids of I	HOI (+1) Hypoiodous acid	—	HIO_3 (+5) Iodic acid	HIO_4 (+7) Periodic acid		

Hydrides

NH ₃	PH ₃	AsH ₃	SbH ₃	BiH ₃
H ₂ O	H ₂ S	H ₂ Se	H ₂ Te	
HF	HCl	HBr	HI	

→

Bond length increases
Thermal stability decreases
Acidic strength increases
Reducing nature increases


ABNORMAL BEHAVIOUR

N, O and F show different properties than their respective period elements due to

- small size
- high electronegativity
- absence of vacant *d*-orbitals.

Oxides of Nitrogen

Formula	Resonance Structures	Bond Parameters
N ₂ O		$\text{N}\equiv\text{N}-\text{O}$ 113 pm 119 pm (Linear)
NO		$:\text{N}\equiv\text{O}:$ 115 pm (Linear)
N ₂ O ₃		 (Planar)
NO ₂		 (Angular)
N ₂ O ₄		 (Planar)
N ₂ O ₅		 (Planar)


IMPORTANT COMPOUNDS**Dinitrogen (N₂)****Preparation :**

- **Properties :** Colourless, odourless, tasteless, non-toxic, unreactive due to triple bond; $\text{N}\equiv\text{N}$.

- **Uses :** For manufacturing of HNO₃, NH₃, etc.

Ammonia (NH₃)

- **Preparation :** By Haber's process,



- **Properties :** Its aqueous solution is basic in nature.
 $\text{NH}_3 + \text{H}_2\text{O} \longrightarrow \text{NH}_4^+ + \text{OH}^-$

- **Uses :** For manufacturing of nitrogenous fertilizers, HNO₃, Na₂CO₃, etc.

Titrimetric analysis has a wide applications in the various fields. It is used to determine the acid content of alcoholic beverages, calcium content in dairy products, total water hardness, vitamin C in fruit juices, etc.

Titrimetric / Volumetric Analysis

Simple Titration

- Acid-base titration :** The concentration of an acid is estimated by adding a solution of standard base. The equivalence point is detected by adding a few drops of suitable indicator to the acid solution.
- Redox titration :** The concentration of oxidant is estimated by adding reductant or vice versa. For example, Fe^{2+} ions can be estimated by titrating against acidified KMnO_4 solution.
- Precipitation titration :** Cations and anions combine to form a compound of very low solubility. $\text{AgNO}_3 + \text{NaCl} \rightarrow \text{AgCl} + \text{NaNO}_3$ (White ppt.)
- Complexometric titration :** Titrate combines with the titrant to form a complex salt. $\text{CuSO}_4 + 4\text{NH}_4\text{OH} \rightarrow [\text{Cu}(\text{NH}_3)_4]\text{SO}_4 + 4\text{H}_2\text{O}$

If a mixture of NaOH & NaHCO₃ is taken

- Equivalents of NaOH = equivalents of HCl required upto phenolphthlein end point.
- Equivalents of NaHCO_3 = equivalents of HCl required from phenolphthlein to methyl orange end point.

If a mixture of Na₂CO₃ & NaHCO₃ is taken

- 1/2 Equivalents of Na_2CO_3 ($n = 2$) = equivalents of HCl required upto phenolphthalein end point.
- 1/2 Equivalents of Na_2CO_3 ($n = 2$) + equivalents of NaHCO_3 ($n = 1$) initially = Equivalents of HCl required from phenolphthalein end point to methyl orange end point.

Back Titration

Reactant A of known concentration is titrated with reactant B (V_1 litre) of known concentration. Equivalents of B reacted with A = $N_1 V_1$. In another flask, reactant C of unknown concentration is reacted with A and excess of A is titrated with B (V_2 litre). Equivalents of B reacted with excess of A = $N_1 V_2$. Equivalents of pure C = $N_1 V_1 - N_1 V_2$. Moles of pure C = $\frac{N_1 V_1 - N_1 V_2}{n}$

Here, n-factor

For acids, n = basicity of acid

For bases, n = acidity of base

In case of redox reactions:

n -factor = change in oxidation number per molecule = total number of electrons transferred per mole

Mass of pure C = $\frac{N_1 V_1 - N_1 V_2}{n} \times \text{Molar mass of C}$

% Purity of C = $\frac{N_1 V_1 - N_1 V_2}{n} \times \frac{\text{Molar mass of C}}{w} \times 100$

where, N_1 = Normality of B, w = Mass of C

In the given calculation, we have taken alkali mixture of NaOH and Na_2CO_3 but other alkali mixtures can also be taken.

If a mixture of NaOH, Na₂CO₃ & NaHCO₃ is taken


- Equivalents of NaOH + 1/2 equivalents of Na_2CO_3 ($n = 2$) = Equivalents of HCl required upto phenolphthalein end point
- 1/2 equivalents of Na_2CO_3 ($n = 2$) + Equivalents of NaHCO_3 initially ($n = 1$) = Equivalents of HCl required from phenolphthalein end point to methyl orange end point.

Double Titration

The purpose of double titration is to determine the percentage composition of an alkali mixture or an acid mixture. **Process:**

- Dissolve mixture of Na_2CO_3 and NaOH in water to make alkaline solution, add two indicators phenolphthalein and methyl orange.
- Now this solution is titrated against HCl.

Reactions Taking Place



Calculations

At methyl orange end point:

- Moles of HCl consumed by NaHCO_3 = $M_1 V_2$ = Moles of NaHCO_3 formed from Na_2CO_3 = Moles of Na_2CO_3 in the mixture
- Mass of Na_2CO_3 in the mixture = $M_1 V_2 \times 106$
- % of Na_2CO_3 in mixture = $\frac{M_1 V_2 \times 106}{w} \times 100$


At phenolphthalein end point:

- Moles of HCl used in eq. (i) = $M_1 V_1 - M_1 V_2$ = Moles of NaOH
- Mass of NaOH = $(M_1 V_1 - M_1 V_2) \times 40$
- % of NaOH = $\frac{(M_1 V_1 - M_1 V_2) \times 40}{w} \times 100$ where, V_1 = volume of HCl used when phenolphthalein changes colour, V_2 = volume of HCl required from phenolphthalein colour change to methyl orange colour change w = Mass of Na_2CO_3 -NaOH mixture


Redox reactions are very important in day to day life, these found vast applications in photosynthesis, production of chemicals, extraction of metals, electrochemical cells and quantitative analysis. Some important reduction processes are discussed here.

Reduction Processes

Reduction of Oxygen Containing Compounds




Reduction of Unsaturated Compounds




Important Named Reduction Reactions and Reagents

- $\text{Zn}-\text{Hg}/\text{H}_2\text{O} \Rightarrow$ Clemmensen reduction
- $\text{NH}_2-\text{NH}_2/\text{H}_2\text{O} \Rightarrow$ Wolff-Kishner reduction
- $\text{Al}(\text{O}-\text{CH}(\text{CH}_3)_2)_3 + \text{CH}_3\text{CH}(\text{OH})-\text{CH}_3 \Rightarrow$ Meerwein-Ponndorf-Verley reduction (MPV Reduction)
- DIBAL-H \Rightarrow Di-iso-butyl aluminium hydride (Highly reactive, hard to control, explodes on heating, grinding or on exposure to water)
- $\text{H}_2[\text{SnCl}_4] \Rightarrow$ Stephen's reduction (RCN to RCHO)
- $\text{Pd} + \text{BaSO}_4 + \text{quinoline}$ in boiling xylene \Rightarrow Lindlar's catalyst
- Reduction of alkynes to *trans*-alkene using $\text{Li}, \text{Na}/\text{Liq. NH}_3 \Rightarrow$ Birch reduction


Reduction to Alcohol




Reduction of Benzoyl Chloride




Reduction of Epoxide




Reduction of N-containing Compounds



Reduction of Halides



Reduction of Unsaturated Aldehydes



Nitric Acid (HNO₃)

- **Preparation** : By Ostwald process,

$$4\text{NH}_3 + 5\text{O}_2 \xrightarrow[500\text{ K, 9 bar}]{\text{Pt, Rh gauge}} 4\text{NO} + 6\text{H}_2\text{O}$$

$$2\text{NO} + \text{O}_2 \rightleftharpoons 2\text{NO}_2$$

$$3\text{NO}_2 + \text{H}_2\text{O} \longrightarrow 2\text{HNO}_3 + \text{NO}$$
- **Properties** : It is a strong oxidising agent as it decomposes to give nascent oxygen easily.
- **Oxidation of non-metals** :

$$2\text{HNO}_3 (\text{conc.}) \longrightarrow \text{H}_2\text{O} + 2\text{NO}_2 + [\text{O}]$$

$$2\text{HNO}_3 (\text{dil.}) \longrightarrow \text{H}_2\text{O} + 2\text{NO} + 3[\text{O}]$$

$$\text{S}_8 + 48\text{HNO}_3 \longrightarrow 8\text{H}_2\text{SO}_4 + 48\text{NO}_2 + 16\text{H}_2\text{O}$$

$$\text{C} + 4\text{HNO}_3 \longrightarrow \text{CO}_2 + 4\text{NO}_2 + 2\text{H}_2\text{O}$$

$$\text{P}_4 + 20\text{HNO}_3 \longrightarrow 4\text{H}_3\text{PO}_4 + 20\text{NO}_2 + 4\text{H}_2\text{O}$$

$$\text{I}_2 + 10\text{HNO}_3 \longrightarrow 2\text{HIO}_3 + 10\text{NO}_2 + 4\text{H}_2\text{O}$$
- The concentrated nitric acid renders metals like iron, cobalt, nickel, chromium, etc passive.
- $\text{Metal} + \text{HNO}_3 \longrightarrow \text{Metal nitrate} + [\text{H}]$
- Causes nitration of organic compounds.

Phosphorus (P₄)

- Discrete tetrahedral
- Very reactive
- Translucent waxy solid

- Iron grey luster
- Less reactive than white P
- Polymeric structure (P₄ units linked)

- Two forms $\left\{ \begin{array}{l} \alpha\text{-black P (at 803 K)} \\ \beta\text{-black P (at 473 K)} \end{array} \right.$
- Very less reactive
- Opaque monoclinic or rhombohedral crystals


Oxygen	Ozone
$:\ddot{\text{O}}=\ddot{\text{O}}:$	$ \begin{array}{c} 1.278 \text{ \AA} \\ \diagdown \quad \diagup \\ \text{:}\ddot{\text{O}}\text{---}\text{O}\text{---}\ddot{\text{O}}\text{:} \\ \diagup \quad \diagdown \\ 1.278 \text{ \AA} \\ 116.8^\circ \end{array} $
Laboratory Preparation	
$2\text{KClO}_3 \xrightarrow{420\text{ K}} 2\text{KCl} + 3\text{O}_2$	$3\text{O}_2 \rightleftharpoons 2\text{O}_3 - \text{energy}$ 5-10% ozone mixture is obtained, known as ozonised oxygen.
Properties	
$2\text{H}_2 + \text{O}_2 \xrightarrow[\text{or electric discharge}]{1073\text{ K}} 2\text{H}_2\text{O}$ $2\text{NO} + \text{O}_2 \longrightarrow 2\text{NO}_2$ $\text{P}_4 + 5\text{O}_2 \longrightarrow \text{P}_4\text{O}_{10}$ $4\text{Al} + 3\text{O}_2 \longrightarrow 2\text{Al}_2\text{O}_3$	$\text{O}_3 + 2\text{H}^+ + 2\text{e}^- \longrightarrow \text{O}_2 + \text{H}_2\text{O}; E^\circ = +2.07\text{ V}$ $2\text{HCl} + \text{O}_3 \longrightarrow \text{H}_2\text{O} + \text{Cl}_2 + \text{O}_2$ $\text{PbS} + 4\text{O}_3 \longrightarrow \text{PbSO}_4 + 4\text{O}_2$ $\text{I}_2 + \text{H}_2\text{O} + 5\text{O}_3 \longrightarrow 2\text{HIO}_3 + 5\text{O}_2$ Coloured substance + [O] \longrightarrow Colourless substance (Bleaching action)

INFOSHOTS

van der Waals' forces !

Very weak van der Waals' forces between individual atoms are measured for the first time. Individual noble gas atoms were fixed within a molecular network and the interactions with a single xenon atom were determined that was positioned at the tip of an atomic force microscope. As expected, the forces varied according to the distance between the two atoms, but in some cases, the forces were several times larger than theoretically calculated.

Allotropic Forms of Sulphur



Sulphuric Acid

- Lead chamber process :**

$$2\text{SO}_2 + \text{O}_2 + 2\text{H}_2\text{O} \xrightarrow[\text{(Catalyst)}]{\text{NO}} 2\text{H}_2\text{SO}_4$$
- Contact process :**

$$2\text{SO}_2 + \text{O}_3 \xrightarrow{\text{Catalyst}} 2\text{SO}_3$$

$$\text{H}_2\text{SO}_4 + \text{SO}_3 \longrightarrow \text{H}_2\text{S}_2\text{O}_7 \xrightarrow{+\text{H}_2\text{O}} 2\text{H}_2\text{SO}_4$$

(oleum)
- Properties**
 - As oxidising agent (due to nascent O) :**

$$\text{H}_2\text{SO}_4 \xrightarrow{444^\circ\text{C}} \text{H}_2\text{O} + \text{SO}_2 + [\text{O}]$$

$$\text{C} + 2\text{H}_2\text{SO}_4 \longrightarrow \text{CO}_2 + 2\text{SO}_2 + 2\text{H}_2\text{O}$$

$$\text{S}_8 + 16\text{H}_2\text{SO}_4 \longrightarrow 24\text{SO}_2 + 16\text{H}_2\text{O}$$

$$\text{P}_4 + 10\text{H}_2\text{SO}_4 \longrightarrow 4\text{H}_3\text{PO}_4 + 10\text{SO}_2 + 4\text{H}_2\text{O}$$

$$2\text{KI} + 2\text{H}_2\text{SO}_4 \longrightarrow \text{K}_2\text{SO}_4 + \text{SO}_2 + \text{I}_2 + 2\text{H}_2\text{O}$$
 - As dibasic acid :**

$$\text{Na}_2\text{CO}_3 + \text{H}_2\text{SO}_4 \longrightarrow \text{Na}_2\text{SO}_4 + \text{H}_2\text{O} + \text{CO}_2$$

$$2\text{NaCl} + \text{H}_2\text{SO}_4 \longrightarrow \text{Na}_2\text{SO}_4 + 2\text{HCl}$$

Bleaching Powder ($\text{CaOCl}_2 \cdot \text{H}_2\text{O}$)

- Preparation :**

$$\text{Ca}(\text{OH})_2 + \text{Cl}_2 \longrightarrow \text{CaOCl}_2 + \text{H}_2\text{O}$$

Dry slaked lime
Preparation is carried out in Hasenclever's plant and Beckmann's plant.
- Properties**

$$6\text{CaOCl}_2 \xrightarrow{\text{On long standing}} \text{Ca}(\text{ClO}_3)_2 + 5\text{CaCl}_2$$
- Oxidising properties :**

$$\text{CaOCl}_2 + \text{H}_2\text{S} \longrightarrow \text{CaCl}_2 + \text{H}_2\text{O} + \text{S}$$

$$3\text{CaOCl}_2 + 2\text{NH}_3 \longrightarrow 3\text{CaCl}_2 + 3\text{H}_2\text{O} + \text{N}_2$$
- Bleaching action :**
 Coloured matter + [O] \longrightarrow Colourless product

INTERHALOGEN COMPOUNDS

Type XX'_1 (Linear shape)	Type XX'_3 (T-shape)	Type XX'_5 (Square pyramidal shape)	Type XX'_7 (Pentagonal bipyramidal shape)
ClF	ClF ₃	ClF ₅	
BrF, BrCl	BrF ₃	BrF ₅	
ICl, IBr, IF	ICl ₃ , IF ₃	IF ₅	IF ₇

COMPOUNDS OF XENON

Molecule	Electron pairs	Hybridisation	Shape
XeF ₂	5	sp^3d	Linear
XeF ₄	6	sp^3d^2	Square planar
XeF ₆	7	sp^3d^3	Distorted octahedral
XeOF ₂	5	sp^3d	T-shape
XeOF ₄	6	sp^3d^2	Square pyramidal
XeO ₃	4	sp^3	Pyramidal
XeO ₂ F ₂	5	sp^3d	See-saw
XeO ₄	4	sp^3	Tetrahedral

Your favourite MTG Books/Magazines available in WEST BENGAL at

- Progressive Book Centre - Kharagpur
Ph: 03222-279956; Mob: 9932619526, 9434192998
- International Book Trust - Kolkata
Ph: 033-22414947, 24792343; Mob: 9830360012
- Rukmani Agencies - Kolkata Ph: 033-24666173, 224839473; Mob: 9830037811
- Every Books - Kolkata
Ph: 033-22418590, 22194699; Mob: 9830162977, 8599985926
- Katha - Kolkata Ph: 033-22196313; Mob: 9830257999
- Saraswati Book Store - Kolkata Ph: 22198108, 22190784; Mob: 9831349473
- Chhatra Pustak Bhawan - Medinipur Mob: 9609996099, 9332341750
- Novelty Books - Siliguri Ph: 0353-2525445; Mob: 7797938077
- Om Traders - Siliguri Ph: 0353-6450299; Mob: 9434310035, 9749048104

Visit "MTG IN YOUR CITY" on www.mtg.in to locate nearest book seller OR write to info@mtg.in OR call **0124-6601200** for further assistance.

SPEED PRACTICE


- From the following information,
 $X + \text{H}_2\text{SO}_4 \longrightarrow Y$ (a colourless and irritating gas)
 $Y + \text{K}_2\text{Cr}_2\text{O}_7 + \text{H}_2\text{SO}_4 \longrightarrow$ green coloured solution
 Identify the pair X and Y.
 (a) Cl^- , HCl (b) SO_3^{2-} , SO_2
 (c) S^{2-} , H_2S (d) CO_3^{2-} , CO_2
- In the commercial electrochemical process for aluminium extraction, the electrolyte used is
 (a) $\text{Al}(\text{OH})_3$ in NaOH solution
 (b) an aqueous solution of $\text{Al}_2(\text{SO}_4)_3$
 (c) a molten mixture of Al_2O_3 and Na_3AlF_6
 (d) a molten mixture of $\text{AlO}(\text{OH})$ and $\text{Al}(\text{OH})_3$
- Which of the following is correct statement?
 (a) Phosphates have no biological significance in humans.
 (b) Between nitrates and phosphates, phosphates are less abundant in earth's crust.
 (c) Between nitrates and phosphates, nitrates are less abundant in earth's crust.
 (d) Oxidation of nitrates is possible in soil.
- Composition of azurite mineral is
 (a) $\text{CuCO}_3 \cdot \text{CuO}$
 (b) $\text{Cu}(\text{HCO}_3)_2 \cdot \text{Cu}(\text{OH})_2$
 (c) $2\text{CuCO}_3 \cdot \text{Cu}(\text{OH})_2$
 (d) $\text{CuCO}_3 \cdot 2\text{Cu}(\text{OH})_2$
- The number of $\text{S}=\text{O}$ and $\text{S}-\text{OH}$ bonds present in peroxodisulphuric acid and pyrosulphuric acid are respectively
 (a) (4 and 2) and (4 and 2)
 (b) (2 and 4) and (2 and 4)
 (c) (4 and 2) and (2 and 4)
 (d) (2 and 2) and (2 and 2)

(JEE Main Online 2017)
- In the construction of aircrafts, an alloy, elektron is often used. Its composition is
 (a) 15% Mg and 85% Al
 (b) 95% Mg and 5% Zn
 (c) 95% Mg and 5% Al
 (d) 80% Al and 20% Zn
- Which of the following is not correct?
 (a) $3\text{O}_2 \xrightleftharpoons[\text{discharge}]{\text{Silent electric}} 2\text{O}_3$; $\Delta H = -284.5 \text{ kJ mol}^{-1}$
 (b) Ozone undergoes addition reaction with unsaturated carbon compounds.
 (c) Sodium thiosulphate reacts with I_2 to form sodium tetrathionate and sodium iodide.
 (d) Ozone oxidises lead sulphide to lead sulphate.
- The process is used to remove reducible oxides from metals. Molten impure metal is stirred by a wooden rods. The hydrocarbon gases reduce the oxides. The process is called
 (a) zone refining (b) cupellation
 (c) poling (d) leaching.
- Roasted copper pyrite on smelting with sand produces
 (a) FeSiO_3 as fusible slag and Cu_2S as matte
 (b) CaSiO_3 as infusible slag and Cu_2O as matte
 (c) $\text{Ca}_3(\text{PO}_4)_2$ as fusible slag and Cu_2S as matte
 (d) $\text{Fe}_3(\text{PO}_4)_2$ as infusible slag and Cu_2S as matte.
- The correct statement(s) about the oxoacids, HClO_4 and HClO , is(are)
 (a) the conjugate base of HClO_4 is weaker base than H_2O
 (b) the central atom in both HClO_4 and HClO is sp^3 hybridized
 (c) HClO_4 is formed in the reaction between Cl_2 and H_2O
 (d) HClO_4 is more acidic than HClO because of the resonance stabilization of its anion.

(JEE Advanced 2017)
- Hydrolysis of XeF_4 and CaNCN give respectively
 (a) XeO_3 and CaCO_3 (b) XeO_2 and CaCN_2
 (c) XeO_3 and CaCN_2 (d) XeOF_2 and CaCO_3
- Xenon hexafluoride reacts with potassium fluoride to yield
 (a) $[\text{XeF}_4]^{2+} [\text{KF}_3]^{2-}$ (b) XeF_4
 (c) $[\text{XeF}_5]^+ [\text{KF}_2]^-$ (d) $\text{K}^+ [\text{XeF}_7]^-$

13. Pb, which is present as impurity in silver, is removed by
 (a) Parke's process
 (b) Pattinson process
 (c) cupellation
 (d) addition of Zn in molten silver.

14. Which of the following diagrams is correctly related to the extraction of Mg from sea water?




15. When copper is heated with conc. HNO_3 it produces
 (a) $\text{Cu(NO}_3)_2$, NO and NO_2
 (b) $\text{Cu(NO}_3)_2$ and N_2O
 (c) $\text{Cu(NO}_3)_2$ and NO_2
 (d) $\text{Cu(NO}_3)_2$ and NO (NEET 2016)

16. Out of $\text{H}_2\text{S}_2\text{O}_3$, $\text{H}_2\text{S}_2\text{O}_6$, H_2SO_5 and $\text{H}_2\text{S}_2\text{O}_8$ peroxy acids are
 (a) $\text{H}_2\text{S}_2\text{O}_3$, $\text{H}_2\text{S}_4\text{O}_6$ (b) $\text{H}_2\text{S}_4\text{O}_6$, H_2SO_5
 (c) H_2SO_5 , $\text{H}_2\text{S}_2\text{O}_8$ (d) $\text{H}_2\text{S}_2\text{O}_3$ and $\text{H}_2\text{S}_2\text{O}_8$

17. The process of pickling is to
 (a) cover metal surface with a less reactive metal
 (b) electroplate a metal with a non-metal
 (c) cool a red hot steel by dipping it in oil
 (d) remove layers of basic oxides from metal surface before electroplating.


18. Which series of reactions correctly represents chemical relations related to iron and its compound?



(JEE Main 2014)

19. Pitch blende is
 (a) ore of Ti (b) ore of U
 (c) ore of W (d) ore of B.

20. Which is incorrectly matched?



21. The chemical composition of slag formed during the smelting process in the extraction of copper is




22. The non-metal that does not exhibit positive oxidation state is




(JEE Main Online 2016)

23. Which of the following allotropic forms of phosphorus contain given structure?




- (a) White phosphorus (b) Red phosphorus
 (c) Black phosphorus (d) Yellow phosphorus

24. Which of the following is correct?



25. In the extraction of copper from its sulphide ore, the metal is finally obtained by the reduction of cuprous oxide with



(NEET 2015)

MPP-5 CLASS XI

ANSWER KEY

- | | | | | | | | | | |
|-----|-----------|-----|---------|-----|-------|-----|-----|-----|-------|
| 1. | (a) | 2. | (a) | 3. | (d) | 4. | (a) | 5. | (c) |
| 6. | (d) | 7. | (b) | 8. | (c) | 9. | (d) | 10. | (b) |
| 11. | (d) | 12. | (a) | 13. | (a) | 14. | (a) | 15. | (c) |
| 16. | (b) | 17. | (a) | 18. | (b) | 19. | (b) | 20. | (a,d) |
| 21. | (a,b,c,d) | 22. | (a,b,c) | 23. | (a,b) | 24. | (4) | 25. | (7) |
| 26. | (3) | 27. | (d) | 28. | (a) | 29. | (b) | 30. | (b) |

26. When an inorganic compound reacts with SO_2 in aqueous medium produces (A). (A) on reaction with Na_2CO_3 gives the compound (B) which with sulphur gives a substance (C) used in photography. The compound (C) is

- (a) $\text{Na}_2\text{S}_2\text{O}_3$ (b) Na_2SO_4
(c) Na_2S (d) $\text{Na}_2\text{S}_2\text{O}_7$

27. In which pair of ions both the species contain S — S bond?

- (a) $\text{S}_4\text{O}_6^{2-}$, $\text{S}_2\text{O}_3^{2-}$ (b) $\text{S}_2\text{O}_7^{2-}$, $\text{S}_2\text{O}_8^{2-}$
(c) $\text{S}_4\text{O}_6^{2-}$, $\text{S}_2\text{O}_7^{2-}$ (d) $\text{S}_2\text{O}_7^{2-}$, $\text{S}_2\text{O}_3^{2-}$

(NEET 2017)

28. Compounds formed when the noble gases get entrapped in the cavities of crystal lattices of certain organic and inorganic compounds are known as

- (a) interstitial compounds
(b) clathrates
(c) hydrates (d) picrates.

29. Cassiterite is concentrated by


- (a) levigation
(b) electromagnetic separation
(c) floatation
(d) liquefaction.

30. In which of the following reactions HNO_2 acts as an oxidising agent?

- (a) $\text{MnO}_4^- + \text{NO}_2^- + \text{H}^+ \longrightarrow \text{Mn}^{2+} + \text{NO}_3^- + \text{H}_2\text{O}$
(b) $\text{KI} + \text{H}_2\text{SO}_4 + \text{HNO}_2 \longrightarrow \text{K}_2\text{SO}_4 + \text{NO} + \text{I}_2 + \text{H}_2\text{O}$
(c) $\text{Cl}_2 + \text{HNO}_2 \longrightarrow \text{HCl} + \text{NO}_2$
(d) $\text{O}_3 + \text{NO}_2^- + \text{H}^+ \longrightarrow \text{O}_2 + \text{NO}_3^- + \text{H}_2\text{O}$

SOLUTIONS

1. (b): $\text{SO}_3^{2-} + 2\text{H}^+ \longrightarrow \text{SO}_2 + \text{H}_2\text{O}$




2. (c): The electrolysis of pure alumina faces some difficulties. Pure alumina is a bad conductor of electricity. The fusion temperature of pure alumina is about 2000°C . At this temperature, when electrolysis is carried out on the fused mass, the metal formed vaporises, as the boiling point of aluminium is 1800°C . These difficulties are overcome by using a mixture containing alumina, cryolite (Na_3AlF_6), and fluorspar (CaF_2).

3. (c): Between nitrates and phosphates, nitrates are more soluble in water and hence, are less abundant in earth's crust.

4. (c): Azurite is a basic copper carbonate ore having composition $2\text{CuCO}_3 \cdot \text{Cu}(\text{OH})_2$.


5. (a): Peroxodisulphuric acid :



No. of S = O bonds = 4

No. of S — OH bonds = 2


Pyrosulphuric acid :



No. of S = O bonds = 4 ; No. of S — OH bonds = 2

6. (c): Elektron is magnesium alloy containing 0% to 9.5% of some of the following elements in varying proportion aluminium (< 9.5%), yttrium (5.25%), silver (2.5%), etc.


7. (a): The given reaction is endothermic.



8. (c)

9. (a): During smelting, silica combines with FeO to form fusible FeSiO_3 slag and Cu_2S as matte.


10. (a,b,d) : Structures of HClO and HClO_4 are :



(a) HClO_4 is a stronger acid than H_3O^+ . Therefore, conjugate base of HClO_4 , i.e., ClO_4^- , is weaker base than H_2O .


(b) The hybridisation of central atom in both HClO and HClO_4 is sp^3 .

(c) Reaction of Cl_2 with water forms HOCl which decomposes to give nascent oxygen.




(d) HClO_4 is more acidic than HClO as ClO_4^- is more stable than ClO^- due to resonance.

11. (a): $6\text{XeF}_4 + 12\text{H}_2\text{O} \xrightarrow{\text{Partial hydrolysis}} 2\text{XeO}_3 + 24\text{HF} + 4\text{Xe} + 3\text{O}_2$







12. (d): XeF_6 behaves as F^- acceptor to form fluoroanions.




13. (c)

14. (b): Sea water contains Mg^{2+} ions.





15. (c): $\text{Cu} + 4\text{HNO}_3(\text{conc.}) \longrightarrow \text{Cu}(\text{NO}_3)_2 + 2\text{NO}_2 + 2\text{H}_2\text{O}$




17. (d)

18. (a): Combustion of Fe to give Fe_3O_4 which upon reduction with CO first gives FeO and finally Fe. Option (b) and (c) are wrong because FeSO_4 and $\text{Fe}_2(\text{SO}_4)_3$ on heating gives Fe_2O_3 , SO_2 , SO_3 and not Fe.




Option (d) is wrong because in presence of air, reduction of FeCl_3 to FeCl_2 cannot occur.

19. (b)




21. (b): When smelting is done in the blast furnace, most of the ferrous oxide is converted to ferric oxide. With silica, it forms ferrous silicate, which is the slag.








22. (c): Fluorine is the most electronegative element and it shows only -1 oxidation state.

23. (b): Polymeric form of $(\text{P}_4)_n$ is red phosphorus.

24. (c)



It is an example of auto-reduction.



28. (b)

29. (b): Cassiterite is tinstone (SnO_2) which is non-magnetic and contains wolframite, FeWO_4 (magnetic) impurities. These are separated by electromagnetic separation.

30. (b): HNO_2 oxidises KI to iodine (I_2).


Your favourite MTG Books/Magazines available in RAJASTHAN at

- Competition Book House - Alwar Ph: 0144-2338391; Mob: 9460607836
- Nakoda Book Depot - Bhilwara Ph: 01482-239653; Mob: 9214983594
- Alankar Book Depot - Bhiwadi Ph: 01493-222294; Mob: 9414707462
- Uttam Pustak Kendra - Bikaner Mob: 8955543195, 9414572625
- Yadu Books & Stationers - Bikaner Mob: 9251653481
- Goyal Books & Stationers - Jaipur Ph: 0141-2742454; Mob: 9414326406, 9929638435
- India Book House - Jaipur Ph: 0141-2314983, 2311191, 2651784; Mob: 9829014143, 9414079983
- Ravi Enterprises - Jaipur Ph: 0141-2602517, 2619958, 2606998; Mob: 9829060694
- Shri Shyam Pustak Mandir - Jaipur Ph: 0141-2317972; Mob: 9928450717
- Sarvodaya Book Stall - Jodhpur Ph: 0291-2653734, 35; Mob: 8107589141
- Bhandari Stationers - Kota Ph: 0744-2327576, 2391678; Mob: 9001094271, 9829038758
- Raj Traders - Kota Ph: 0744-2429090; Mob: 9309232829, 9214335300
- Vardhman Sports & Stationers - Kota Mob: 9461051901, 9351581238, 9828139717
- Jhuria Book Shop - Sikar Mob: 9784246419, 9460838235, 8432943550
- Popular Book Depot - Udaipur Ph: 2442881, 0487-2329436, 2421467; Mob: 9388513335, 9847922545

Visit "MTG IN YOUR CITY" on www.mtg.in to locate nearest book seller OR write to info@mtg.in OR call

0124-6601200 for further assistance.

EXAMINER'S MIND CLASS XII



The questions given in this column have been prepared strictly on the basis of NCERT Chemistry for Class XII. This year JEE (Main & Advanced)/NEET/AIIMS have drawn their papers heavily from NCERT books.

Section - I	Q. 1 to 10 Only One Option Correct Type MCQs.
Section - II	Q. 11 to 13 More than One Options Correct Type MCQs.
Section - III	Q. 14 to 17 Paragraph Type MCQs having Only One Option Correct.
Section - IV	Q. 18 & 19 Matching List Type MCQs having Only One Option Correct.
Section - V	Q. 20 to 22 Assertion Reason Type MCQs having Only One Option Correct. Mark the correct choice as : (a) If both assertion and reason are true and reason is the correct explanation of assertion. (b) If both assertion and reason are true but reason is not the correct explanation of assertion. (c) If assertion is true but reason is false. (d) If both assertion and reason are false.
Section - VI	Q. 23 to 25 Integer Value Correct Type Questions having Single Digit Integer Answer, ranging from 0 to 9 (both inclusive).

THE p-BLOCK ELEMENTS (GROUP 15 TO 18)

SECTION - I

Only One Option Correct Type


- Under normal conditions, which of the following hydrides is non-combustible?
(a) NH_3 (b) PH_3 (c) AsH_3 (d) SbH_3
- Aqueous solutions of hydrogen sulphide and sulphur dioxide when mixed together, yield
(a) sulphur and water
(b) sulphur trioxide and water
(c) hydrogen peroxide and sulphur
(d) hydrogen and sulphurous acid.
- Which among the following factors is most important in making fluorine the strongest oxidising agent?
(a) Electron affinity (b) Ionisation energy
(c) Hydration energy
(d) Bond dissociation energy
- The heat of vaporisation, polarisation and solubility in water of noble gases are in the order
(a) $\text{He} < \text{Ne} < \text{Ar} < \text{Kr} < \text{Xe}$
(b) $\text{Xe} < \text{Kr} < \text{Ar} < \text{Ne} < \text{He}$
(c) $\text{He} < \text{Ne} < \text{Ar} < \text{Xe} < \text{Kr}$
(d) $\text{He} < \text{Ne} \approx \text{Ar} < \text{Kr} \approx \text{Xe}$
- Sodium hydroxide solution reacts with phosphorus to give phosphine. To bring about this reaction, we need
(a) white phosphorus and dil. NaOH
(b) white phosphorus and conc. NaOH
(c) red phosphorus and dil. NaOH
(d) red phosphorus and conc. NaOH.
- Which of the following properties of H_2SO_4 in respective reaction is wrong?

$$\begin{array}{c} \text{COOH} \\ | \\ \text{COOH} \end{array} \xrightarrow{\text{H}_2\text{SO}_4} \text{CO} + \text{CO}_2 + \text{H}_2\text{O};$$
 H_2SO_4 acts as dehydrating agent.
(b) $\text{Cu} + 2\text{H}_2\text{SO}_4 \longrightarrow \text{CuSO}_4 + 2\text{H}_2\text{O} + \text{SO}_2;$
 H_2SO_4 acts as oxidising agent.
(c) $\text{Na}_2\text{S} + \text{H}_2\text{SO}_4 \longrightarrow \text{Na}_2\text{SO}_4 + \text{H}_2\text{S};$
 H_2SO_4 acts as an acid.
(d) $2\text{HBr} + \text{H}_2\text{SO}_4 \longrightarrow 2\text{H}_2\text{O} + \text{SO}_2 + \text{Br}_2;$
 H_2SO_4 acts as reducing agent.
- A black compound of manganese reacts with a halogen acid to give greenish yellow gas. When excess of this gas reacts with NH_3 and unstable trihalide is formed. In this process, the oxidation state of nitrogen changes from
(a) -3 to $+3$ (b) -3 to 0
(c) -3 to $+5$ (d) 0 to -3

JEE (ADVANCED)

mtg

Dry runs are here!



FEATURES:

- 17 years solved papers with detailed solutions
- 10 Model Test Papers
- Chapter-wise indexing of questions

₹475

Now, create your own pre-JEE. Just like pre-boards. With previous years' papers and model test papers for JEE (Advanced), complete with detailed solutions, identify your areas of weakness and work on addressing them in time. Multiple test papers ensure you do your dry runs again and again, till such time you feel confident of taking on the best. For it will indeed be the best you compete with in JEE (Advanced). So what are you waiting for? **Order MTG's JEE Advanced Explorer today.**



Scan now with your
smartphone or tablet
Application to read
QR codes required

Available at all leading book shops throughout India. To buy online visit www.mtg.in.

For more information or for help in placing your order, call 0124-6601200 or email: info@mtg.in

8. The stability of interhalogen compounds follows the order
 (a) $\text{IF}_3 > \text{BrF}_3 > \text{ClF}_3$ (b) $\text{BrF}_3 > \text{IF}_3 > \text{ClF}_3$
 (c) $\text{ClF}_3 > \text{BrF}_3 > \text{IF}_3$ (d) $\text{ClF}_3 > \text{IF}_3 > \text{BrF}_3$
9. The reaction between NH_2^- and N_2O gives
 (a) NO (b) N_2O_5
 (c) NH_2NH_2 (d) N_3^-
10. Which of the following statements regarding sulphur is incorrect?
 (a) S_2 molecule is paramagnetic.
 (b) The vapour at 200°C consists mostly of S_8 rings.
 (c) At 600°C the gas mainly consists of S_2 molecules.
 (d) The oxidation state of sulphur is never less than +4 in its compounds.

SECTION - II

More than One Options Correct Type




11. Which of the following statements are incorrect?
 (a) Dry slaked lime reacts with Cl_2 to give calcium chlorate.
 (b) With excess chlorine ammonia forms nitrogen trichloride.
 (c) Sulphur reacts with Cl_2 to give SCl_4 .
 (d) With hot and conc. NaOH, Cl_2 gives sodium chlorate.
12. Which statements regarding argon in solid state are correct?
 (a) Its atoms form a covalent network.
 (b) The forces between the molecules are weak van der Waals' forces.
 (c) Its molecules are monoatomic.
 (d) It is good conductor of electricity.
13. Ammonia on reaction with hypochlorite anion can form
 (a) NO (b) NH_4Cl (c) N_2H_4 (d) HNO_2

SECTION - III

Paragraph Type

Paragraph for Questions 14 and 15

Industrially sulphuric acid is produced by the following steps :



Since the reaction between SO_3 and H_2O is violent, therefore, SO_3 is passed into 98% H_2SO_4 to produce oleum ($\text{H}_2\text{S}_2\text{O}_7$).

14. Pure H_2SO_4 does not react with metal because
 (a) pure H_2SO_4 does not contain any water
 (b) pure H_2SO_4 acts as an oxidising agent
 (c) pure H_2SO_4 is strongly H-bonded
 (d) pure H_2SO_4 does not contain any SO_3 .
15. $\text{H}_2\text{SO}_4 + \text{PCl}_5 \longrightarrow (X) \xrightarrow{\text{H}_2\text{O}}$ two strong acids
 Therefore (X) is
 (a) SO_2Cl_2 (b) SOCl
 (c) ClSO_3H (d) POCl_3

Paragraph for Questions 16 and 17

The direct reaction of xenon with fluorine leads to a series of compounds with oxidation numbers +2, +4 and +6. XeF_4 reacts violently with water to give XeO_3 . The compounds of xenon exhibit rich stereochemistry and their geometries can be deduced considering the total number of electron pairs in the valence shell.

16. The structure of XeO_3 is
 (a) linear (b) planar
 (c) pyramidal (d) T-shaped.
17. XeF_4 and XeF_6 are expected to be
 (a) oxidising (b) reducing
 (c) unreactive (d) strongly basic.

SECTION - IV

Matching List Type

18. Match the List I with List II and select the correct answer using the codes given below the lists :

List I

- (P) P_4O_6
 (Q) P_4O_{10}
 (R) N_2O_3
 (S) N_2O_5
 (a) (P-i); (Q-ii, iv); (R-iii); (S-iv)
 (b) (P-ii); (Q-i, iv); (R-iii); (S-i, iv)
 (c) (P-i); (Q-i, iii); (R-iv); (S-i, ii, iii)
 (d) (P-ii); (Q-i, iii); (R-iv); (S-i)

List II

- (i) White crystalline solid
 (ii) Waxy solid with garlic smell
 (iii) Blue coloured liquid
 (iv) Coordinate bonds

19. Match the List I with List II and select the correct answer using the codes given below the lists :

List I

- (A) Helium
 (B) Argon
 (C) Neon
 (D) Xenon
 (a) A-q; B-s; C-p; D-r
 (c) A-r; B-q; C-s; D-p

List II

- (p) Ionisation energy comparable to O_2
 (q) Provides inert atmosphere in metallurgy
 (r) Cryogenic
 (s) For advertising sign
 (b) A-q; B-p; C-s; D-r
 (d) A-r; B-s; C-q; D-p

SECTION - V

Assertion Reason Type

20. **Assertion :** H_3PO_3 and H_3PO_4 are tribasic acids as they contain three hydrogen atoms each.
Reason : Both H_3PO_3 and H_3PO_4 are reducing in nature.
21. **Assertion :** Reaction of SO_2 and H_2S in the presence of Fe_2O_3 catalyst gives elemental sulphur.
Reason : In this reaction H_2S is acting as a reducing agent.

22. **Assertion :** HClO_4 is less acidic than HBrO_4 .
Reason : HClO_4 ionises less in water than HBrO_4 .

SECTION - VI

Integer Value Correct Type

23. What is the oxidation state of chlorine in HClO_4 ?
24. Number of P—OH bonds present in pyrophosphoric acid is
25. Amongst the following, the maximum number of isoelectronic molecules/ions are
 XeO_3 , BrO_3^- , ClF , XeF_2 , OF_2 , XeF_4 , ICl_4^- , ClO^- , IBr_2^-

ALCOHOLS, PHENOLS AND ETHERS

SECTION - I

Only One Option Correct Type


1. When 2,3-dimethyl-2-butanol undergoes acid catalysed dehydration the minor product is
 (a) 2,3-dimethyl-1-butene
 (b) 2,3-dimethyl-2-butene
 (c) 3,3-dimethyl-1-butene
 (d) none of these.
2. What is the function of diethyl ether in Grignard reagent preparation?
 1. To act as catalyst
 2. To act as solvent
 3. To provide lone pair of electrons for coordination
 4. To act as an acid
 (a) 1,2 (b) 2,3 (c) 3,4 (d) 2,4
3. Which of the following is not true in case of reaction with heated copper at 300°C ?
 (a) Phenol \longrightarrow Benzyl alcohol
 (b) Primary alcohol \longrightarrow Aldehyde
 (c) Secondary alcohol \longrightarrow Ketone
 (d) Tertiary alcohol \longrightarrow Olefin

4. The reaction of $\text{CH}_3\text{CH}=\text{CH}-\text{C}_6\text{H}_4-\text{OH}$ with HBr gives

- (a) $\text{CH}_3\text{CHBrCH}_2-\text{C}_6\text{H}_4-\text{OH}$
- (b) $\text{CH}_3\text{CH}_2\text{CHBr}-\text{C}_6\text{H}_4-\text{OH}$
- (c) $\text{CH}_3\text{CHBrCH}_2-\text{C}_6\text{H}_3(\text{Br})-\text{OH}$
- (d) $\text{CH}_3\text{CH}_2\text{CHBr}-\text{C}_6\text{H}_3(\text{Br})-\text{OH}$

5. For which of the following parameters the structural isomers $\text{C}_2\text{H}_5\text{OH}$ and CH_3OCH_3 would be expected to have same values? (Assume ideal behaviour)
 (a) Vapour pressure at the same temperature
 (b) Heat of vaporisation
 (c) Gaseous densities at the same temperature and pressure
 (d) Boiling points
6. Which of the following statements are correct?
 I. Glycerol on reaction with oxalic acid at 110°C (383 K) and followed by hydrolysis gives formic acid and glycerol.
 II. Glycerol on reaction with oxalic acid at 230°C (503 K) and followed by heating gives allyl alcohol.
 III. Glycerol on oxidation with dil. HNO_3 gives a mixture of glyceric and tartaric acid.
 IV. Glycerol on oxidation with conc. HNO_3 gives glyceric acid.
 (a) I and II (b) I and III
 (c) III and IV (d) I, II, III, IV
7. Isopropyl benzene is oxidised in the presence of air to give a compound 'A'. When compound 'A' is treated with dilute mineral acid, the aromatic product formed is
 (a) phenol (b) benzene
 (c) benzaldehyde (d) acetophenone.
8. Intramolecular rearrangement of phenyl acetate to give *o*- and *p*-hydroxyacetophenone in the presence of anhydrous AlCl_3 is known as
 (a) Reimer-Tiemann reaction
 (b) Kolbe's reaction
 (c) Fries rearrangement
 (d) Claisen rearrangement.

9. Identify X in the following sequence :



- (a) $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$ (b) $\text{CH}_3\text{CHOHCH}_3$
 (c) $\text{CH}_3\text{OCH}_2\text{CH}_3$ (d) $\text{CH}_3\text{CH}_2\text{CHO}$

10. What amount of bromine will be required to convert 2 g of phenol into 2, 4, 6-tribromophenol?

- (a) 4.00 g (b) 6.00 g
 (c) 10.08 g (d) 20.44 g


SECTION - II

More than One Options Correct Type

11. Phenol is less acidic than

- (a) acetic acid (b) *p*-methoxyphenol
 (c) *p*-nitrophenol (d) ethanol.

12. Which of the following compounds may undergo Reimer-Tiemann reaction?



13. Which of the following compounds will give a yellow precipitate with iodine and alkali?


- (a) Acetophenone (b) Methyl acetate
 (c) Acetamide (d) 2-Hydroxypropane

SECTION - III

Paragraph Type

Paragraph for Questions 14 and 15


Reimer-Tiemann reaction introduces aldehyde group on to the aromatic ring of phenol, *ortho* to the hydroxyl group. This reaction involves electrophilic aromatic substitution. This is a general method for the synthesis of a substituted salicylaldehydes as depicted below.



14. Which one of the following reagents is used in the above reaction?

- (a) aq. NaOH + CH_3Cl
 (b) aq. NaOH + CH_2Cl_2
 (c) aq. NaOH + CHCl_3
 (d) aq. NaOH + CCl_4


15. The structure of the intermediate (I) is




Paragraph for Questions 16 and 17



Compound (A), $\text{C}_{10}\text{H}_{12}\text{O}_3$ is insoluble in water, dil. HCl and dilute aq. NaHCO_3 . It is however soluble in dilute NaOH. A solution of (A) in dilute NaOH was boiled and the distillate collected in a solution of NaOI, where a yellow precipitate was formed. The alkaline residue in the distillation flask was acidified with dil. H_2SO_4 to precipitate a solid (B), which was boiled, steam distilled and collected. The pure steam distilled product (B) was found to have the formula, $\text{C}_7\text{H}_6\text{O}_3$ and it was dissolved in aq. NaHCO_3 with effervescence.

16. The structural formula of compound (A) is



17. The structural formula of compound (B) is





(d) none of these.

SECTION - IV

Matching List Type

18. Match the List I with List II and select the correct answer using the codes given below the lists :

List I (Compounds)	List II (Methods to distinguish)
(P) Methanol and ethanol	1. Lucas reagent
(Q) Phenol and cyclohexanol	2. Sodium metal
(R) <i>n</i> -Butyl alcohol and <i>tert</i> -butyl alcohol	3. Iodoform test
(S) Ethanol and diethyl ether	4. Ferric chloride

P	Q	R	S
(a) 2	4	1	3
(b) 4	1	2	3
(c) 1	2	3	4
(d) 3	4	1	2

19. Match the List I with List II and select the correct answer using the codes given below the lists :

List I (Reaction)	List II (Product)
(P) Reimer-Tiemann reaction	1. Benzoquinone
(Q) Kolbe's reaction	2. Salicylaldehyde
(R) Williamson synthesis	3. Salicylic acid
(S) Oxidation of phenol	4. Ether

	P	Q	R	S
(a)	3	4	2	1
(b)	3	4	1	2
(c)	4	3	1	2
(d)	2	3	4	1

SECTION - V

Assertion Reason Type

20. **Assertion :** Phenol is more reactive than benzene.
Reason: In case of phenol, the intermediate carbocation is more resonance stabilised.
21. **Assertion :** The ease of dehydration of alcohols follows the order : Primary > Secondary > Tertiary.
Reason: Dehydration proceeds through the formation of oxonium ions.
22. **Assertion :** *tert*-Butyl methyl ether is not prepared by the reaction of *tert*-butyl bromide with sodium methoxide.
Reason : Sodium methoxide is a weak nucleophile.

SECTION - VI


Integer Value Correct Type

23. How many of structurally isomeric pentyl alcohols will give immediate turbidity in Lucas test?
24.
How many moles of HI consumed in above reaction?
25. How many of the following ethers cannot be prepared by Williamson's synthesis?
 $\text{CH}_3\text{OCH}_2\text{CH}_3$, $\text{C}_6\text{H}_5\text{OCH}_3$, $\text{C}_6\text{H}_5\text{OCH}_2\text{CH}_3$, $(\text{C}_6\text{H}_5)_2\text{O}$, $(\text{CH}_3)_3\text{COCH}_3$, $(\text{CH}_3)_3\text{COCH}_2\text{CH}_3$, $(\text{CH}_3)_3\text{COC}(\text{CH}_3)_3$, $(\text{C}_2\text{H}_5)_2\text{O}$, $\text{C}_6\text{H}_5\text{CH}_2\text{OC}_6\text{H}_5$


SOLUTIONS

THE *p*-BLOCK ELEMENTS (GROUP 15 TO 18)

1. (a)
2. (a) : $\text{SO}_2 + 2\text{H}_2\text{S} \longrightarrow 3\text{S} + 2\text{H}_2\text{O}$
3. (c) : Although both hydration energy and bond dissociation energy make F_2 a strong oxidising agent but the effect of hydration energy is more important.
4. (a)
5. (a) : $\text{P}_4 + 3\text{NaOH} + 3\text{H}_2\text{O} \longrightarrow 3\text{NaH}_2\text{PO}_2 + \text{PH}_3 \uparrow$
Caustic Sodium Phosphine
soda hypophosphite
6. (d) : Br^- has been changed to Br_2 . H_2SO_4 is acting as an oxidising agent.
7. (a) : $\text{MnO}_2 + 4\text{HCl} \longrightarrow \text{MnCl}_2 + \text{Cl}_2 + 2\text{H}_2\text{O}$
Greenish
yellow





8. (a): The stability of interhalogens increases as the size difference and polarity of bond increases. Thus, the correct order of stability is $\text{IF}_3 > \text{BrF}_3 > \text{ClF}_3$.






10. (d): The oxidation state of sulphur in its compounds varies from -2 to +6.


- 11.(a,c): If chlorine is passed through dry slaked lime, bleaching powder is produced.




12. (b,c)




14. (c): Pure H_2SO_4 does not act as oxidising agent, because its molecules are associated due to strong hydrogen bonds.



15. (a): Two hydroxyl groups of sulphuric acid can be replaced by chlorine when treated with PCl_5 .




16. (c): In XeO_3 there are total of 4 electron pairs around central atom. Out of which 3 are bonding electron pairs and one is non-bonding electron pair. Thus, it has sp^3 hybridisation and pyramidal shape.




17. (a): All the xenon fluorides are strongly oxidising.

18. (b) 19. (c)



H_3PO_3 is reducing in nature but H_3PO_4 is not a reducing agent as in H_3PO_4 , phosphorus is in +5 oxidation state.


21. (b):



22. (d)

23. (7): The oxidation state of Cl in HClO_4 is +7.


24. (4):



4(P—O—H) bonds are present in pyrophosphoric acid.

25. (3): ClF ($17 + 9 = 26$), OF_2 ($8 + 2 \times 9 = 26$), ClO^- ($17 + 8 + 1 = 26$)

ALCOHOLS, PHENOLS AND ETHERS



2. (b)

3. (a): Phenol does not react with copper at 300°C .




Mad about rehearsing?



₹ 400



₹ 400



₹ 400


Tune. Fine tune. Reach the peak of your readiness for JEE with MTG's 40+16 Years Chapterwise Solutions. It is undoubtedly the most comprehensive 'real' question bank, complete with detailed solutions by experts.

Studies have shown that successful JEE aspirants begin by familiarising themselves with the problems that have appeared in past JEEs as early as 2 years in advance. Making it one of the key ingredients for their success. How about you then? Get 40+16 Years Chapterwise Solutions to start your rehearsals early. Visit www.mtg.in to order online.




Available at all leading book shops throughout the country.
For more information or for help in placing your order:
Call 0124-6601200 or email: info@mtg.in

Visit
www.mtg.in
for latest offers
and to buy
online!




5. (c): As $\text{C}_2\text{H}_5\text{OH}$ and CH_3OCH_3 are isomers, they have the same molecular mass and have same values of gaseous densities at the same temperature and pressure.

6. (d)



8. (c)

9. (b): Since $\text{C}_3\text{H}_6\text{O}$ on treatment with $\text{I}_2 + \text{NaOH}$ gives CHI_3 , therefore, $\text{C}_3\text{H}_6\text{O}$ must be a methyl ketone, i.e. CH_3COCH_3 . If this is so then, X must be $\text{CH}_3\text{CHOHCH}_3$.




94 grams of phenol requires bromine = 474 g
 \therefore 2 grams of phenol would require amount of bromine = $\frac{474}{94} \times 2 = 10.08$ g

11. (a,c)

12. (a,b)

13. (a,d): Compounds which contain $\text{CH}_3\text{CO}-$ or $\text{CH}_3\text{CHOH}-$ group attached to C or H on treatment with I_2 and alkali give yellow ppt. of iodoform.





Thus, acetophenone and 2-hydroxypropane give iodoform but acetamide and methyl acetate which although contain $\text{CH}_3\text{CO}-$ group but that is not attached to C or H and hence, do not give iodoform.

14. (c): For introduction of $-\text{CHO}$ group, during Reimer-Tiemann reaction, a mixture of aq. $\text{NaOH} + \text{CHCl}_3$ is used.

15. (b)

16. (b):




18. (d)

19. (d)

20. (a)

21. (d): The ease of dehydration of alcohols can be explained on the basis of stability of the intermediate carbocation. Greater the stability of the carbocation formed, greater will be the rate of reaction. The order of stability of carbocation formed is




This is due to the electron releasing (+I) effect of the alkyl group. Therefore, the ease of dehydration of alcohols follows the order:

Tertiary > secondary > primary alcohols

22. (c): Sodium ethoxide is a strong nucleophile.

23. (1): Only $\text{H}_3\text{C}-\overset{\text{CH}_3}{\underset{\text{OH}}{\text{C}}}-\text{CH}_2\text{CH}_3$, which is a tertiary alcohol, gives immediate turbidity with Lucas reagent.

24. (2):



25. (2): Williamson's synthesis requires that the alkyl halide should be 1° and alkoxide ion may be 1° , 2° or 3° . Thus, two ethers which cannot be prepared by Williamson's synthesis are $(\text{C}_6\text{H}_5)_2\text{O}$, $(\text{CH}_3)_3\text{COC}(\text{CH}_3)_3$ ◆◆

ACE

YOUR WAY CBSE XII



Series 4

CHAPTERWISE PRACTICE PAPER : THE *d*- AND *f*-BLOCK ELEMENTS | COORDINATION COMPOUNDS

Time Allowed : 3 hours

Maximum Marks : 70

GENERAL INSTRUCTIONS

- (i) All questions are compulsory.
- (ii) Q. no. 1 to 5 are very short answer questions and carry 1 mark each.
- (iii) Q. no. 6 to 10 are short answer questions and carry 2 marks each.
- (iv) Q. no. 11 to 22 are also short answer questions and carry 3 marks each.
- (v) Q. no. 23 is a value based question and carry 4 marks.
- (vi) Q. no. 24 to 26 are long answer questions and carry 5 marks each.
- (vii) Use log tables if necessary, use of calculators is not allowed.

1. Why do Zr and Hf exhibit similar properties?
 2. Why is Ca-EDTA used for the treatment of lead poisoning?
 3. What properties of tungsten make it suitable for use as filaments in light bulbs?
 4. What is the coordination number of Fe in $[\text{Fe}(\text{EDTA})]$?
 5. State the kind of isomerism possible for the following :
 - (i) $[\text{Cr}(\text{en})_3]^{3+}$
 - (ii) $[\text{Cr}(\text{NH}_3)_4(\text{Cl})(\text{Br})]\text{Br}$
 6. What will be the correct order of absorption of wavelength of light in the visible region for the complexes $[\text{Co}(\text{NH}_3)_6]^{3+}$, $[\text{Co}(\text{CN})_6]^{3-}$ and $[\text{Co}(\text{H}_2\text{O})_6]^{3+}$?
 7. Write the formula of the oxo-metal anions of the first series of the transition metals in which the metal exhibits the oxidation state equal to its group number.
 8. Specify the oxidation numbers of the metals in the following coordination entities :
 - (a) $[\text{Co}(\text{H}_2\text{O})(\text{CN})(\text{en})_2]^{2+}$
 - (b) $[\text{PtCl}_4]^{2-}$
 - (c) $[\text{Cr}(\text{NH}_3)_3\text{Cl}_3]$
 - (d) $[\text{CoBr}_2(\text{en})_2]^+$
 9. Write the electronic configurations of the elements with the atomic numbers 61, 91, 101 and 109.
- OR**
- Write the chemical equations for the reactions of $\text{Cr}_2\text{O}_7^{2-}$ with excess of (a) Fe^{2+} and (b) SO_3^{2-} .
10. Explain the crystal field splitting in square planar complexes.
 11. How would you account for the following?
 - (i) The atomic radii of the third series of transition elements are virtually the same as those of the corresponding members of the second series.
 - (ii) The E° value for $\text{Mn}^{3+}/\text{Mn}^{2+}$ couple is much more positive than that for $\text{Cr}^{3+}/\text{Cr}^{2+}$ couple or $\text{Fe}^{3+}/\text{Fe}^{2+}$ couple.
 - (iii) The highest oxidation state of a metal is exhibited in its oxide or fluoride.
 12. Calculate the crystal field stabilisation energy of a d^8 ion in a square planar field for both strong and weak field cases.

13. (i) Why are Sm^{2+} , Eu^{2+} and Yb^{2+} good reducing agents?
 (ii) Can lanthanum ($Z = 57$) exhibit +4 oxidation state?
 (iii) Why are +3 oxidation state of gadolinium ($Z = 64$) and lutetium ($Z = 71$) especially stable?
14. Name the following compounds :
 (a) $[\text{Co}(\text{en})_2(\text{CN})_2]\text{ClO}_3$
 (b) $\text{K}_4[\text{Co}(\text{CN})_6]$
 (c) $[\text{Ni}(\text{NH}_3)_6]_3[\text{Co}(\text{NO}_2)_6]_2$
15. Name the members of the lanthanoid series which exhibit + 4 oxidation states and those which exhibit +2 oxidation states. Try to correlate this type of behaviour with the electronic configuration of these elements.
16. Write the correct formulae for the following coordination compounds :
 (a) $\text{CrCl}_3 \cdot 6\text{H}_2\text{O}$ (violet, with 3 chloride ions/unit formula)
 (b) $\text{CrCl}_3 \cdot 6\text{H}_2\text{O}$ (light green colour, with 2 chloride ions/unit formula)
 (c) $\text{CrCl}_3 \cdot 6\text{H}_2\text{O}$ (dark green colour, with 1 chloride ion/unit formula)
- OR**
- For the square coplanar complex $[\text{Pt}(\text{NH}_3)(\text{NH}_2\text{OH})\text{py}(\text{NO}_2)]^+$, how many geometrical isomers are possible? Draw the structures of these isomers.
17. What are interstitial compounds? Explain them with reference to transition metals and mention their two important properties.
18. What can be inferred from the magnetic moment values of the following complex species?
- | Example | Magnetic Moment (B.M.) |
|---|------------------------|
| (i) $\text{K}_4[\text{Mn}(\text{CN})_6]$ | 1.7 |
| (ii) $[\text{Fe}(\text{H}_2\text{O})_6]^{2+}$ | 4.9 |
| (iii) $\text{K}_2[\text{MnCl}_4]$ | 5.9 |
19. Decide giving reasons which one will exhibit the property indicated in the given pair.
 (i) Sc^{3+} or Cr^{3+} : Paramagnetism
 (ii) V or Mn : More number of oxidation states
 (iii) V^{4+} or V^{5+} : Diamagnetism
20. Discuss briefly giving an example in each case the role of coordination compounds in
 (i) biological systems (ii) medicinal chemistry
 (iii) analytical chemistry.
21. Among lanthanoids, Ln(III) compounds are predominant. However, occasionally in solutions or in solid compounds, +2 and +4 ions are also obtained.
22. Draw geometrical isomers of the following :
 (a) $[\text{CoCl}_2(\text{en})_2]^+$ (b) $[\text{Co}(\text{NH}_3)_2\text{Cl}_2(\text{en})]^+$
 (c) $[\text{Co}(\text{NH}_3)\text{Cl}(\text{en})_2]^{2+}$
23. Raman after passing M.Sc. in Chemistry joined a textile factory as a junior scientist. He came to know that factory is using chlorine based bleaching agents for bleaching of textiles. Raman suggested the chief production manager of the factory to use potassium permanganate instead of chlorine for bleaching purpose.
 Now answer the following questions :
 (i) Which property of KMnO_4 makes it a bleaching agent?
 (ii) Write reduction half reaction of KMnO_4 in acidic medium.
 (iii) Why is KMnO_4 better bleaching agent than chlorine?
 (iv) What value is associated with the suggestion of Raman?
24. (a) Give the formula of each of the following coordination entities :
 (i) Co^{3+} ion is bound to one Cl^- , one NH_3 molecule and two bidentate ethylene diamine (*en*) molecules.
 (ii) Ni^{2+} ion is bound to two water molecules and two oxalate ions.
- Write the name and magnetic behaviour of each of the above coordination entities.
 (b) Identify low spin and high spin complex out of these two.
- OR**
- Indicate the oxidation state, electronic configuration, coordination number and magnetic moment for each of the following complexes.
 (i) $\text{K}[\text{Cr}(\text{H}_2\text{O})_2(\text{C}_2\text{O}_4)_2] \cdot 3\text{H}_2\text{O}$
 (ii) $[\text{Co}(\text{NH}_3)_5\text{Cl}]\text{Cl}_2$ (iii) $[\text{CrCl}_3(\text{py})_3]$
 (iv) $\text{Cs}[\text{FeCl}_4]$ (v) $\text{K}_4[\text{Mn}(\text{CN})_6]$
25. (i) A black mineral (A) on treatment with dilute sodium cyanide solution in the presence of air gives a clear solution containing (B) and (C).
 (ii) The solution of (B) on reaction with zinc gives precipitate of metal (D).
 (iii) (D) dissolved in dil. HNO_3 and the resulting solution gives a white precipitate of (E) with dil. HCl.

- (iv) (E) on fusion with sodium carbonate gives (D).
 (v) (E) dissolved in aqueous solution of ammonia giving a colourless solution of (F).

Identify A to F and give chemical equations involved.

OR

- (a) First ionisation energy of copper is higher than those of alkali metals, while second and third ionisation energies are lower. Explain.
 (b) A mixed oxide of iron and chromium, $\text{FeO} \cdot \text{Cr}_2\text{O}_3$ is fused with sodium carbonate in the presence of air to form a yellow compound (A). On acidification, the compound (A) forms an orange coloured compound (B) which is a strong oxidising agent. Identify
 (i) the compounds (A) and (B)
 (ii) write balanced chemical equations for each step.
26. (a) Arrange the following complexes in the increasing order of conductivity of their solutions : $[\text{Co}(\text{NH}_3)_3\text{Cl}_3]$, $[\text{Co}(\text{NH}_3)_4\text{Cl}_2]\text{Cl}$, $[\text{Co}(\text{NH}_3)_6]\text{Cl}_3$, $[\text{Co}(\text{NH}_3)_5\text{Cl}]\text{Cl}_2$.
 (b) Why do compounds having similar geometry have different magnetic moment?
 (c) $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ is blue in colour while CuSO_4 is colourless. Why?


OR

- (a) Why are different colours observed in octahedral and tetrahedral complexes for the same metal and same ligands?
 (b) What is the relationship between observed colour of the complex and the wavelength of light absorbed by the complex?
 (c) $\text{CoSO}_4\text{Cl} \cdot 5\text{NH}_3$ exists in two isomeric forms 'A' and 'B'. Isomer 'A' reacts with AgNO_3 to give white precipitate, but does not react with BaCl_2 . Isomer 'B' gives white precipitate with BaCl_2 but does not react with AgNO_3 . Answer the following questions :
 (i) Identify 'A' and 'B' and write their structural formulae.
 (ii) Name the type of isomerism involved.
 (iii) Give the IUPAC name of 'A' and 'B'.


SOLUTIONS

1. Due to lanthanoid contraction, Zr and Hf have nearly similar size *i.e.*, Zr (160 pm), Hf (159 pm), thus, they exhibit similar properties.
 2. The complex of calcium with EDTA is used for the treatment of lead poisoning. Lead readily replaces calcium in the complex and lead-EDTA complex thus formed is finally eliminated from the body through urine.
 3. High melting point (3410°C), moderate electrical resistance, low volatility, makes tungsten an important component in light bulbs.
 4. EDTA is a hexadentate ligand, thus coordination number of Fe is 6 in $[\text{Fe}(\text{EDTA})]$.
 5. (i) Optical isomerism
 (ii) Geometrical isomerism
 6. As strength of ligand increases crystal field splitting energy (CFSE) increases.
 Order of strength of ligands :
 $\text{I}^- < \text{Br}^- < \text{Cl}^- < \text{NO}_3^- < \text{F}^- < \text{OH}^- < \text{ox} < \text{H}_2\text{O} < \text{py} = \text{NH}_3 < \text{en} < \text{dipy} < \text{o-phen} < \text{NO}_2^- < \text{CN}^- < \text{CO}$.
- Now, $\Delta E = \frac{hc}{\lambda}$.
- So, as CFSE increases, ΔE increases and λ decreases. Thus, the correct order of absorption of wavelength of light in the visible region is :
 $[\text{Co}(\text{H}_2\text{O})_6]^{3+} > [\text{Co}(\text{NH}_3)_6]^{3+} > [\text{Co}(\text{CN})_6]^{3-}$
7. ScO_2^- ; Group number = Oxidation state of Sc = 3.
 TiO_3^{2-} ; Group number = Oxidation state of Ti = 4
 VO_3^- ; Group number = Oxidation state of V = 5
 $\text{Cr}_2\text{O}_7^{2-}$; Group number = Oxidation state of Cr = 6
 CrO_4^{2-} ; Group number = Oxidation state of Cr = 6
 MnO_4^- ; Group number = Oxidation state of Mn = 7
8. (a) $x + (-1) = +2$ or $x = +3$
 (b) $x + 4(-1) = -2$ or $x = +2$
 (c) $x + 3(-1) = 0$ or $x = +3$
 (d) $x + 2(-1) = +1$ or $x = +3$
9. Atomic number (61) = $_{54}[\text{Xe}] 4f^5 6s^2$
 Atomic number (91) = $_{86}[\text{Rn}] 5f^2 6d^1 7s^2$
 Atomic number (101) = $_{86}[\text{Rn}] 5f^{13} 7s^2$
 Atomic number (109) = $_{86}[\text{Rn}] 5f^{14} 6d^7 7s^2$
- OR
- (a) $6\text{Fe}^{2+} + \text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ \longrightarrow 6\text{Fe}^{3+} + 2\text{Cr}^{3+} + 7\text{H}_2\text{O}$
 (b) $\text{Cr}_2\text{O}_7^{2-} + 3\text{SO}_3^{2-} + 8\text{H}^+ \longrightarrow 3\text{SO}_4^{2-} + 2\text{Cr}^{3+} + 4\text{H}_2\text{O}$
10. The square planar arrangement can be derived from octahedral by removing two ligands from z-axis. On removing ligands from z-axis, d_{z^2} , d_{xz} and d_{yz} orbitals become more stable. As the lobes of $d_{x^2-y^2}$ point towards the ligands, the orbital has highest energy and the lobes of d_{xy} orbital lie

between the ligands, it is thus next highest orbital in energy. The lobes of d_{xz} and d_{yz} are least affected as they lie out of the plane of the complex. The d_z^2 orbital is somewhat affected by electrostatic field of the ligands.



11. (i) The atomic radii of the third series of transition elements are virtually the same as those of the corresponding members of the second series due to lanthanoid contraction.
- (ii) $Mn^{3+}(3d^4)$ is less stable than $Mn^{2+}(3d^5)$ which has stable half-filled configuration. Fe^{3+} is more stable than Fe^{2+} due to half-filled configuration. Cr^{3+} has stable t_{2g}^3 configuration. So, E° value for the Mn^{3+}/Mn^{2+} couple is much more positive.
- (iii) The highest oxidation state of a metal is exhibited in its oxide or fluoride. Due to the high electronegativity and small size of oxygen and fluorine.
12. The terms "strong field" for square planar complexes means that the splitting of the two highest orbitals is greater than the pairing energy. Thus, the strong and weak field configurations are



$$\text{CFSE (for strong field)} = 4(-0.51\Delta_o) + 2(-0.43\Delta_o) + 2(0.23\Delta_o) + P = -2.44\Delta_o + P$$

$$\text{CFSE (for weak field)} = 4(-0.51\Delta_o) + 2(-0.43\Delta_o) + 1(+0.23\Delta_o) + 1(+1.23\Delta_o) + P = -1.44\Delta_o + P$$

13. (i) The most stable oxidation state of lanthanides is +3. Hence, ions in +2 state tend to change to +3 oxidation state by loss of electron and act as reducing agents.
- (ii) La^{3+} has a stable configuration of an inert gas $[Xe]5d^06s^0$. To obtain +4 oxidation state, the stable configuration is to be disturbed which is not possible under ordinary conditions and hence, La^{4+} does not exist.
- (iii) This is because gadolinium in +3 oxidation state has half filled $4f$ -subshell ($4f^7$) and lutetium in +3 state has completely filled $4f$ -subshell ($4f^{14}$) which are very stable configurations.

14. (a) Dicyanobis(ethylenediamine)cobalt(III) chlorate
(b) Potassium hexacyanocobaltate(II)
(c) Hexaamminenickel(II) hexanitrocobaltate(III)
15. +4 oxidation state : Cerium (Ce), Praseodymium (Pr), Terbium (Tb), Dysprosium (Dy)
+2 oxidation state : Neodymium (Nd), Promethium (Pm), Samarium (Sm), Europium (Eu), Thulium (Tm) and Ytterbium (Yb)
+4 and +2 oxidation states of lanthanoid elements arise mainly from the extra stability of empty, half-filled and fully-filled f -subshells. For example, Ce(IV) is $4f^0$ and has noble gas configuration. Similarly, Tb (IV) and Eu (II) have $4f^7$ configuration. Yb(II) has $4f^{14}$ configuration.

16. The given three coordination compounds are examples of special form of ionisation, called hydrate isomerism. Water can exist as a part of coordination entity or a part outside the coordination entity.
- (a) $[Cr(H_2O)_6]Cl_3$: all three Cl^- ions are outside the coordination entity.
- (b) $[CrCl(H_2O)_5]Cl_2 \cdot H_2O$: one Cl^- ion is part of coordination entity and two are outside of it.
- (c) $[CrCl_2(H_2O)_4]Cl \cdot 2H_2O$: two Cl^- ions are part of coordination entity and one is present outside of it.

The Cl^- ions outside the coordination entity are ionisable and forms white precipitate with $AgNO_3$ while Cl^- ions which are part of coordination entity are not ionisable.

OR

Three geometric isomers are possible for the given coordination compound. Any other possible configuration would merely be a rotation or reflection of one of these.

NEET के ENTRANCE EXAM में हिंदी माध्यम विद्यार्थियों के लिए सफलता मंत्र

NCERT TEXTBOOK पर BASED भौतिकी, रसायन
और जीव विज्ञान की OBJECTIVE पुस्तकें

NEET पाठ्यक्रम पर आधारित और हमारे SUBJECT EXPERTS द्वारा निर्मित भौतिकी, रसायन और जीव विज्ञान के CHAPTERWISE OBJECTIVE TYPE प्रश्नों का अभ्यास कर इन तीनों विषयों में महारत हासिल कर NEET परीक्षाओं में सफलता प्राप्त करें और विजयी बनें।



₹ 600



₹ 550



₹ 750

ये तीनों पुस्तकें ही पढ़ना क्यों जरूरी हैं?

- प्रश्नों को शीघ्रता और सरलता से हल करने के लिए CHAPTERWISE SYNOPSIS (मुख्य बिंदु)
- छात्रों की प्रगति जाँचने के लिए TOPICWISE OBJECTIVE प्रश्न
- NCERT EXEMPLAR (प्रश्न प्रदर्शिका) के प्रश्न विस्तृत उत्तरों के साथ
- सभी राष्ट्रीय एवं राज्य स्तरीय प्रतियोगिताओं के लिए अत्यंत उपयोगी, सरल एवं महत्वपूर्ण पुस्तकें
- AIIMS | JEE की तैयारी हेतु अभिकथन एवं तर्क प्रारूप प्रश्न
- स्वमूल्यांकन हेतु पाँच अभ्यास प्रश्न पत्र



NEET की सबसे महत्वपूर्ण CHAPTERWISE QUESTION BANK SERIES की पुस्तकें



₹ 200



₹ 200




₹ 275

- 100% NEET पाठ्यक्रम पर आधारित
- 17 वर्षों (2017-2001) के NEET/AIPMT के CHAPTERWISE प्रश्नों के उत्तर सरल व्याख्या सहित



Visit www.MTG.in to buy online. Or visit a leading bookseller near you.
For more information, e-mail info@mtg.in or call 1800 300 23355 (toll-free) today.



17. Small atoms like B, C, N, etc., can occupy positions in the voids among the atoms of transition metals. Such combinations are termed as interstitial compounds. They are usually non-stoichiometric materials having formulae like $\text{VH}_{0.56}$, $\text{TiH}_{1.7}$, TiC , etc. Steel is an interstitial compound in which voids among iron atoms are occupied by carbon atoms. Two important properties :

- (i) Transition metals become more harder and rigid.
- (ii) Melting points of interstitial compounds are higher than parent metals.

18. Calculate the magnetic moment by applying formula; $\mu = \sqrt{n(n+2)}$

- (i) The magnetic moment of 1.7 B.M. corresponds to $n = 1$. Thus, in $\text{K}_4[\text{Mn}(\text{CN})_6]$ there is only one unpaired electron in $3d$ -subshell. Thus, the distribution of five $3d$ -electrons in $\text{Mn}(\text{II})$ is $(t_{2g})^5$.
- (ii) The magnetic moment of 4.9 B.M. indicates that there are four unpaired electrons in the complex. Thus, the six $3d$ -electrons in $\text{Fe}(\text{II})$ are distributed as $(t_{2g})^4 (e_g)^2$.
- (iii) The magnetic moment of 5.9 B.M. indicates that there are five unpaired electrons in the complex. Hence, the five $3d$ -electrons in $\text{Mn}(\text{II})$ are distributed as $(t_{2g})^3 (e_g)^2$ in the given complex.

19. (i) $\text{Sc} : [\text{Ar}]3d^1 4s^2$, $\text{Sc}^{3+} : [\text{Ar}]$;

It has no unpaired electron.

$\text{Cr} : [\text{Ar}]3d^5 4s^1$, $\text{Cr}^{3+} : [\text{Ar}]3d^3$;

It has three unpaired electrons. Hence, Cr^{3+} is paramagnetic.

- (ii) $\text{V} : [\text{Ar}]3d^3 4s^2$, $\text{Mn} : [\text{Ar}]3d^5 4s^2$; Mn exhibits more number of oxidation states.

The expected oxidation states in V are +2, +3, +4, +5.

The expected oxidation states in Mn are +2, +3, +4, +5, +6, +7.

- (iii) $\text{V}^{4+} : [\text{Ar}] 3d^1$; One unpaired electron
 $\text{V}^{5+} : [\text{Ar}]$; No unpaired electron
 Hence, V^{5+} is diamagnetic in nature.

20. (i) Chlorophyll is a coordination compound of magnesium. Haemoglobin is a coordination compound of iron. Vitamin B_{12} , cyanocobalamin, the anti-pernicious anaemia factor, is a coordination compound of cobalt.


- (ii) There is growing interest in the use of chelate therapy in medicinal chemistry. Excess of copper and iron are removed by the chelating ligands *D*-penicillamine and desferrioxime *B* via the formation of coordination compounds. EDTA is used in the treatment of lead poisoning. Some coordination compounds of platinum effectively inhibit the growth of tumours *e.g.* *cis*-platin.

- (iii) Coordination compounds find use in many qualitative and quantitative chemical analysis. The familiar colour reactions given by metal ions with a number of ligands (especially chelating ligands) as a result of formation of coordination entities, form the basis for their detection and estimation by classical and instrumental methods of analysis. Examples of such reagents include EDTA, DMG (dimethylglyoxime), α -nitroso- β -naphthol, etc.


21. All lanthanoids exhibit a common stable oxidation state of +3. In addition, some lanthanoids show +2 and +4 oxidation states also in solution or in solid compounds. These are shown by those elements which by doing so attain the stable f^0 (empty f -subshell), f^7 (half filled f -subshell) and f^{14} (fully filled f -subshell) configurations.


For example,

- (i) Ce and Tb exhibit +4 oxidation state. Cerium (Ce) and terbium (Tb) attain f^0 and f^7 configurations respectively when they get +4 oxidation states, as shown below :




- (ii) Eu and Yb exhibit +2 oxidation state. Europium and ytterbium get f^7 and f^{14} configurations in +2 oxidation states as shown below :







23. (i) Oxidising property of KMnO_4 makes it a bleaching agent.
 (ii) Reduction half reaction of KMnO_4 in acidic medium
 $\text{MnO}_4^- + 8\text{H}^+ + 5\text{e}^- \longrightarrow \text{Mn}^{2+} + 4\text{H}_2\text{O}$
 $E^\circ = +1.52 \text{ V}$
 (iii) Chlorine may form some harmful compounds during bleaching of textiles. These compounds may cause water pollution whereas the bleaching by KMnO_4 is harmless.
 (iv) Self awareness and concern towards environment.

24. (a) (i) $[\text{Co}(\text{NH}_3)(\text{Cl})(\text{en})_2]^{2+}$
 Name : Amminechloridobis(ethane-1, 2-diamine) cobalt(III) ion



Since there are no unpaired electrons, complex is diamagnetic.

(ii) $[\text{Ni}(\text{H}_2\text{O})_2(\text{C}_2\text{O}_4)_2]^{2-}$
 Name : Diaquadioxalatonickelate(II) ion



The complex has two unpaired electrons and therefore, it will be paramagnetic.


(b) $[\text{Co}(\text{NH}_3)(\text{Cl})(\text{en})_2]^{2+}$ is low spin complex and $[\text{Ni}(\text{H}_2\text{O})_2(\text{C}_2\text{O}_4)_2]^{2-}$ is high spin complex.

OR

(i) $\text{K}[\text{Cr}(\text{H}_2\text{O})_2(\text{C}_2\text{O}_4)_2] \cdot 3\text{H}_2\text{O}$

Oxidation state = +3, coordination number = 6

Electronic configuration : t_{2g}^3




$$\mu = \sqrt{n(n+2)} = \sqrt{3 \times 5} = \sqrt{15} = 3.87 \text{ B.M.}$$

(ii) $[\text{Co}(\text{NH}_3)_5\text{Cl}]\text{Cl}_2$

Oxidation state of Co = +3, coordination number = 6

Electronic configuration : t_{2g}^6




$\mu = 0 \text{ B.M.}$

(iii) $[\text{CrCl}_3(\text{py})_3]$

Oxidation state = +3, coordination number = 6

Electronic configuration : t_{2g}^3




$$\mu = \sqrt{n(n+2)} = \sqrt{3 \times 5} = \sqrt{15} = 3.87 \text{ B.M.}$$

(iv) $\text{Cs}[\text{FeCl}_4]$

Oxidation state = +3, coordination number = 4

Electronic configuration : $t_{2g}^3 e_g^2$




$$\mu = \sqrt{n(n+2)} = \sqrt{5 \times 7} = \sqrt{35} = 5.92 \text{ B.M.}$$

(v) $\text{K}_4[\text{Mn}(\text{CN})_6]$

Oxidation state = +2, coordination number = 6

Electronic configuration : t_{2g}^5



$$\mu = \sqrt{n(n+2)} = \sqrt{1 \times 3} = 1.732 \text{ B.M.}$$

25. (i) $\text{Ag}_2\text{S} + 4\text{NaCN} + 2\text{O}_2 \longrightarrow 2\text{Na}[\text{Ag}(\text{CN})_2] + \text{Na}_2\text{SO}_4$
 (A) (B) (C)
- (ii) $2\text{Na}[\text{Ag}(\text{CN})_2] + \text{Zn} \longrightarrow \text{Na}_2[\text{Zn}(\text{CN})_4] + 2\text{Ag}\downarrow$
 (B) (D)
- (iii) $2\text{Ag} + 4\text{HNO}_3 \longrightarrow 2\text{AgNO}_3 + 2\text{NO}_2 + 2\text{H}_2\text{O}$
 (D)
 $\text{AgNO}_3 + \text{HCl} \longrightarrow \text{AgCl}\downarrow + \text{HNO}_3$
 (E)
- (iv) $2\text{AgCl} + \text{Na}_2\text{CO}_3 \longrightarrow 2\text{Ag} + 2\text{NaCl} + \text{CO}_2 + 1/2\text{O}_2$
 (E) (D)
- (v) $\text{AgCl} + 2\text{NH}_4\text{OH} \longrightarrow [\text{Ag}(\text{NH}_3)_2]\text{Cl} + 2\text{H}_2\text{O}$
 (E) (F)
- Thus, A = Ag_2S ; B = $\text{Na}[\text{Ag}(\text{CN})_2]$
 C = Na_2SO_4 ; D = Ag; E = AgCl; F = $[\text{Ag}(\text{NH}_3)_2]\text{Cl}$


OR

- (a) Copper has small size compared to alkali metals in their period. Its electronic configuration is $[\text{Ar}]3d^{10}4s^1$. As filled 3d-subshell is less effective in shielding than s- and p-subshells, so s-electron in copper is more tightly held by the nucleus than that in alkali metals. Hence, its first ionisation energy is higher than for alkali metals. However, second and third ionisation energy values of copper are lower as compared to those of alkali metals due to removal of electrons from diffused d-orbitals.
- (b) $4\text{FeO} \cdot \text{Cr}_2\text{O}_3 + 8\text{Na}_2\text{CO}_3 + 7\text{O}_2 \longrightarrow 8\text{Na}_2\text{CrO}_4 + 2\text{Fe}_2\text{O}_3 + 8\text{CO}_2$
 (A)
 Sodium chromate
 (Yellow)
- $2\text{Na}_2\text{CrO}_4 + \text{H}_2\text{SO}_4 \longrightarrow \text{Na}_2\text{Cr}_2\text{O}_7 + \text{Na}_2\text{SO}_4 + \text{H}_2\text{O}$
 (A) (B)
 (Yellow) (Orange)

The compound (B) is oxidising agent.

26. (a) These complexes can ionise in solution as :
 $[\text{Co}(\text{NH}_3)_3\text{Cl}_3]$ - does not ionise.
 $[\text{Co}(\text{NH}_3)_4\text{Cl}_2]\text{Cl} \longrightarrow [\text{Co}(\text{NH}_3)_4\text{Cl}_2]^+ + \text{Cl}^-$
 (2 ions)
 $[\text{Co}(\text{NH}_3)_6]\text{Cl}_3 \longrightarrow [\text{Co}(\text{NH}_3)_6]^{3+} + 3\text{Cl}^-$ (4 ions)
 $[\text{Co}(\text{NH}_3)_5\text{Cl}]\text{Cl}_2 \longrightarrow [\text{Co}(\text{NH}_3)_5\text{Cl}]^{2+} + 2\text{Cl}^-$
 (3 ions)

As the number of ions in solution increases, their conductivity also increases. Therefore, conductivity follows the order :




- (b) The compounds having similar geometry may have different number of unpaired electrons due to the presence of weak and strong field ligands in complexes. If CFSE is high, the complex will show low value of magnetic moment and if CFSE is low, the complex will show high value of magnetic moment. For example, the $[\text{CoF}_6]^{3-}$ is paramagnetic but $[\text{Co}(\text{NH}_3)_6]^{3+}$ is diamagnetic.
- (c) In $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, water acts as ligand. As a result it causes crystal field splitting. Hence, d-d transition is possible in $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ and shows colour. In the anhydrous CuSO_4 due to the absence of water (ligand), crystal field splitting is not possible and hence, no d-d transition occurs and shows no colour.

OR

- (a) Lower wavelength is absorbed in octahedral complex than tetrahedral complex for the same metal and ligand. In tetrahedral coordination entity, formation of the d-orbital splitting is inverted and is smaller as compared to the octahedral field splitting. Thus, the energy of tetrahedral complex is smaller than compared to octahedral complex. $\Delta_t = (4/9)\Delta_o$
- (b) When white light falls on the complex, some part of it is absorbed. Higher the crystal field splitting, lower will be the wavelength absorbed by the complex. The observed colour of complex is the colour generated from the wavelength left over.
- (c) $\text{CoSO}_4 \cdot 5\text{NH}_3$:
- (i) Isomer A reacts with AgNO_3 but not with BaCl_2 , it shows it has Cl^- ion outside the coordination sphere.
 Hence, A = $[\text{Co}(\text{NH}_3)_5\text{SO}_4]\text{Cl}$
 Isomer B reacts with BaCl_2 but not with AgNO_3 , it shows it has SO_4^{2-} outside the coordination sphere.
 Hence, B = $[\text{Co}(\text{NH}_3)_5\text{Cl}]\text{SO}_4$
- (ii) Ionisation isomerism
- (iii) A = Pentaamminesulphatocobalt(III) chloride and
 B = Pentaamminechlorocobalt(III) sulphate



Concerned about your performance in **Class XII** Boards?



Well, fear no more, help is at hand.....

To excel, studying in right direction is more important than studying hard. Which is why we created the Excel Series. These books – for Physics, Chemistry, Biology & Mathematics – have been put together totally keeping in mind the prescribed syllabus and the pattern of CBSE's Board examinations, so that students prepare and practice with just the right study material to excel in board exams.

Did you know nearly all questions in CBSE's 2017 Board Examination were a part of our Excel books? That too fully solved !

HIGHLIGHTS:

- Comprehensive theory strictly based on NCERT, complemented with illustrations, activities and solutions of NCERT questions
- Practice questions & Model Test Papers for Board Exams
- Value based questions
- Previous years' CBSE Board Examination Papers (Solved)
- CBSE Board Papers 2017 Included



Scan now with your smartphone or tablet*

Visit
www.mtg.in
for latest offers
and to buy
online!



Available at all leading book shops throughout the country.
For more information or for help in placing your order:
Call 0124-6601200 or email: info@mtg.in
*Application to read QR codes required

MPP-5 MONTHLY Practice Problems

Class XII

This specially designed column enables students to self analyse their extent of understanding of specified chapters. Give yourself four marks for correct answer and deduct one mark for wrong answer. Self check table given at the end will help you to check your readiness.

The *d*- and *f*- Block Elements | Coordination Compounds

Total Marks : 120

Time Taken : 60 Min.

NEET / AIIMS

Only One Option Correct Type

- The basic character of the transition metal monoxides follow the order
 - CrO > VO > FeO > TiO
 - TiO > VO > CrO > FeO
 - TiO > FeO > VO > CrO
 - VO > CrO > TiO > FeO
- Which of the following statements is correct?
 - Geometrical isomerism is not observed in complexes of coordination number four having tetrahedral geometry.
 - Square planar complexes generally do not show geometrical isomerism.
 - The square planar complexes of general formulae Ma_3b or Mab_3 exhibit *cis-trans* isomerism.
 - The platinum glycinate complex, $[Pt(gly)_2]$ does not show geometrical isomerism.
- There are three unpaired electrons in $[Co(H_2O)_6]^{2+}$ and calculated value of magnetic moment is 3.87 B.M which is quite different from the experimental value of 4.40 B.M. This is because of
 - increase in number of unpaired electrons
 - some contribution of the orbital motion of the electron to the magnetic moment
 - change in orbital spin of the electron
 - d-d* transition.
- In which of the following metal complexes, central metal atom have zero valency?
 - $[Cu(NH_3)_4]SO_4$
 - $K_3[Fe(CN)_6]$
 - $[Ni(CO)_4]$
 - $[Pt(NH_3)_2Cl_2]$
- If zinc pieces are added to $FeCl_3$ solution, its colour changes from yellow to light green. It is because
 - $FeCl_3$ solution is acidic due to hydrolysis and zinc added produces nascent hydrogen causing reduction of $FeCl_3$ to $FeCl_2$
 - $FeCl_3$ solution is basic and zinc added produces H_2 which reduces $FeCl_3$ to $FeCl_2$
 - zinc oxidises $FeCl_3$ to $FeCl_2$
 - none of the above is correct.
- The two compounds pentaamminesulphatocobalt(III) bromide and pentaamminesulphatocobalt(III) chloride represent
 - linkage isomerism
 - ionisation isomerism
 - coordination isomerism
 - none of these.
- Which compound is coloured due to charge transfer spectra?
 - $K_2Cr_2O_7$
 - $KMnO_4$
 - CrO_3
 - All of these
- Crystal field stabilisation energies of octahedral and tetrahedral complexes having high spin d^6 configuration respectively are
 - $0.6 \Delta_o, 0.6 \Delta_t$
 - $0.4 \Delta_o, 0.4 \Delta_t$
 - $0.4 \Delta_o, 0.6 \Delta_t$
 - $0.6 \Delta_o, 0.4 \Delta_t$
- Pick out the correct statements from the following :
 - Cobalt(III) is more stable in octahedral complexes.
 - Zinc forms coloured complexes.
 - Most of the *d*-block elements and their compounds are ferromagnetic.
 - Osmium shows +8 oxidation state.

- (5) Cobalt (II) is more stable in octahedral complexes.
 (a) 1 and 2 (b) 1 and 3
 (c) 2 and 5 (d) 1 and 4

10. Which of the following characteristics is true for $[\text{Ni}(\text{en})_2]^{2+}$ complex?
 (a) Paramagnetism, dsp^2 , square planar, coordination number of Ni = 2
 (b) Diamagnetism, dsp^2 , square planar, coordination number of Ni = 4
 (c) Diamagnetism, sp^3 , tetrahedral, coordination number of Ni = 4
 (d) Paramagnetism, sp^3 , tetrahedral, coordination number of Ni = 4
11. On heating KMnO_4 , which one among the following is not formed?
 (a) K_2MnO_4 (b) O_2
 (c) MnO_2 (d) MnO
12. Choose the correct name for $[\text{Co}(\text{NH}_3)_5\text{Cl}]\text{Cl}_2$.
 (a) Pentaamminechlorocobalt(II) chloride
 (b) Pentaamminechlorocobalt(III) chloride
 (c) Pentaamminechlorocobalt(III) dichloride
 (d) Pentaamminecobalt(III) chloride

Assertion & Reason Type

Directions : In the following questions, a statement of assertion is followed by a statement of reason. Mark the correct choice as :

- (a) If both assertion and reason are true and reason is the correct explanation of assertion.
 (b) If both assertion and reason are true but reason is not the correct explanation of assertion.
 (c) If assertion is true but reason is false.
 (d) If both assertion and reason are false.

13. **Assertion :** KMnO_4 is stored in dark coloured bottles.

Reason : On heating with alkalis, KMnO_4 is converted to manganate.

14. **Assertion :** F^- ion is a weak ligand and forms outer orbital complex.

Reason : F^- ion cannot force the electrons of d_{z^2} and $d_{x^2-y^2}$ orbitals of the inner shell to occupy d_{xy} , d_{yz} and d_{zx} orbitals of the same shell.

15. **Assertion :** There is an increase in oxidation potential of Fe^{2+} when it combines with a ligand CN^- .

Reason : Ionic activity of Fe^{2+} increases during complex formation of $[\text{Fe}(\text{CN})_6]^{4-}$.

JEE MAIN / JEE ADVANCED

Only One Option Correct Type

16. A complex containing Mn^{2+} and Br^- ions was found to have a magnetic moment of 5.92 B.M. The

probable formula and geometry of the complex is most likely to be

- (a) $[\text{MnBr}_4]^{2-}$, tetrahedral
 (b) $[\text{MnBr}_6]^{3-}$, octahedral
 (c) $[\text{MnBr}_4]^{2-}$, square planar
 (d) $[\text{MnBr}_5]^{3-}$, trigonal planar

17. Which of the following orders are correct?


- (I) $5d - 5d > 4d - 4d > 3d - 3d$ (orbital overlapping)
 (II) $\text{Mn}^{2+} > \text{Fe}^{2+} > \text{Cr}^{3+} > \text{Cu}^{2+}$ (magnetic moment)
 (III) $\text{Sc}^{3+} > \text{Zn}^{2+} > \text{V}^{5+} > \text{Ti}^{4+}$ (magnetic nature)
 (IV) $\text{Zn} > \text{Cu} > \text{Fe} > \text{Co}$ (atomic size)

- (a) Only I, II (b) Only I, IV
 (c) Only I, II, IV (d) Only II, III, IV

18. On treatment of $[\text{Pt}(\text{NH}_3)_4]^{2+}$ with concentrated HCl, two compounds (I) and (II) having the same formula, $[\text{Pt}(\text{NH}_3)_2\text{Cl}_2]$ are obtained, (I) can be converted into (II) by boiling with dilute HCl. A solution of (I) reacts with oxalic acid to form $[\text{Pt}(\text{NH}_3)_2(\text{C}_2\text{O}_4)]$ whereas (II) does not react. Choose the correct option from the following :

- (a) (I) *cis*, (II) *trans*; both tetrahedral
 (b) (I) *cis*, (II) *trans*; both square planar
 (c) (I) *trans*, (II) *cis*; both tetrahedral
 (d) (I) *trans*, (II) *cis*; both square planar

19. FeCr_2O_4 (chromite) is converted to Cr by the following steps :



I, II and III are

- | I | II | III |
|---|--------------------------------|--------------|
| (a) $\text{Na}_2\text{CO}_3/\text{air}, \Delta$ | C | C |
| (b) $\text{NaOH}/\text{air}, \Delta$ | C, Δ | Al, Δ |
| (c) $\text{NaOH}/\text{air}, \Delta$ | C, Δ | Mg, Δ |
| (d) conc. $\text{H}_2\text{SO}_4, \Delta$ | $\text{NH}_4\text{Cl}, \Delta$ | C, Δ |

More than One Options Correct Type

20. $\text{K}_4[\text{Fe}(\text{CN})_6]$ is used in the identification of

- (a) Fe^{2+} ions (b) Fe^{3+} ions
 (c) Cu^{2+} ions (d) Cd^{2+} ions.

21. Mark the incorrect statements.

- (a) Potassium dichromate oxidises a secondary alcohol into a ketone.
 (b) Potassium permanganate is a weaker oxidising agent than potassium dichromate.
 (c) Potassium permanganate is a stronger oxidising agent than potassium dichromate.
 (d) Potassium dichromate oxidises a secondary alcohol into an aldehyde.

22. In basic medium, the amount of Ni^{2+} in a solution can be estimated with the dimethylglyoxime reagent. The correct statements about the reaction and the product are
- in ammoniacal solution, Ni^{2+} salts give cherry-red precipitate of nickel(II) dimethylglyoximate
 - two dimethylglyoximate units are bound to one Ni^{2+}
 - in the complex, two dimethylglyoximate units are hydrogen bonded to each other
 - each dimethylglyoximate unit forms a six-membered chelate ring with Ni^{2+} .
23. When an acidified dichromate solution is shaken with a pinch of Na_2O_2
- a blue colouration is observed
 - orange red colouration finally changes to green
 - oxygen gas is evolved
 - bluish green precipitate is formed.

Integer Answer Type

24. The volume (in mL) of 0.1 M AgNO_3 required for complete precipitation of chloride ions present in 30 mL of 0.01 M solution of $[\text{Cr}(\text{H}_2\text{O})_5\text{Cl}]\text{Cl}_2$, as silver chloride, is close to
25. Amongst the following, the number of metals which do not form amalgams is
Ti, V, Cr Mn, Fe, Co, Ni, Pt, Cu, Zn
26. The complex, $\text{Ca}_2[\text{M}(\text{CN})_6]$ has spin only magnetic moment 2.83 B.M. and the complex $\text{K}_2[\text{MCl}_4]$ has spin only magnetic moment of 4.9 B.M. How many electrons are present in valence shell d -orbitals of the neutral gaseous atom M ?

Comprehension Type

The lanthanoids exhibit a principal oxidation state of +3. The +3 oxidation states in lanthanum, gadolinium and lutetium are stable because these ions have an empty, a half-filled and completely filled $4f$ -subshell respectively. But some of the lanthanoids exhibit different oxidation states.

27. Good oxidising agent out of the following lanthanoids is
- Ce^{3+}
 - Sm^{2+}
 - Ce^{4+}
 - Eu^{2+}
28. For which of the following lanthanoids, +2 oxidation state exists in aqueous solution?
- Ce, Yb, La
 - Ce, Eu, Ho
 - Er, Tb, Sm
 - Sm, Eu, Yb

Matrix Match Type

29. Match the magnetic moments listed in Column II with the electronic configurations of the complexes listed in Column I and choose the correct option using the codes given below.

Column I		Column II	
(A) d^3 (octahedral)		(p) 0.0 B.M.	
(B) d^5 (octahedral, low spin)		(q) $2\sqrt{6}$ B.M.	
(C) d^6 (octahedral, low spin)		(r) $\sqrt{15}$ B.M.	
(D) d^6 (octahedral, outer orbital)	(s) $\sqrt{3}$ B.M.		
A	B	C	D
(a) p	r	q	s
(b) r	s	p	q
(c) r	p	q	s
(d) q	r	p	s

30. Match the Column I with Column II and choose the correct option using the codes given below.

Column I		Column II	
(A) Aquated Mn^{2+} ion		(p) White	
(B) $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$		(q) Blue	
(C) Aquated V^{4+} ion		(r) Pink	
(D) Anhydrous CuSO_4		(s) Green	
A	B	C	D
(a) p	q	r	s
(b) p	s	q	r
(c) r	s	q	p
(d) q	p	r	s



Keys are published in this issue. Search now! ☺




SELF CHECK

No. of questions attempted
No. of questions correct
Marks scored in percentage

Check your score! If your score is

> 90%	EXCELLENT WORK !	You are well prepared to take the challenge of final exam.
90-75%	GOOD WORK !	You can score good in the final exam.
74-60%	SATISFACTORY !	You need to score more next time.
< 60%	NOT SATISFACTORY!	Revise thoroughly and strengthen your concepts.

Boost your fundamentals with MTG's Objective series



Deep knowledge and crystal clear understanding of fundamentals is key to success. MTG's Objective series is created keeping just this insight in mind for Class XI & XII students preparing to compete in entrance exams. Put together by MTG's renowned editorial team, these unique books ensure students get just the start they need.

HIGHLIGHTS:

- 5,255+ pages covering the latest syllabus of NEET and other entrance exams
- Check-Your-Grasp questions for self-assessment
- NCERT xtract from NCERT books
- Question Banks including questions from previous years' test papers (solved) of various exams like AIIMS, AIPMT / NEET, AFMC, BHU, AMU, J&K CET, UGET-Manipal, Odisha, UP-CPMT, WB JEE, Kerala PMT, etc.
- Detailed solutions to MCQs for clear understanding
- Additional information for students for exams such as AIIMS, NEET etc.



Scan now with your
smartphone or tablet*



Available at all leading book shops throughout India.
For more information or for help in placing your order:
Call 0124-6601200 or email info@mtg.in

*Application to read QR codes required

Visit
www.mtg.in
for latest offers
and to buy
online!

CHEMISTRY Musing

PROBLEM SET 50

Chemistry Musing was started from August '13 issue of Chemistry Today. The aim of Chemistry Musing is to augment the chances of bright students preparing for JEE (Main and Advanced) / NEET / AIIMS / JIPMER with additional study material. In every issue of Chemistry Today, 10 challenging problems are proposed in various topics of JEE (Main and Advanced) / NEET. The detailed solutions of these problems will be published in next issue of Chemistry Today. The readers who have solved five or more problems may send their solutions. The names of those who send atleast five correct solutions will be published in the next issue. We hope that our readers will enrich their problem solving skills through "Chemistry Musing" and stand in better stead while facing the competitive exams.


JEE MAIN/NEET

- A cylinder of gas is assumed to contain 11.2 kg of butane. If a normal family needs 20,000 kJ of energy per day for cooking, how long will the cylinder last? (Given : the heat of combustion of butane is 2658 kJ mol^{-1})
 (a) 27 days (b) 3700 min
 (c) 26 days (d) 5000 s
- The equilibrium constant K_p for the reaction,


$$\text{N}_2 + 3\text{H}_2 \rightleftharpoons 2\text{NH}_3$$
 is 1.64×10^{-4} at 400°C . What will be the equilibrium constant at 500°C if the heat of reaction in this temperature range is $-25,140$ calories?
 (a) 1.4×10^{-5} (b) 3.6×10^{-6}
 (c) 2.8×10^{-10} (d) 10^{-10}
- $\text{PhMe} \xrightarrow[\text{H}_3\text{O}^+]{\text{KMnO}_4/\text{OH}^-} (\text{B}) \xrightarrow[\text{or PCl}_5]{\text{SOCl}_2} (\text{C}) \xrightarrow{\text{PhCOONa}} (\text{D})$
 Compound (D) is
 (a) PhCOCl (b) PhCONH_2
 (c) PhCOOH (d) $(\text{PhCO})_2\text{O}$
- Two compounds have the empirical formula, $\text{Cr}(\text{NH}_3)_3(\text{NO}_2)_3$. In aqueous solution, one of these conducts electricity while the other does not. What is the probable structure of the compound which conducts electricity?
 (a) $[\text{Cr}(\text{NH}_3)_3(\text{NO}_2)_3]$ (b) $[\text{Cr}(\text{NH}_3)_2(\text{NO}_2)_3]\text{NH}_3$
 (c) $[\text{Cr}(\text{H}_2\text{O})(\text{NH}_3)_3(\text{NO}_2)_2]\text{NO}_2$
 (d) None of these
- Choose the correct arrangement of the following in decreasing order of $\left(\frac{m}{o+p}\right)$ ratio.
 I. PhCH_3 II. PhCBr_3
 III. PhCCl_3 IV. PhCl_3
 (a) $\text{II} > \text{III} > \text{I} > \text{IV}$ (b) $\text{IV} > \text{II} > \text{III} > \text{I}$
 (c) $\text{III} > \text{II} > \text{I} > \text{IV}$ (d) $\text{I} > \text{III} > \text{II} > \text{IV}$

JEE ADVANCED


- An optically active compound (A) of formula, $\text{C}_9\text{H}_{12}\text{O}_2$ produced the following compound when refluxed with KMnO_4 .



The original compound showed these properties also :



Which of the following is the structure of (A)?




COMPREHENSION

- A yellow powder X is burnt in a stream of fluorine to obtain a colourless gas, Y which is thermally stable and chemically inert. Its molecule has octahedral geometry. Another colourless gas, Z with same constituent atoms as that of Y is obtained when sulphur dichloride is heated with sodium fluoride. Its molecule has trigonal bipyramidal geometry.
- The yellow powder X is
 (a) $\text{K}_2\text{Cr}_2\text{O}_7$ (b) FeCl_3
 (c) K_2CrO_4 (d) S
 - The colourless gases Y and Z respectively are
 (a) SF_4, SF_4 (b) SF_6, SF_4
 (c) NaF, SF_6 (d) $\text{S}_2\text{F}_2, \text{NaF}$

INTEGER VALUE

- Total number of aldol condensation products of the following reaction is



- The reaction, $\text{SO}_2\text{Cl}_2 \xrightarrow{k_1} \text{SO}_2 + \text{Cl}_2$ is a first order reaction with $k_1 = 2.2 \times 10^{-5} \text{ sec}^{-1}$ at 302°C . The percentage of SO_2Cl_2 that will get decomposed in 90 minutes when the reaction is carried out at 302°C is $(8.2 + y)$. The value of y is




JEE Advanced

PRACTICE PROBLEMS

SECTION 1 (Maximum Marks : 18)


This section contains 6 multiple choice questions. Each question has 4 choices (a), (b), (c) and (d), out of which only one is correct. For each question you will be awarded 3 marks if you have darkened only the bubble corresponding to the correct answer and zero mark if no bubble is darkened. In all other cases, minus one (-1) mark will be awarded.

1. The enthalpy changes at 298 K in successive breaking of O—H bonds of HOH are



The bond enthalpy of O—H bond is

- (a) 498 kJ mol⁻¹ (b) 463 kJ mol⁻¹
 (c) 428 kJ mol⁻¹ (d) 70 kJ mol⁻¹
2. An ideal monoatomic gas is taken round the cycle ABCDA as shown in the figure. The work done during the cycle is




- (a) $-PV$
 (b) $-2PV$
 (c) $-\frac{1}{2}PV$
 (d) zero.
3. A given mass of gas expands from the state X to state Y by three paths 1, 2 and 3 as shown in the figure. If w_1 , w_2 and w_3 respectively be the work done by the gas along these three paths then
- (a) $w_1 > w_2 > w_3$ (b) $w_2 < w_1 < w_3$
 (c) $w_1 = w_2 = w_3$ (d) $w_1 < w_2 < w_3$
4. Identify the reaction in which the heat liberated corresponds to the heat of formation (ΔH).
- (a) $\text{C}_{(\text{diamond})} + \text{O}_{2(g)} \rightarrow \text{CO}_{2(g)} + \text{heat}$
 (b) $2\text{H}_{2(g)} + \text{O}_{2(g)} \rightarrow 2\text{H}_2\text{O}_{(g)} + \text{heat}$
 (c) $\text{C}_{(\text{diamond})} + 2\text{H}_{2(g)} \rightarrow \text{CH}_{4(g)} + \text{heat}$
 (d) $\text{S}_{(\text{rhombic})} + \text{O}_{2(g)} \rightarrow \text{SO}_{2(g)} + \text{heat}$

5. The entropy change in the fusion of one mole of a solid melting at 27°C is (latent heat of fusion = 2930 J mol⁻¹)

- (a) 9.77 J K⁻¹ mol⁻¹ (b) 10.73 J K⁻¹ mol⁻¹
 (c) 2930 J K⁻¹ mol⁻¹ (d) 108.5 J K⁻¹ mol⁻¹

6. Two moles of a monoatomic ideal gas are taken through a cyclic process starting from A as shown in the figure. The volume ratios are $\frac{V_B}{V_A} = 2$ and $\frac{V_D}{V_A} = 4$.



If the temperature T_A at A is

27°C, calculate the total heat absorbed (in calories) in the cyclic process.

- (a) 1080 (b) 900 (c) 600 (d) 1200

SECTION 2 (Maximum Marks : 08)


This section contains 2 multiple choice questions. Each question has 4 choices (a), (b), (c) and (d), out of which one or more than one answer is correct. For each question you will be awarded 4 marks if you have darkened only the bubble corresponding to the correct answer and zero mark if no bubble is darkened. No negative marking in this section.

7. Five moles of an ideal gas at 293 K are expanded isothermally from an initial pressure of 0.4 kPa to a final pressure of 0.1 kPa against a constant external pressure of 0.1 kPa. Select the correct option(s).
- (a) $q = 0$ and $w \neq 0$
 (b) $q = 0$ and $w = 0$
 (c) $\Delta E = 0$ and $\Delta H \neq 0$
 (d) $\Delta E = 0$ and $\Delta H = 0$
8. In an insulated container, 1 mole of a liquid (molar volume 100 mL) at 1 bar is steeply taken to 100 bar when volume of liquid decreases by 1 mL. Select the correct option(s).
- (a) $\Delta U = 99 \text{ bar mL}$ (b) $\Delta U = 100 \text{ bar mL}$
 (c) $\Delta H = 9900 \text{ bar mL}$ (d) $\Delta H = 100 \text{ bar mL}$

SECTION 3 (Maximum Marks : 24)

This section contains 6 integer type questions. Answers have to be given in between 0 to 9 in the form of nearest integer. Each question carries 4 marks if you have darkened the correct answer and no negative mark will be awarded for an incorrectly bubbled answer.

9. 3.5 g of a gas (mol. wt. = 28) was burnt in excess of oxygen at 298 K in a constant volume calorimeter. The temperature of the calorimeter was found to increase to 298.45 K. The enthalpy of combustion (in kJ mol^{-1}) is (Heat capacity of calorimeter = 2.5 kJ K^{-1})
10. The heat of formation of ethane is -19.46 kcal . Bond energies of H—H, C—H and C—C bonds are 104.18, 99.0 and 80.0 kcal respectively. The heat of atomization of graphite can be expressed in scientific notation as $p \times 10^q$ calories. Identify p .
11. For a hypothetical reaction, $A_{2(g)} + B_{2(g)} \rightleftharpoons 2AB_{(g)}$ ΔG° and ΔS° are 20 kJ/mol and $-20 \text{ J K}^{-1} \text{ mol}^{-1}$ respectively at 200 K. If ΔC_p is $20 \text{ J K}^{-1} \text{ mol}^{-1}$ then ΔH° at 400 K is x . Find $\frac{x}{4}$?



12. An ideal gas is taken around the cycle ABCA as shown in the figure. Work done in the cyclic process = xP_1V_1 . What is the value of x ?
13. In a thermodynamic process, helium gas obeys the law $\frac{T}{p^{2/5}} = \text{constant}$. The heat given to n moles of He in order to raise the temperature from T to $2T$ is
14. In the process :
 $\text{H}_2\text{O}_{(s)} (-10^\circ\text{C}, 1 \text{ atm}) \rightarrow \text{H}_2\text{O}_{(l)} (10^\circ\text{C}, 1 \text{ atm})$
 C_p for ice = $9 \text{ cal deg}^{-1} \text{ mol}^{-1}$,
 C_p for H_2O = $18 \text{ cal deg}^{-1} \text{ mol}^{-1}$
 Latent heat of fusion of ice = $1440 \text{ cal mol}^{-1}$ at 0°C
 The entropy change for the above process will be

SECTION 4 (Maximum Marks : 06)

This section contains 1 paragraph. Based upon paragraph, 2 multiple choice questions have to be answered. For each question you will be awarded 3 marks if you have darkened only the bubble corresponding to the correct answer and zero mark if no bubble is darkened. In all other cases, minus one (-1) mark will be awarded.

PARAGRAPH

Chemical reactions can be viewed from a thermodynamic perspective. Change in enthalpy (ΔH) and entropy (ΔS) are the two driving forces that determine whether a reaction is spontaneous. Gibbs free energy (ΔG) combines these two factors in one equation :

$$\Delta G = \Delta H - T\Delta S$$

Compound	$\Delta H_f^\circ(\text{kJ/mol})$	$S^\circ(\text{J/mol-K})$
$\text{N}_{2(g)}$	0	192
$\text{H}_{2(g)}$	0	131
$\text{NH}_{3(g)}$	-46	192

15. What is the value of ΔH° for the reaction,
 $\text{N}_{2(g)} + 3\text{H}_{2(g)} \rightarrow 2\text{NH}_{3(g)}$?
- (a) -46 kJ (b) $-92,000 \text{ J}$
 (c) 0 J (d) 138 kJ
16. What is ΔS° for the reaction, $\text{N}_{2(g)} + 3\text{H}_{2(g)} \rightarrow 2\text{NH}_{3(g)}$?
- (a) -201 J/K (b) -192 J/K
 (c) -131 J/K (d) 201 J/K

SECTION 5 (Maximum Marks : 16)

This section contains 2 questions. Each question contains statements given in 2 columns. Statements in the first column have to be matched with statements in the second column. For each question in this section, you will be awarded 8 marks if you have darkened all the bubbles corresponding only to the correct answer and 2 marks for each row. No negative mark will be awarded for an incorrectly bubbled answer.

17. Match the Column I with Column II.
- | Column I | Column II |
|---|--|
| (A) $\text{Ice} \rightleftharpoons \text{H}_2\text{O}_{(l)}$ at room temperature 25°C | (p) $\Delta H = -\text{ve}, \Delta S = +\text{ve}$ |
| (B) $\text{H}_2\text{O}_{(l)} \rightleftharpoons \text{Ice}$ at room temperature 25°C | (q) $\Delta G = +\text{ve}$ |
| (C) $2\text{O}_{3(g)} \rightarrow 3\text{O}_{2(g)}$ | (r) $\Delta H = +\text{ve}, \Delta S = -\text{ve}$ |
| (D) $3\text{O}_{2(g)} \rightarrow 2\text{O}_{3(g)}$ | (s) $\Delta G = -\text{ve}$ |
18. Match the Column I with Column II.
- | Column I | Column II |
|--|---|
| (A) $\text{Ice} \rightleftharpoons \text{Water}$ at 273 K | (p) $\Delta H < 0, \Delta S < 0$ |
| (B) $\text{NH}_{3(g)} + \text{HCl}_{(g)} \rightarrow \text{NH}_4\text{Cl}_{(s)}$ | (q) $\Delta G = 0$ |
| (C) $2\text{NH}_{3(g)} \rightarrow 3\text{H}_{2(g)} + \text{N}_{2(g)}$ | (r) $\Delta n > 0, \Delta H > \Delta U$ |
| (D) $\text{CO}_{2(g)} \rightarrow \text{Dry ice}$ | (s) $\Delta n < 0, \Delta H < \Delta U$ |

ANSWER KEY

1. (b) 2. (a) 3. (d) 4. (d) 5. (a)
 6. (d) 7. (d) 8. (b,c) 9. (9) 10. (2)
 11. (5) 12. (3) 13. (0) 14. (6) 15. (b)
 16. (a) 17. (A) - (s), (B) - (q), (C) - (p, s); (D) - (q, r)
 18. (A) - (q), (B) - (p, s), (C) - (r); (D) - (p, s)

For detailed solutions to the
JEE Advanced Practice Problems
 visit our website: www.vidyalankar.org





CONCEPT BOOSTER

Hi all!! Hope you all are doing well. This is the last part of the consecutive 'SHORTCUT OF INORGANIC CHEMISTRY' article. Inorganic Chemistry is endless but what I have tried is to give you an idea, how to read Inorganic Chemistry. Hope you liked it. Keep practicing harder and harder.

*Arunava Sarkar

HEATING EFFECTS


HEATING EFFECT OF DICHROMATE SALTS





HEATING EFFECT OF PHOSPHATE SALTS

Shortcut : Remove only water and if any gas is possible (like NH_3 but not O_2 and oxides of P) then that too.



- Now, phosphoric acid is H_3PO_4 . So, the first salt that comes to our mind is NaH_2PO_4 (one acidic hydrogen substitution). It is 1° phosphate salt.



(Just remove H_2O ; don't think much.)







- Ammonium containing phosphate salts, of course evolve NH_3 along with water on heating. For example,



HEATING EFFECT OF OXALATE SALTS

- In general, more reactive metal's oxalate salts give both CO and CO_2 on heating whereas less reactive metals give CO_2 only.










HEATING EFFECT OF ACETATE SALTS

A few points must be remembered here :

- Formation of acetone is inevitable as this is a general method of preparation of acetone.
- Smaller alkaline earth metals like Be, Mg (first two in the group) cannot hold bigger CO_3^{2-} tightly with them and they prefer to exist in the form of oxide. So, for them, metal oxide and CO_2 will be produced. But, for relatively larger metals like Ca, Ba, etc. carbonate salts will be produced.
- For alkali metals, carbonate salts are obtained.

Let us check :








HEATING EFFECT OF FORMATE SALTS

Three major points to be remembered :


- Lower alkali metals give corresponding oxalates.
- Alkaline earth metals give formaldehyde (in general).
- Silver and mercury salts as usual give metal back along with formic acid. Let us see :



HEATING EFFECT OF BORIC ACID



HEATING EFFECT OF OXALIC ACID



MPP-5 CLASS XII

ANSWER KEY

- | | | | | |
|-----------|-------------|-------------|---------|-----------|
| 1. (b) | 2. (a) | 3. (b) | 4. (c) | 5. (a) |
| 6. (d) | 7. (d) | 8. (c) | 9. (d) | 10. (b) |
| 11. (d) | 12. (b) | 13. (b) | 14. (a) | 15. (c) |
| 16. (a) | 17. (c) | 18. (b) | 19. (b) | 20. (b,c) |
| 21. (b,d) | 22. (a,b,c) | 23. (a,b,c) | 24. (6) | 25. (2) |
| 26. (5) | 27. (c) | 28. (d) | 29. (b) | 30. (c) |

Buy digital copies
of your favorite
magazine



Visit Now : digital.mtg.in

Study Anytime ! Anywhere !



mtg



ADVANCED CHEMISTRY BLOC

(AN INSIGHT TO MOT)

Mukul C. Ray, Odisha

LINEAR COMBINATION OF ATOMIC ORBITALS (LCAO)

The definition goes like this.

“The simplest trial functions that work reasonably well in molecular orbital theory turn out to be linear combination of atomic orbitals, or LCAO”.


But what is the trial function? In molecular orbital theory (MOT), you actually do not solve Schrödinger (ö is a German umlaut, read it as Schroedinger) equation. It's not that easy. Instead you use a trial function, an "educated guess" as to what the solution might be. Suppose you want to know the value of x in the equation, $4x + 5 = 70$ without solving. Without solving you just guess a value of x and see how it works. In MOT, the estimating procedure is analogous. In order to determine how well a trial function for an orbital works in MOT, you calculate energy. Your trial wave function should try to approach as close as to the energy of the actual orbital. In MOT, the trial wave function for molecular orbitals are weighted linear sum of valence atomic orbitals of atoms in a molecule. When I say weighted that means you need not necessarily take equal contribution of atomic orbital wave functions of combining atoms.

Number of atomic orbitals combining is equal to the number of molecular orbitals formed. It's not new, I am sure, for you that is atomic orbitals combine to give σ_{1s} and σ^*_{1s} , former having lower energy and later higher energy than combining atomic orbitals.

In VBT, we treat electrons as if they reside in the atomic orbital. This is an over simplification. We tried in a better way in MOT. The bonding molecular orbital arises out of constructive interference between the atomic orbitals because both orbitals have the same

phase. The antibonding orbital arises out of destructive interference.

You can compare the situation like the following :



Say, you are walking along the arrow mark you will be able to notice remarkably neat pattern of alternating maxima and minima produced from two identical sources. This is superposition of waves from two loud speakers. The atomic orbital wave functions almost behave similarly.


The key point to understand is that, every electron which enters a bonding molecular orbital stabilises the molecule or polyatomic ion and every electron which enters an antibonding molecular orbital destabilises it. The emphasis on electron pair has been removed. Also, an electron when removed from bonding orbital, destabilises the system and an electron when added to antibonding orbital also destabilises the system but adding an electron to antibonding orbital destabilises it more.

Applications


○ Stability : $H_2 > H_2^+ > H_2^-$

Even though the bond order of H_2^+ and H_2^- are same, H_2^- is less stable. The paragraph just above explains this observation.

- For second period diatomic molecules, O_2 and F_2 , the sequence of energy of molecular orbital is $(\sigma 2s) (\sigma^* 2s) (\sigma 2p_z) (\pi 2p_x = \pi 2p_y) (\pi^* 2p_x = \pi^* 2p_y) (\sigma^* 2p_z)$. It's simple to handle, O_2 , O_2^+ , O_2^- , O_2^{2-} , F_2 , etc. You just feel the electrons from lower end considering each molecular orbital takes maximum 2 electrons and for degenerate molecular orbital, Hund's rule is applicable.
- For second period diatomic molecules on the left of oxygen like B_2 , C_2 , N_2 , the energy sequence is slightly altered. The reason is that the energy gap between $2s$ and $2p$ is less in these atoms B, C, N, etc.




Also you must note the stability sequence :



The explanation is the same as for H_2^+ and H_2^- .

- For CO, the two hybrid orbitals are directed at 180° to each other.




Solution Senders of Chemistry Musing

Set - 49



- Ravinder Kashapogu, Telangana
- Naveen Soni, Gujarat
- Akila Iyer, Bengaluru

One hybrid orbital is directed dimetrically opposite to the CO bonding region. Hence, it does not form bond but retains a pair of electron (σ_3).





σ_3 is primarily of non-bonding nature. The pair of electrons present in σ_3 is the carbon based lone pair, which is approximately 126 pm far from carbon in opposite direction to oxygen. This lowers the dipole moment of CO molecule significantly down to 0.112 D.

When CO ionises to CO^+ , the carbon-oxygen bond length decreases from 112.8 pm to 111.5 pm. Some author have an opinion that bond order has increased from 3 to 3.5 in CO^+ . But this is not very reasonable as second period elements can never form more than three bonds. (Though Re forms quadruple bond in $Re_2Cl_8^{2-}$ ion). The better explanation comes from VBT. A triple bond that has partial dative bond character develops pure covalent character in order to distribute the net positive charge on both the atoms. As an evidence, we can see that dative bonds are longer than isoelectronic covalent bonds. For example,



Also,



So, when the dative bond of CO changes to covalent bond in CO^+ , the bond length decreases.



CHEMISTRY MUSING

SOLUTION SET 49

1. (a): Let radius of hollow sphere X be r .

$$\therefore \text{Edge length } (a) = 4r / \sqrt{3}$$

$$\text{Volume of unit cell} = a^3 = (4r / \sqrt{3})^3$$


Volume of X unoccupied by Y (having radius = $r/2$)

$$\text{in unit cell} = 2 \times \left[\frac{4}{3} \pi r^3 - \frac{4}{3} \pi \left(\frac{r}{2} \right)^3 \right]$$

$$\therefore \frac{\text{Volume of X unoccupied by Y in unit cell}}{\text{Volume of unit cell}}$$


$$= \frac{\frac{4}{3} \pi \times \frac{7r^3}{8} \times 2}{\left(\frac{4r}{\sqrt{3}} \right)^3} = \frac{7\pi\sqrt{3}}{64}$$

2. (c):




P = Be

Q = B



Formic acid is different from other monocarboxylic acids as it can act as a reducing agent, thus it reduces Fehling solution to red cuprous oxide.



5. (c): $\text{BOH} + \text{HCl} \rightarrow \text{BCl} + \text{H}_2\text{O}$

At equivalence point, $M_1V_1 = M_2V_2$

$$\frac{2}{5} \times 2.5 = \frac{2}{15} \times V_2, \quad V_2 = 7.5 \text{ mL}, \quad V_{\text{HCl}} = 7.5 \text{ mL}$$

Moles of BOH in 2.5 mL = 0.001

Moles of HCl in 7.5 mL = 0.001

\therefore Moles of salt formed = 0.001

Total volume = 2.5 + 7.5 = 10 mL = 0.01 L

$$\text{Conc. of salt} = \frac{0.001}{0.01} = 0.1 \text{ mole/litre}$$

Hydrolysis of salt takes place:

$$\text{pH} = 7 - \frac{1}{2} (\text{p}K_b + \log C)$$

$$\text{pH} = 7 - \frac{1}{2} (12 - 1)$$

$$[\because \text{p}K_b = -\log 10^{-12} = 12 \text{ and } \log 0.1 = -1]$$

$$\text{pH} = 1.5 \Rightarrow -\log[\text{H}^+] = 1.5$$

$$\Rightarrow \log [\text{H}^+] = -1.5$$

$$[\text{H}^+] = 3.2 \times 10^{-2} \text{ M}$$

6. (d): 374 kJ is required for 1 mole

$$\therefore 22.44 \text{ kJ is required for} = \frac{1}{374} \times 22.44 \text{ moles} = 0.06$$




$$\text{Moles} = \frac{\text{mass}}{\text{Atomic mass}}$$



$$0.06 = \frac{8}{\text{Atomic mass}}, \quad \text{Atomic mass} = 133.33$$

Moles of $M_{(g)}^+$ formed = 0.06

$$\text{Thus, atoms of } M \text{ converted to } M_{(g)}^+ = 0.06 \times 6.023 \times 10^{23} = 3.613 \times 10^{22} \text{ atoms}$$

7. (b):





Total molality after the addition of Hg(CN)_2
 $=$ molality of K^+ + molality of CN^-
 $+ \text{molality of } [\text{Hg(CN)}_{x+2}]^{x-}$
 $= 0.1892 + (0.1892 - 0.095x) + 0.095$
 $= (0.4734 - 0.095x)$

Now, $K_f = \frac{\Delta T_f}{m} \Rightarrow 1.86 = \frac{0.53}{0.4734 - 0.095x}$
 $x = 2.$

10. (2): The kinetic energy of the electron is provided by accelerating potential,

$V = \frac{h^2}{\lambda^2 \times 2 \times e \times m}$
 $\lambda = 8.7 \text{ pm} = 8.7 \times 10^{-12} \text{ m}$
 $e = 1.6 \times 10^{-19} \text{ C}$
 $m = 9.1 \times 10^{-31} \text{ kg}$
 $h = 6.63 \times 10^{-34} \text{ J s}$

$V = \frac{(6.63 \times 10^{-34} \text{ J s})^2}{(8.7 \times 10^{-12} \text{ m})^2 \times 2 \times (1.6 \times 10^{-19} \text{ C}) \times 9.1 \times 10^{-31} \text{ kg}}$
 $= 1.99 \times 10^4 \text{ kg m}^2 \text{ s}^{-2} \text{ C} \approx 2 \times 10^4 \text{ V}$

ATTENTION COACHING INSTITUTES: a great offer from MTG

MTG offers "Classroom Study Material" for JEE (Main & Advanced), NEET and FOUNDATION MATERIAL for Class 6, 7, 8, 9, 10, 11 & 12 with YOUR BRAND NAME & COVER DESIGN.

This study material will save you lots of money spent on teachers, typing, proof-reading and printing. Also, you will save enormous time. Normally, a good study material takes 2 years to develop. But you can have the material printed with your logo delivered at your doorstep.

Profit from associating with MTG Brand – the most popular name in educational publishing for JEE (Main & Advanced)/NEET/PMT....

Order sample chapters on Phone/Fax/e-mail.


Phone : 0124-6601200

09312680856

e-mail : sales@mtg.in | www.mtg.in




CLASSROOM STUDY MATERIAL



Your logo here

CROSSWORD




Readers can send their responses at editor@mtg.in or post us with complete address by 25th of every month to win exciting prizes. Winners' name with their valuable feedback will be published in next issue.

ACROSS

- A mildly toxic, white crystalline organic compound belonging to the indole family. It occurs naturally in faeces and coal tar and has a strong faecal odour. (7)
- An aminopyridine that functions as a centrally acting non-opioid analgesic that was originally used as an analgesic for acute and chronic pain. (10)
- A gelatinous precipitate formed by coagulation of a sol. (6)
- A German scientist who prepared almost perfectly semipermeable membrane using gelatinous copper ferrocyanide, $\text{Cu}_2[\text{Fe}(\text{CN})_6]$. (6)
- Naturally occurring alloy of gold and silver (upto 45% Ag). It resembles pure gold in appearance. (8)
- An atomic unit of energy which is equal to e^2/a_0 (where e is the charge of an electron and a_0 is the atomic unit of length). (7)
- An arsenic derivative that causes blisters and used in chemical warfare. (8)
- An instrument used to determine fermentation efficiency of yeast by measuring the amount of carbon dioxide produced from a given quantity of sugar. (9)
- The action of heating a substance until it burns away rapidly. (12)
- _____ law which expresses the dependence of a reaction enthalpy on temperature. (9)
- A stable heterocyclic octacirculene based on thiophene. (9)
- A polycyclic alkane with a structure similar to basket. (9)
- Fraction which is defined as the mass defect divided by the mass number. (7)
- A mineral deposit of gravel, sand and nitrates, found in dry areas of America. (7)





DOWN

- Small, granular, opaque diamonds, used as an abrasive in cutting tools. (4)
- The contraction of a gel accompanied by the separating out of liquid. (9)
- A technique in which a stable atom in a compound is replaced by its radioactive isotope. (9)
- _____ include all quarks and leptons, as well as any composite particle made of an odd number of these, such as all baryons and many atoms and nuclei. (8)
- A metal oxide obtained by heating an ore to high temperature in air. (4)



- A reactor which produces more fissile nuclei than it consumes. (7)
- A thermodynamic property of a real gas which is effective partial pressure. (8)
- Polymethine fibre with elastic properties. (7)
- Equation which gives relation between effect of temperature on surface tension. (6)
- A mixture of two or more components that reaches an equilibrium temperature which is independent of the temperature of any of its components before they are mixed. (10)
- The rule which is a practical aid for determining the configuration of chiral centers of amino acids. (4)
- An organothiophosphate insecticide primarily used on corn. (7)
- An analogue of pyrrole in which an arsenic atom replaces the nitrogen atom. (6)
- The tendency of a solid material to move slowly or deform permanently under the influence of mechanical stress. (5)
- A general name for alloys containing Al, Ni, Co and Fe, used as permanent magnets. (6)
- A very hard natural form of silica, used for knife-edges of balances and in ornaments. (5)

Now, save up to Rs 2,020*



Subscribe to MTG magazines today.

Our 2017 offers are here. Pick the combo best suited for your needs. Fill-in the Subscription Form at the bottom and mail it to us today. If in a rush, log on to www.mtg.in now to subscribe online.

*On cover price of ₹ 30/- each.

For JEE (Main & Advanced), NEET, AIIMS AND JIPMER

About MTG's Magazines

Perfect for students who like to prepare at a steady pace, MTG's magazines-Physics For You, Chemistry Today, Mathematics Today & Biology Today-ensure you practice bit by bit, month by month, to build all-round command over key subjects. Did you know these magazines are the only source for solved test papers of all national and state level engineering and medical college entrance exams?

Trust of over 1 Crore readers since 1982.

- Practice steadily, paced month by month, with very-similar & model test papers
- Self-assessment tests for you to evaluate your readiness and confidence for the big exams
- Content put together by a team

- comprising experts and members from MTG's well-experienced Editorial Board
- Stay up-to-date with important information such as examination dates, trends & changes in syllabi
- All-round skill enhancement –

- confidence-building exercises, new studying techniques, time management, even advice from past JEE/NEET toppers
- **Bonus:** Exposure to competition at a global level, with questions from Intl. Olympiads & Contests

SUBSCRIPTION FORM

Please accept my subscription to:

Note: Magazines are despatched by Book-Post on 4th of every month (each magazine separately).

Tick the appropriate box.

PCMB combo

- 1 yr: ₹ 1,000 (save ₹ 440) 2 yr: ₹ 1,800 (save ₹ 1,080) 3 yr: ₹ 2,300 (save ₹ 2,020)

PCM combo

- 1 yr: ₹ 900 (save ₹ 180) 2 yr: ₹ 1,500 (save ₹ 660) 3 yr: ₹ 1,900 (save ₹ 1,340)

PCB combo

- 1 yr: ₹ 900 (save ₹ 180) 2 yr: ₹ 1,500 (save ₹ 660) 3 yr: ₹ 1,900 (save ₹ 1,340)

Individual magazines

Physics Chemistry Mathematics Biology

- 1 yr: ₹ 330 (save ₹ 30) 2 yr: ₹ 600 (save ₹ 120) 3 yr: ₹ 775 (save ₹ 305)

Enclose Demand Draft favouring **MTG Learning Media (P) Ltd.** payable at New Delhi. You can also pay via Money Orders. Mail this Subscription Form to Subscription Dept., **MTG Learning Media (P) Ltd.**, Plot 99, Sector 44, Gurgaon – 122 003 (HR).

Want the magazines by courier; add the courier charges given below:

- 1 yr: ₹ 240 2 yr: ₹ 450 3 yr: ₹ 600

Tick the appropriate box.

Student Class XI XII Teacher Library Coaching

Name: _____

Complete Postal Address: _____

Pin Code Mobile #


Other Phone # 0

Email: _____

E-mail subscription@mtg.in. Visit www.mtg.in to subscribe online. Call (0)8800255334/5 for more info.

Presenting India's No.1 NEET Guides

STUDENTS'
No.1 CHOICE
for NEET
preparation



MTG's Complete NEET Guides are India's best selling PMT books!! Rich in theoretical knowledge with a vast question bank comprising a wide variety of problems and exercises, these guidebooks ensure students are ready to compete in the toughest of medical entrance tests. 100% NCERT based, the guidebooks have been updated to match the syllabus and the exam pattern for medical entrance exams. No wonder these guidebooks emerged as the bestsellers in a short period of time.

HIGHLIGHTS:

- 100% NCERT based
- Comprehensive Chapterwise theory complemented with concept maps, flowcharts and easy-to-understand illustrations
- Last 10 years' questions (2008-2017) of AIPMT/NEET
- Chapterwise Topicwise MCQs with detailed explanations and solutions
- NEET 2017 Solved Paper included
- Over 50% of questions that appeared in NEET 2017 were from MTG's Complete NEET Guides



Scan now with your
smartphone or tablet*




Available at all leading book shops throughout India.
For more information or for help in placing your order:
Call 0124-6601200 or e-mail:info@mtg.in

*Application to read QR codes required

Visit
www.mtg.in
for latest offers
and to buy
online!

THE PERFECT COACH

Now also available for JEE!



Skill. Passion. Hard work and determination. As a student sitting for the highly competitive JEE, you need all that. However, only a few will win, very likely with the help of a champion coach.

MTG's Champion Series for JEE is just the coach you need. It will guide you in identifying what's important for success and what's not. And then help you check your readiness with its most comprehensive question bank. So you know your strengths and weaknesses right from the word go and course-correct accordingly. Put simply, MTG's Champion Series will help you manage your preparation effort for JEE for maximum outcome. The best part is you study at a pace you're comfortable with.


Because it's all chapterwise, topicwise.



Visit www.MTG.in to buy online. Or visit a leading bookseller near you.
For more information, email info@mtg.in or call 1800 300 23355 (toll-free) today.



The most comprehensive question bank books that you cannot afford to ignore



30 Years' Physics, Chemistry & Biology contain not only chapterwise questions that have appeared over the last 30 years in NEET/AIPMT, but also full solutions, that too by experts. Needless to say, these question banks are essential for any student to compete successfully in NEET.

HIGHLIGHTS:

- Chapterwise questions of last 30 years' (2017-1988) of NEET/AIPMT
- Chapterwise segregation of questions to help you assess the level of effort required to succeed
- An unmatched question bank series with close to 1,000 pages having detailed solutions by experts
- Fully solved questions of NEET 2017 included




Scan now with your
smartphone or tablet*



Available at all leading book shops throughout India.
For more information or for help in placing your order:
Call 0124-6601200 or email info@mtg.in

*Application to read QR codes required

Visit
www.mtg.in
for latest offers
and to buy
online!



Inviting JEE and NEET aspirants

to participate in the SOF National Science Olympiad to see where they stand vis-à-vis competition



**SOF NATIONAL
SCIENCE OLYMPIAD**

NOV. 9 & NOV. 23



With much in common with competitive exams held each year for admission into leading engineering & medical colleges, the SOF National Science Olympiad is a great pre-testing platform for students to assess their chances of success in JEE or NEET etc. Millions of students compete in the SOF Olympiads. To participate, all you need to do is contact your School Principal or your schools SOF Olympiads Coordinator immediately. For more info, log on to www.sofworld.org. Or just scan the QR code with your smartphone or tablet.

Registrations closing soon. Hurry! Contact your Principal / School SOF Olympiads Coordinator today.



iranchembook.ir/edu

THIS IS ALMOST AN EVERY DAY SCENARIO AT V-SPARC

VIT School of Planning and ARchitecture



5 YEAR B.ARCH PROGRAMME VELLORE CAMPUS

IGNITE THE SPARK TODAY!

Valid
NATA score
mandatory

- Practical expertise through continuous hands-on learning
- Site exploration both digital & manual
- Professionals from around the globe impart knowledge
- Direct correlation with industry standards
- Syllabus includes topics from interactions with industry experts
- Application-centric pedagogy
- Global standards for students at par with peers abroad
- Inter-University collaborations

For further details please visit www.vit.ac.in

CONTACT DETAILS

Director - UG Admissions,

Phone: 0416-220 2157/2168, Email: ugadmission@vit.ac.in



VIT[®]
UNIVERSITY
(Estd. u/s 3 of UGC Act 1956)

SOMETIMES A STEP TOWARDS SUCCESS STARTS WITH AN OPPORTUNITY



ANTHE

AAKASH NATIONAL TALENT HUNT EXAM

GET UPTO
100% SCHOLARSHIP*

ANTHE gives students once-in-a-lifetime opportunity to build a successful career & get Scholarship* upto 100% and Cash Awards. Enroll in Aakash for Medical/ Engineering Entrance Exams preparation and to build a strong foundation for Olympiads & other Scholarship Exams. With the inception of ANTHE in 2010, we have groomed thousands of students, producing best results in the country. ANTHE has now become a gateway to success for students.

Date of Examination: **29th October, 2017 (Sunday)**

Exam Fee: ₹ 300/- |

Paytm
Pay Fees
on Paytm!



Last Date of receiving duly filled enrollment form by post / by hand / through online - **22nd October, 2017 (Sunday)**
and with late fee - **24th October, 2017 (Tuesday)**

Test Centres: **All Over India**

Download Enrollment Form from our website: www.aakash.ac.in
or www.anthe.in. The result shall be displayed on our website.



Aakash

Medical | IIT-JEE | Foundations

(Divisions of Aakash Educational Services Pvt. Ltd.)

Registered Office : Aakash Tower, 8, Pusa Road, New Delhi-110005. Ph.: (011) 47623456 | E-mail: anthe@aesl.in

SMS Aakash to 53030 | TOLLFREE: 1800-180-2727, 1800-102-2727 | Helpline: 39454545 | Give a Missed Call: 9599698643