iranchembook.ir/edu

**PRACTICE PAPERS 2018** 

India's #1 CHEMISTRY MONTHLY FOR JEE (Main & Advanced) & NEET www.mtg.in | March 2018 | Pages 92 | ₹ 30

6

MAIN ADVANCED NEET AIIMS CBSE BOARD

> Exam Special

# NEET | JEE Essentials Class XI-XII

CHEMISTRY MUSING



# ADVANCED CHEMISTRY BLOC

 1
 1
 1
 10
 01
 010
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10
 10</t

CONCEPT BOOSTER

CONCEPT

MAP



WWW.LPU.IN

# UNIVERSITY WITH UNBEATABLE PLACEMENT RECORD

chembook ir/edu

From Google to Microsoft, LPU Students are working with top brands at packages of over **₹1 crore.** 

# Think **BIG**

SONIA (LPU Alumni) Software Engineer amazon USA



Thousands of LPU students have got placed in top MNCs in USA, UK, Canada, Australia etc. LPU has been attracting 500+ multinational and national companies like Cognizant, Amazon, HP, Capegemini and many more.



Study engineering minors like, Data Engg., Cyber Security, Internet of Things (IoT)] or nonengineering minors like, Business Economics, Entrepreneurship along with your chosen branch.



Hands-on practical training and certifications in collaboration with world's top brands including Google, Intel, SAP, Oracle, Cisco, Microsoft, and Cadence.

## ADMISSIONS FOR 200+ PROGRAMMES

AFTER 10<sup>TH</sup>, 12<sup>TH</sup>, GRADUATION & POST GRADUATION IN Engineering | Computer Applications | Management | Economics | Law | Agriculture | Hotel Management & Tourism | Biotechnology |

Agriculture | Hotel Management & Tourism | Biotechnology | Pharmaceutical Sciences | Medical Laboratory Sciences | Physiotherapy | Food Technology | Microbiology | Biochemistry | Forensic Sciences | Architecture | Planning | Interior & Furniture Design | Product & Industrial Design | Nutrition and Dietetics | Botany | Zoology | Chemistry | Physics | Mathematics | Commerce | Fashion Design | Multimedia & Animation | Journalism & Film Production | Fine Arts |Performing Arts | Ayurvedic Pharmaceutical Sciences | Political Science | Geography | History | Psychology | Sociology | Public Administration | Library Science | Arts (Humanities) | English and Foreign Languages | Indian Languages | Education | Physical Education



#### HOW TO APPLY? :

Applicants can apply online at www.lpu.in Date of Examination: 10th-30th April 2018 LAST DATE OF APPLICATION : 31-03-2018

#### Toll Free : 1800-102-4430

LOVELY PROFESSIONAL UNIVERSITY (LPU) Jalandhar-Delhi G.T. Road, Phagwara, Punjab admissions@lpu.co.in, SMS LPU to 53030



Transforming Education Transforming India





# Inspired by Love Guided by Knowledge

Life skills and industry-oriented technical training increases your chance of success in campus placements

Bernett Orlando in Google Amrita Alumnus 2013-2017



Multiple Placement Opportunities in top-notch National & Multinational companies



Internship with stipend upto ₹1.0 Lakh p.m.



Highest Salary ₹25.16 Lakhs p.a.

OUR RECRUITERS ABB, Amazon, AstraZeneca, Blue Star, Bosch, Caterpillar, Cerner, Cisco, Cognizant, Dell, Directi, Ericsson, E&Y, FireEye, Ford, GE Digital, Google, Honeywell, Hyundai, IBM, Infosys, Intuit, Juniper, KLA Tencor, Larsen & Toubro, Mahindra & Mahindra, MicroFocus, Microsoft, National Instruments, Nationstar Mortgage, Oracle, Philips, Quess Corp, Reliance Power, Renault Nissan, Roadrunnr, Samsung, SAP, Seagate, TCS, Thermo Fisher, Toyota, United Technologies, Verizon, Wipro, Xome, Zoho to name a few....



AMRITA is RANKED 168 in Asia



www.topuniversities.com/university-rankings/asian-university-rankings/2018

How to

apply a

RANKED 9 in University category by NIRF, MHRD, Govt. of India www.nirfindia.org/UniversityRanking.html

# #1

## SWACHHTA RANKING 2017

**Ranked 'First'** among the cleanest Higher Education institutions in the country in the category of Technical Institutions

SCAN

CODF

www.amrita.edu/btech2018

# AEEE 2018

for Admission to B.Tech. Programmes

offered at Amritapuri, Bengaluru & Coimbatore campuses

Last Date for Receiving Application: 31 March, 2018

For Queries Amritapuri Campus 0476 2809 400 / 402 / 405 admissions@am.amrita.edu Bengaluru Campus 080 2518 3700 admissions@blr.amrita.edu Coimbatore Campus 0422 2685 169 / 170 admissions@amrita.edu

Amrita Vishwa Vidyapeetham, Amrita Nagar P.O., Ettimadai, Coimbatore - 641 112. Tamil Nadu | Accredited by NAAC with 'A' grade

iranchem	book.ir/e	du									
		An In	stitute	<b>For</b> l	Excell	ence In S	<b>Cience</b>	<b>)</b>			
<b>RESULT 2</b>	2017			:	4				FELICI	TATIO	N
	2014	1.11				TOPIALS	11.33				
1			-			000	-04	000		29	8
	<b>#</b> #					Sec.	COD		10-0		
B 3. 2.1		-	Qu	C'A	A	A Q	9.9		-	1	1
	3.40	300	62			FRE	Q+1				
		· •				INT	TI			XI	
REFO		Y		公理	RX.		44			10-14	E.
	Achieve	ment			ALCONTACT OF						
Academic Year	Acmeve	2014	2015	2016	2017	1	12	with			
Selections	Medical	80	98	115	139				1.1.1		
0010013	Engineering	60	76	94	109			-			
COURSE	S OFFE	RED					-				
Two Yea	ars Integr	ated	Progr	amm	e			-			
◆ NEET (Me	edical Entra	nce Ex	(am.)	、		F					
◆ III-JEE (E	ngineering	Entra	nce Ex	(am.)		11	Π				
One Yea	ar Integra	ted P	rogra	mme		F			Th a	<b>4</b> 0h	
NEET Rep	peater Batc	ו				Ш	-				
Test Se	ries NEET	JEE	(Online	/Offlir	ne)	ST	TATIT T	UTORIALS -			
<ul> <li>Special Or Test Serie</li> </ul>	Special Online Distance Learning / Test Series Programme										
<ul> <li>◆ Tab with study material</li> </ul>											
For Free Test Series-Download Lalit Eduapp.											
<ul> <li>Highly Experienced, Best Faculty Team.</li> <li>Well Equiped Digital Class Room.</li> </ul>											
Well Furnished Library & Laboratory.											
The Best Institute in Maharashtra for Medical & Engineering Entrance Exams											
Kalpande Sir's											

An Institute For Excellence In Science

Toshniwal Lay-out, Akola (Maharashtra) Phone : 0724-2456 654, 7720 857 857

### **Opportunity For Faculties**

Course	Faculties	Min.Exp.	Min.Package / Annum.
NEET	Senior	6 yr.	12 Lac
or JEE	Junior	3 yr.	6 Lac
Foundation	Senior	5 yr.	5 Lac
8 <sup>th</sup> ,9 <sup>th</sup> & 10 <sup>th</sup>	Junior	2 yr.	3 Lac

- ★ Special Incentives & Increments According to Performance
- ★ Special Reward to Faculty on Producing Best Ranks.

For Recruitment of Senior & Junior Faculties send your Resume to hrltmaharashtra@gmail.com

6

## iranchembook.ir/edu



www.srmuniv.ac.in

# ALL THAT MATTERS IS TALENT

At SRM we believe in honing potential, in helping students achieve their dreams.

We believe that nothing should come in the way if there is talent. Be it in academics, sports or in some other field. Which is why, we offer both full and partial scholarships to the deserving based on various criteria.

If you dare to dream and have the dedication to realise it, we can help you make it happen.

Scholarships offered to our students:

- SRM founder's scholarship
- Socio-economic scholarship
- SRM merit scholarship
- Differently-abled scholarship
- SRM arts and culture scholarship
  - Take the SRMJEEE (B.Tech) 2018

Online exam dates: April 16-30, 2018 Last date to apply 31<sup>st</sup> March 2018.

To apply, log onto www.srmuniv.ac.in | For queries: Call: +91 44 2745 5510 / +91 44 4743 7500

Join us at: f SRMUniversityOfficial 🛛 🖸 srmuniversityofficial 🛛 🕥 SRM\_Univ

# iranchembook.ir/edu

# **CHEMISTRY** adav

Volume 27

March 2018

**Managing Editor** Mahabir Singh Editor Anil Ahlawat

**Corporate Office:** 

Plot 99, Sector 44 Institutional area, Gurgaon -122 003 (HR). Tel: 0124-6601200 e-mail: info@mtg.in website: www.mtg.in **Regd. Office:** 

No. 3

406, Taj Apartment, Near Safdarjung Hospital, New Delhi - 110029.

S	
	_
Ζ	
1	
10.0	1
7	
-	•
C	

**Competition Edge** 

- **Chemistry Musing Problem Set 56** 8
  - **JEE Advanced Practice Paper** 10
    - **AIIMS Practice Paper** 21
    - Gear Up for JEE Main 31
      - **NEET Practice Paper** 36
        - **Concept Booster** 43
    - **Advanced Chemistry Bloc** 48
- **Chemistry Musing Solution Set 55** 50

#### Class 11

- **Concept Map** 46
- **NEET | JEE Essentials** 52

#### Class 12

- **Concept Map** 47
- **NEET | JEE Essentials** 64
- **CBSE Board Practice Paper** 76

#### Subscribe online at WWW.mtg.in

Individual	Coml	bined Sub	scription	Rates			
	1 yr.	2 yrs.	3 yrs.		1 yr.	2 yrs.	3 yrs.
Mathematics Today	330	600	775	PCM	900	1500	1900
Chemistry Today	330	600	775	PCB	900	1500	1900
Physics For You	330	600	775	PCMB	1000	1800	2300
Biology Today	330	600	775				

Send D.D/M.O in favour of MTG Learning Media (P) Ltd. Payments should be made directly to : MTG Learning Media (P) Ltd, Plot No. 99, Sector 44, Gurgaon - 122003 (Haryana)

We have not appointed any subscription agent

Printed and Published by Mahabir Singh on behalf of MTG Learning Media Pvt. Ltd. Printed at HT Media Ltd., B-2, Sector-63, Noida, UP-201307 and published at 406, Taj Apartment, Ring Road, Near Safdarjung Hospital, New Delhi - 110029.

Editor : Anil Ahlawat

Readers are adviced to make appropriate thorough enquiries before acting upon any advertisements published in this magazine. Focus/Infocus features are marketing incentives. MTG does not vouch or subscribe to the claims and representations made by advertisers. All disputes are subject to Delhi jurisdiction only. Copyright© MTG Learning Media (P) Ltd. All rights reserved. Reproduction in any form is prohibited.



# STRY MUSING

## PROBLEM **SET 56**

hemistry Musing was started from August '13 issue of Chemistry Today. The aim of Chemistry Musing is to augment the ✔ chances of bright students preparing for JEE (Main and Advanced) / NEET / AIIMS / JIPMER with additional study material. In every issue of Chemistry Today, 10 challenging problems are proposed in various topics of JEE (Main and Advanced) / NEET. The detailed solutions of these problems will be published in next issue of Chemistry Today. The readers who have solved five or more problems may send their solutions. The names of those who send atleast five correct solutions will be published in the next issue. We hope that our readers will enrich their problem solving skills through "Chemistry Musing" and stand in better stead while facing the competitive exams.

#### JEE MAIN/NEET

1. A hydrogen electrode placed in a buffer solution of NaCN and HCN in the ratio x : y and y : x has electrode potential values a and b volts respectively, at 25 °C. If the difference (a - b) is 35.52 mV, the ratio y: x is

(c) 3 (a) 2 (b) 3/2 (d) 2/3

2. The normal boiling points of four elements with consecutive atomic numbers are tabulated as :

Elements	W	X	Y	Ζ
Atomic number	N	N + 1	N+2	<i>N</i> + 3
Boiling point (°C)	58	-152	688	1380

One of the important ore of Z is celestine then, which of the following compounds would be the most stable?

(a)  $W_2X$ (b)  $W_2Y$  (c)  $W_2Z$ (d)  $XY_2$ 

- 3. An amine on treatment with  $HNO_2$  evolved  $N_2$  gas. The amine on exhaustive methylation with CH<sub>3</sub>I formed a quaternary salt containing 59.07% iodine. The amine is likely to be
  - (a) CH<sub>3</sub>NH<sub>2</sub> (b) (CH<sub>3</sub>)<sub>2</sub>NH

(c) 
$$C_2H_5NH_2$$
 (d)  $(CH_3)_3N$ 

4. For the given reaction,

$$HO-CH_2-CH_2-CH_2-CH_2CI \xrightarrow{OH} (A)$$

which of the following statements is true?

- (i) (A) is butane-1, 4-diol.
- (ii) (A) is tetrahydrofuran.
- (iii) (*A*) is prepared by  $S_N 1$  reaction.
- (iv) (A) is prepared by intramolecular  $S_N^2$  reaction.
- (a) (i) only (b) (ii) and (iv) only
- (c) (i) and (iv) only (d) (ii) and (iii) only
- 5. Ordinary strong solutions of HCl, HNO<sub>3</sub> and H<sub>2</sub>SO<sub>4</sub> contain roughly
  - (a) 1/5, 2/3 and 3/3 fractions of pure acid in water respectively
  - (b) 2/3, 1/5 and 3/3 fractions of pure acid in water respectively

- (c) 2/3, 3/3 and 1/5 fractions of pure acid in water respectively
- (d) none of these.

#### JEE ADVANCED

6. A given sample of milk turns sour at room temperature (20 °C) in 64 hours. In a refrigerator at 3 °C, milk can be stored three times as long before it sours. How long will it take milk to sour at 40 °C? (a) 192 hr (b) 64 hr (c) 20.5 hr (d) 30 hr

#### COMPREHENSION

For the given reactions sequence,





#### **INTEGER VALUE**

- 9. For how many of the following, the apparent weight increases by applying magnetic field? NO, NO<sub>2</sub>, O<sub>2</sub>, K<sub>3</sub>[Fe(CN)<sub>6</sub>], KO<sub>2</sub>, MnSO<sub>4</sub>, NiSO<sub>4</sub>, CuSO<sub>4</sub>, ZnSO<sub>4</sub>
- 10. During the electrolysis of conc.  $H_2SO_4$ , it was found that H<sub>2</sub>S<sub>2</sub>O<sub>8</sub> and O<sub>2</sub> were liberated in a molar ratio of 3 : 1. If moles of  $H_2$  in terms of moles of  $H_2S_2O_8$ is *a* : *b* then the value of  $3 \times a/b$  is



۰



# iranchembook.ir/edu



# PRACTICE PAPER 2018

#### PAPER - I

#### **SECTION 1 (MAXIMUM MARKS : 28)**

#### This section contains SEVEN questions.

- Each question has FOUR options (a), (b), (c) and (d). ONE OR MORE THAN ONE of these four options is (are) correct.
- For each question, darken the bubble(s) corresponding to all the correct option(s) in the ORS.
- For each question, marks will be awarded in one of the following categories :

Full Marks : +4	If only the bubble(s) corresponding to all the correct option(s) is(are) to be a set of the correct option (s) is (are)
	darkened.
Partial Marks : +	For darkening a bubble corresponding
	to each correct option, provided NO
	incorrect option is darkened.
Zero Mark :	) If none of the bubbles is darkened.
Negative Marks : -2	2 In all other cases.

- For example, if (a), (c), and (d) are all the correct options for a question, darkening all these three will get +4 marks; darkening only (a) and (d) will get +2 marks; and darkening (a) and (b) will get-2 marks, as a wrong option is also darkened.
- **1.** A mixture of two inorganic salts gives following chemical reactivity :
  - (i) Mixture on reaction with dilute H<sub>2</sub>SO<sub>4</sub> produces a colourless and unpleasant gas which turns acidified potassium dichromate paper green.
  - (ii) Mixture on reaction with concentrated H<sub>2</sub>SO<sub>4</sub> gives reddish brown gas which does not produce orange red spots on starch paper.
  - (iii) The mixture gives white precipitate with barium chloride solution which is soluble in dilute HCl.
  - (iv) The sodium carbonate extract of mixture responds to brown ring test.

The mixture contains

- (a)  $SO_3^{2-}$  and  $NO_3^{-}$  anions
- (b)  $S^{2-}$  and  $NO_3^-$  anions
- (c)  $SO_3^{2-}$  and  $NO_2^{-}$  anions
- (d)  $SO_3^{2-}$  and  $Br^-$  anions.
- 2. From the following data, mark the option(s) where  $\Delta H$  is correctly written for the given reaction.

Given :  $H^+_{(aq)} + OH^-_{(aq)} \longrightarrow H_2O_{(l)}$ ;  $\Delta H = -57.3 \text{ kJ/mol}$  $\Delta H_{\text{solution}}$  of  $HA_{(g)} = -70.7 \text{ kJ/mol}$  $\Delta H_{\text{solution}}$  of  $BOH_{(g)} = 20 \text{ kJ/mol}$ 

 $\Delta H_{\text{ionisation}}$  of HA = 15 kJ/mol and BOH is a strong base.

#### Reaction

## $\Delta_r H(\text{kJ/mol})$

- (a)  $HA_{(aq)} + BOH_{(aq)} \longrightarrow BA_{(aq)} + H_2O 42.3$
- (b)  $HA_{(g)} + BOH_{(g)} \longrightarrow BA_{(aq)} + H_2O$  -93
- (c)  $HA_{(g)} \longrightarrow H^+_{(aq)} + A^-_{(aq)}$  -55.7
- (d)  $BOH_{(g)} \longrightarrow B^+_{(aq)} + OH^-_{(aq)} -20$
- **3.** The hemiacetal form of glucose is indicated by
  - (a) reaction with  $(CH_3CO)_2O$
  - (b) oxidation with Tollens' reagent
  - (c) reduction with HI/P
  - (d) glycoside formation.
- $\text{HeCOOEt} \xrightarrow{\text{(i) } \text{NH}_2\text{NH}_2}_{\text{(ii) } \text{HNO}_2} (B) \xrightarrow{\text{MeOH}} (C)$

Which of the following statements are correct about the given reactions sequence?

- (a) The compounds (*B*) and (*C*), respectively, are Me N = C = O and MeNHCOOMe.
- (b) The compounds (*B*) and (*C*), respectively, are  $Et N \equiv C \equiv O$  and MeNH<sub>2</sub>.
- (c) The reaction proceeds via the formation of acyl nitrene (MeCON:) as the intermediate species.
- (d) The reaction proceeds via the formation of acyl nitrene (EtCON:) as the intermediate species.
- **5.** Which of the following substrates will give rearranged product in hydration reaction?

(a) 
$$CH_3 - CH - CH = CH_2$$
  
 $\downarrow$   
 $CH_3$   
(b)  $CH_3 - \ddot{S} - CH_2 - CH = CH_2$ 





So what are you waiting for? Order MTG's JEE Advanced Explorer today.

Scan now with your smartphone or tablet Application to read QR codes required

Available at all leading book shops throughout India. To buy online visit www.mtg.in. For more information or for help in placing your order, call 0124-6601200 or email:info@mtg.in

(c) 
$$CH_3 - CH = CH_2$$
  
(d)  $Ph$   $C = CH - CH_3$   
 $Ph$   $C = CH_3$   
 $CH_3$   
 $C$ 

- 6. Identify compound(s) in which gauche conformer is more stable than staggered.
  - (a) 1, 2-Difluoroethane (b) Chloropropane
  - (c) Ethylene glycol (d) Succinic acid
- 7. Which among the following statements is/are correct? (a)  $XeF_4$  and  $SbF_5$  combine to form salt.
  - (b) He and Ne do not form clathrates.
  - (c) He has lowest b.pt. in its group.
  - (d) He diffuses through rubber and polyvinyl chloride.

#### **SECTION 2 (MAXIMUM MARKS : 15)**

- This section contains FIVE questions.
- The answer to each question is a SINGLE DIGIT INTEGER ranging from 0 to 9, both inclusive.
- For each question, darken the bubble corresponding the correct integer in the ORS.
- For each question, marks will be awarded in one of the following categories :

Full Marks: +3 If only the bubble corresponding to the correct answer is darkened. Zero Marks: 0 In all other cases.

- 8. Number of chelate rings in  $[Cr(ox)_3]^{3-} = a$ Number of chelate rings in  $[Co(en)(NH_3)_2(py)_2]^{3+} = b$ Number of chelate rings in  $[Fe(EDTA)]^{-} = c$ Number of chelate rings in brown ring complex  $[Fe(H_2O)_5(NO)]SO_4 = d$ then the value of (a + b + c + d) is
- 9. If edge fraction unoccupied in ideal antifluorite structure is x. Calculate the value of Zv

where, 
$$Z = \frac{x}{0.097}$$

- 10. *m*-Phenylenediamine when treated with NaNO<sub>2</sub> and HCl at 0-5 °C undergoes self-coupling reaction to produce a trimer, known as Bismarck brown. The total number of nitrogen atoms in the final product is
- 11. A decapeptide (mol. wt. 796) on complete hydrolysis gives glycine, alanine and phenylalanine. Glycine contributes 47.0% to the total weight of the hydrolysed products. The number of glycine units present in the decapeptide is
- 12. When (S)-2-chloropentane reacts with NaOH in 75 per cent ethanol and 25 per cent acetone follows rate law :

What will be percentage of S<sub>N</sub>1 product when concentration of  $[OH^-] = 1.5$  molar?

#### **SECTION 3 (MAXIMUM MARKS : 18)**

- This section contains SIX questions of matching type.
- This section contains TWO tables (each having 3 columns and 4 rows)
- Based on each table, there are THREE questions
- Each question has FOUR options (a), (b), (c) and (d). ONLY ONE of these four options is correct
- For each question, darken the bubble corresponding to the correct option in the ORS
- For each question, marks will be awarded in one of the following categories :
  - Full Marks : +3 If only the bubble corresponding to the correct option is darkened. Zero Marks :
  - 0 If none of the bubbles is darkened Negative Marks : -1 In all other cases.

#### Answer Q. 13 to 15 by appropriately matching the information given in the three columns of the following table.

Columns 1, 2 and 3 contain number of moles, volumes at STP and number of atoms, respectively.

Column 1	Column 2	Column 3
(I) 0.125	(i) 7840 mL	(P) $13.25 \times 10^{23}$
(II) 0.35	(ii) 11.2 dm <sup>3</sup>	(Q) $6.32 \times 10^{23}$
(III) 1.1	(iii) 2.8 L	(R) $6.023 \times 10^{23}$
(IV) 0.50	(iv) 24640 cc <sup>3</sup>	(S) $3.011 \times 10^{23}$

- **13.** The correct combination for  $16 \text{ g of } NO_2$  is (a) (I) (ii) (R) (b) (II) (iii) (S) (c) (II) (i) (Q) (d) (IV) (iii) (P)
- 14. Oxygen gas occupies 10.5 L at 27 °C under a pressure of 2.5 atm. At STP correct match for the sample will be

(a)	(IV) (ii) (R)	(b) (III) (iv) (P)
(c)	(II) (i) (Q)	(d) (I) (iii) (S)

15. For 32 g of sulphur, correct combination will be (a) (I) (ii) (P) (b) (I) (iii) (R) (c) (II) (iii) (Q) (d) (III) (iv) (S)

Answer Q. 16 to 18 by appropriately matching the information given in the three columns of the following table.





Columns 1, 2 and 3 contain reactants, intermediates and products, respectively.

Column 1	Column 2	Column 3
(I) $\stackrel{R}{H} \subset = O$	(i) γ-Hydroxy acid	$(P) \xrightarrow{R}_{R} NH$
(II) RCONH <sub>2</sub>	(ii) Hemiacetal	(Q) 1°-amine
(III) CHO CHO	(iii) Alkyl isocyanate	$(R) \stackrel{R}{\underset{H}{\searrow}} C \stackrel{OR'}{\underset{OR'}{\bigvee}}$
$(IV) \\ 0 \\ R - C - N \begin{pmatrix} CH_3 \\ CH_3 \end{pmatrix}$	(iv) Carbene	(S) Phthalide

#### **16.** Correct matching for addition of R' – OH will be

(a)	(1) (iii) $(S)$	(b) (l) (ii) (R)
(c)	(III) (ii) (P)	(d) (III) (i) (Q)

17. Correct matching for intramolecular Cannizzaro reaction will be (a) (III) (ii) (P) (b) (IV) (iii) (R) (c) (III) (i) (S)

(d) (I) (iv) (Q)

- 18. Correct matching for Hoffmann bromamide degradation reaction is
  - (a) (IV) (iii) (Q)
  - (b) (IV) (iv) (P) (c) (II) (i) (R)
  - (d) (II) (iii) (Q)

PAPER - II

1/2

#### SECTION 1 (Maximum Marks : 21) This section contains SEVEN questions.

- Each question has FOUR options (a), (b), (c) and (d). ONLY ONE
- of these four options is correct.
- For each question, darken the bubble corresponding to the correct option in the ORS.
- For each question, marks will be awarded in one of the following categories :

Full Marks : +3 If only the bubble corresponding to the correct option is darkened. Zero Marks : 0 If none of the bubbles is darkened. Negative Marks : -1 In all other cases.

1. For a 3s-orbital, 
$$\psi(3s) = \frac{1}{9\sqrt{3}} \left(\frac{1}{a_0}\right)^{1/2}$$
  
(6 - 6\sigma + \sigma^2) e^{-\sigma/2}

where,  $\sigma = \frac{2Zr}{3a_0}$ . What is the maximum radial distance of node from nucleus?

(a)  $\frac{2}{3} \frac{(3+\sqrt{3})a_0}{Z}$  (b)  $\frac{3}{2} \frac{(3+\sqrt{3})a_0}{Z}$ (c)  $\frac{3}{2} \frac{(3-3\sqrt{3})a_0}{Z}$  (d)  $\frac{3}{2} \frac{(3-\sqrt{3})a_0}{Z}$ 

2. In first order reaction,  $A \longrightarrow B$ , A is optically active and *B* is optically inactive. A series of experiments were conducted on a solution of A :

Time	0	60 min	∞
Optical rotation	82°	77°	2°

Assume some impurity is present, calculate the optical rotation after 5 hours.

(Given ln 1.066 =	$0.064, e^{0.32} = 1.377)$
(a) 60	(b) 30
(c) 20	(d) 120

3. A fixed mass 'm' of a gas is subjected to transformation of states from K to L to M to N and back to *K* as shown in the figure :



The pair of isochoric processes among the transformation of states is

(a) K to L and L to M (b) L to M and N to K

(c) L to M and M to N(d) M to N and N to K.

- Devise a series of reactions to convert 4. ethyl 3-oxobutanoate to ethyl 4-oxopentanoate. Select reagents and conditions from the following table, listing them in the order of use :
  - (1) Sodium ethoxide in ethanol
  - (2) Ethanol + Acid catalyst
  - (3)  $H_3O^+$ ; heat (4)  $CO_2$  then  $H_3O^+$
  - (5) Mg in ether (6)  $PBr_3$
  - (7) NaBH<sub>4</sub> in alcohol (8) CH<sub>2</sub>I<sub>2</sub> in ether; Zn Cu
  - (9)  $BrCH_2COOC_2H_5$  (10)  $(CH_3CO)_2O$ ; Pyridine
  - (a) 1, 9, 3 then 2 (b) 7, 6, 5, 10 then 2
  - (c) 3, 7, 6, 5, 10 then 2 (d) 8, 3 then 2

- 5. FeO crystal has a simple cubic structure and each edge of the unit cell is 5 Å. Taking density of the oxide as 4 g/cc, the number of  $Fe^{2+}$  and  $O^{2-}$  ions present in each unit cell are

  - (a)  $4Fe^{2+}$  and  $4O^{2-}$  (b)  $6Fe^{2+}$  and  $6O^{2-}$ (c)  $2Fe^{2+}$  and  $2O^{2-}$  (d)  $1Fe^{2+}$  and  $1O^{2-}$
- 6. The bond angle between two hybrid orbitals is 180°. The percentage s-character of hybrid orbital is
  - (a) 50 (b) 75
  - (c) 33 (d) 66
- 7. What is the complementary RNA sequence for the **DNA segment AATCAGTT?** 
  - (a) AAUCAGUU (b) CCAUCGAA
  - (c) AACUGAUU (d) UUAGUCAA

#### **SECTION 2 (MAXIMUM MARKS : 28)**

- This section contains SEVEN questions.
- Each question has FOUR options (a), (b), (c) and (d). ONE OR MORE THAN ONE of these four option(s) is(are) correct.
- For each question, darken the bubble(s) corresponding to all the correct option(s) in the ORS.
- For each question, marks will be awarded in one of the following categories :

ull Marks :	+ 4	If only the bubble(s) corresponding
		to all the correct option(s) is(are)
		darkened.
Partial Marks :	+1	For darkening a bubble corresponding
		to each correct option, provided NO
		incorrect option is darkened.
ero Marks :	0	If none of the bubbles is darkened.

Negative Marks : -2 In all other cases.

- For example, if (a), (c), and (d) are all the correct options for a question, darkening all these three will get +4 marks; darkening only (a) and (d) will get +2 marks; and darkening (a) and (b) will get -2 marks, as a wrong option is also darkened.
- Which statements are correct about lactose? 8.
  - (a)  $(C_1 \beta)$  (OH) of glucose is linked with  $(C_4 - OH)$  of galactose.
  - (b)  $(C_1 \beta)$  (OH) of galactose is linked with (C<sub>4</sub>—OH) of  $\beta$ -glucose.
  - (c) It is hydrolysed both by amylase and lactase.
  - (d) It exhibits mutarotation.
- 9. Which of the following would give Hoffmann alkene?





- **10.** Select the correct statement(s).
  - (a) On Pauling scale, the difference in EN of two atoms A and B in SI units is

$$(EN_A - EN_B) = 0.1017 \sqrt{\Delta_{A-B}}$$

- (b) On Pauling scale, the difference in EN of two atoms A and B in kcal  $mol^{-1}$  is  $(EN_A - EN_B) = 0.208 \sqrt{\Delta_{A-B}}$
- (c) The Mulliken's EN values are scaled down to match the Pauling value by dividing  $\left(\frac{IP + EA}{2}\right)$ in eV by 2.8
- (d) The Mulliken's EN values are scaled down to match the Pauling value by multiplying  $\left(\frac{IP+EA}{2}\right)$  in eV by 3.17

$$11. \qquad O \qquad \xrightarrow{\text{dil. H}_2\text{SO}_4} P_1 + P_2$$

- $P_1$  and  $P_2$  products are identified by
- (a) Tollens' reagent (b) Iodoform test
- (c)  $Br_2 + H_2O$  test (d) 1% of alkaline KMnO<sub>4</sub>.
- **12.** Mark the incorrect statement(s).
  - (a) Potassium dichromate oxidises a secondary alcohol into a ketone.
  - (b) Potassium permanganate is a weaker oxidising agent than potassium dichromate.
  - (c) Potassium dichromate oxidises a secondary alcohol into aldehyde.
  - (d) Alkaline KMnO<sub>4</sub> solution oxidises tertiary alcohol to a mixture of a ketone and an acid.
- **13.** Select the correct statement(s).
  - (a) Chelation effect is maximum for five or six membered rings.
  - (b) Complex ions in which ligands can be interchanged rapidly are said to be non-labile.
  - (c) For a given ion and ligand, greater the charge on the metal ion, greater is the stability.
  - (d)  $[Fe(CN)_6]^{4-}$  and  $[Fe(H_2O)_6]^{2+}$  can be distinguished by magnetic moment.



F

F

## iranchembook.ir/edu

- 14. Which of the following statements is/are correct?
  - (a) Alcoholic group is tested by CAN test.
  - (b) Accurate method to determine the molecular weight of organic compound is 'cryoscopic method'.
  - (c) If there is 54% silver in a silver salt of a dibasic acid, then its (dibasic acid) molar mass is  $186 \text{ g mol}^{-1}$ .
  - (d) Diazonium salts form azo dyes with alkaline compounds.

#### SECTION 3 (Maximum Marks : 12)

- This section contains TWO paragraphs.
- Based on each paragraph, there are TWO questions.
- Each question has Four options (a), (b), (c) and (d). ONLY ONE of these four options is correct.
- For each question, darken the bubble corresponding to the correct option in the ORS.
- For each question, marks will be awarded in one of the following categories :
  - **Full Marks :** +3 If only the bubble corresponding to the correct option is darkened.

Zero Marks: 0 In all other cases.

#### Paragraph 1

Tollens' test is given by aldehydes.

$$Ag^{+} + e^{-} \longrightarrow Ag; \qquad E_{red}^{\circ} = +0.800 \text{ V}$$

$$C_{6}H_{12}O_{6} + H_{2}O \longrightarrow C_{6}H_{12}O_{7} + 2H^{+} + 2e^{-}$$

$$E_{ox}^{\circ} = -0.05 \text{ V}$$

$$[Ag(NH_{3})_{2}]^{+} + e^{-} \longrightarrow Ag + 2NH_{3} \quad E_{red}^{\circ} = +0.373 \text{ V}$$

$$use\left(\frac{F}{RT}\right) = 38.9 \text{ V}^{-1}$$

**15.** Calculate (ln *K*) for,

$C_6H_{12}O_6 + 2Ag^+ + H_2O$	$\longrightarrow C_6H_{12}O_7 + 2H^+ + 2Ag$
(a) 55.6	(b) 29.6
(c) 66	(d) 58.35

**16.** NH<sub>3</sub> is used in this reaction rather than any other base. What is the correct reason for this?

- (a)  $[Ag(NH_3)_2]^+$  is a weaker oxidising agent than  $Ag^+$ .
- (b) NH<sub>3</sub> prevents the decomposition of gluconic acid.
- (c) Ag precipitates gluconic acid as its silver salt.
- (d) NH<sub>3</sub> changes the standard reduction potential of [Ag(NH<sub>3</sub>)<sub>2</sub>]<sup>+</sup>.

#### Paragraph 2

Shown below is a synthesis of elm bark beetle pheromone, multistriatin.



Based on the above synthesis, answer the following questions :

- **17.** Tosylation is done in conversion of *A* to *B* in order to
  - (a) easily remove the -OH group of A
  - (b) reduce the -OH group
  - (c) oxidise the -OH group
  - (d) condense alcohols.
- **18.** Structural formula of (*C*) is



#### SOLUTIONS

PAPER - I

1. 
$$(\mathbf{a,c}):$$
 (i) Na<sub>2</sub>SO<sub>3</sub> + H<sub>2</sub>SO<sub>4</sub>  $\longrightarrow$  Na<sub>2</sub>SO<sub>4</sub> + H<sub>2</sub>O  
+ SO<sub>2</sub>↑  
K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub> + H<sub>2</sub>SO<sub>4</sub> + 3SO<sub>2</sub>  $\longrightarrow$  K<sub>2</sub>SO<sub>4</sub>  
+ Cr<sub>2</sub>(SO<sub>4</sub>)<sub>3</sub> + H<sub>2</sub>O  
(Green)  
(ii)  $4NO_2^- + 2H_2SO_4 \longrightarrow 4NO_2^+ + O_2^+ + SO_4^- + 2H_2O$ 

As it does not produce orange red spots on starch paper, it cannot be  $Br_2$ . So, the anion is  $NO_3^-$ .

(iii) 
$$SO_3^{2^-} + Ba^{2^+} \longrightarrow BaSO_3 \downarrow$$
 (white)  
 $SO_3^{2^-} + 2H^+ \longrightarrow SO_2 \uparrow + H_2O$   
(iv)  $2NO_3^- + 4H_2SO_4 + 6Fe^{2^+} \longrightarrow 6Fe^{3^+} + 2NO \uparrow$   
 $+ 4SO_4^{2^-} + 4H_2O$   
 $Fe^{2^+} + 5H_2O + NO \longrightarrow [Fe(H_2O)_5NO]^{2^+}$ 

CHEMISTRY TODAY | MARCH '18

15

2. (a, b, c) : (a) 
$$\Delta_r H = 15 - 57.3 = -42.3 \text{ kJ/mol}$$
  
(b)  $HA_{(g)} + BOH_{(g)} \xrightarrow{\Delta_r H} BA_{(aq)} + H_2O$   
 $-70.7 \text{ kJ} \qquad 20 \text{ kJ} \qquad -42.3 \text{ kJ}$   
 $\Delta_r H = -70.7 + 20 - 42.3 = -93 \text{ kJ/mol}$   
(c)  $HA_{(g)} \xrightarrow{\Delta_r H} H^+_{(aq)} + A^-_{(aq)}$   
 $-70.7 \qquad 15 \qquad -70.7 + 15 = -55.7 \text{ kJ/mol}$   
(d)  $BOH_{(g)} \longrightarrow BOH_{(aq)}; \Delta H = 20 \text{ kJ/mol}$   
 $BOH_{(aq)} \longrightarrow B^+_{(aq)} + OH^-_{(aq)}; \Delta H = -57.3 \text{ kJ/mol}$   
Total  $\Delta_r H = 20 - 57.3 = -37.3 \text{ kJ/mol}$   
3. (a, d) : Glucose  $\xrightarrow{\text{anhydride, pyridine}} Pentaacetate derivative Form oxime  $\sqrt{NH_2OH}$   
No reaction$ 

Glycoside is functionally acetal which is formed from hemiacetal.

(a,c) : It is an example of Lossen rearrangement **4**. reaction.



Delocalisation of bond-pair e of C-H bond in parallel ABMO of C-F is possible; leads to stability in gauche form



Due to +*I* nature of  $-CH_3$ ,  $\delta$ + created on it while -*I* nature of -Cl induce  $\delta$ - on it. Thus, dipole-dipole interaction leads to stability of gauche conformation.



7. (a, b, c, d)

8. (9):  $a = 3, b = 1, c = 5, d = 0 \Longrightarrow (a + b + c + d) = 9$ 

9. (3): Fraction of edge unoccupied (x) = 
$$\frac{a-2r}{a}$$

$$a = 2\sqrt{2}r , \ x = \frac{2(\sqrt{2}-1)}{2\sqrt{2}} = \frac{0.414}{1.414} = 0.293$$
$$Z = \frac{x}{0.097} = \frac{0.293}{0.097} = 3$$

10. (8):





 $0.125 \text{ mol} \equiv 0.125 \times 22.4 \text{ L} = 2.8 \text{ L}$  $0.125 \text{ mol contains} = 0.125 \times 6.023 \times 10^{23} \text{ molecules}$ 0.125 mol of S<sub>8</sub> contains =  $0.125 \times 6.023 \times 10^{23} \times 8$ =  $6.023 \times 10^{23}$  atoms **16.** (b):  $\underset{H}{\overset{R}{\longrightarrow}} C = O + H^+ \longrightarrow$  $\begin{bmatrix} R & f \\ H & C = O - H & \longleftrightarrow & R & f \\ H & C & O - H & \longleftrightarrow & H & C - OH \end{bmatrix}$  $\xrightarrow{R'OH} \stackrel{R}{\underset{H}{\overset{}}} C \xrightarrow{OH} \stackrel{R}{\underset{H}{\overset{}}} C \xrightarrow{OH} \stackrel{H^+}{\underset{H}{\overset{}}} H \xrightarrow{R} C \xrightarrow{OH} \stackrel{H^+}{\underset{Hemiacetal}{\overset{}}} H$  $\begin{array}{c} R \\ H \\ C \\ OR' \\ OR' \\ OR' \\ Acetal \end{array} \xrightarrow{R \\ OR' \\ Acetal \end{array} \xrightarrow{OR'} C \\ C \\ OR' \\ Acetal \end{array}$ MtG **ONLINE TEST SERIES** Practice Part Syllabus/ Full Syllabus 40 Mock Tests for E Main Now on your android Smart phones with the same login of web portal. Log on to test.pcmbtoday.com



- 1. (b): At nodes, probability of finding of electron is zero.
  - $\therefore \quad \psi^2 = 0 \text{ or } \psi = 0$

$$\therefore \quad 6-6 \ \sigma+\sigma^2=0; \ \sigma=3\pm\sqrt{3}$$

For maximum distance  $r = \frac{3}{2} \frac{(3 + \sqrt{3})a_0}{Z}$ 

CHEMISTRY TODAY | MARCH '18

2. (a): 
$$kt = \ln \frac{\alpha_0}{\alpha_t}$$
  
 $k \times 60 = \ln \frac{80}{75} \implies k = \frac{1}{60} \ln \frac{80}{75}$   
 $k \times 5 \times 60 = \ln \frac{80}{r_t}; 5\ln \frac{80}{75} = \ln \frac{80}{r_t}; \ln \frac{80}{r_t} = 0.32$   
 $r_t = 58$   
Optical rotation observed after 5 hrs = 58 + 2 = 60  
3. (b): In isochoric process, volume is constant.  
4. (a):  $CH_3CCH_2COC_2H_5 \xrightarrow{CH_3CH_2OH_3} \xrightarrow{CH_3CH_2OC_2H_5} \xrightarrow{CH_3CH_2OO}_{(1)}$   
 $CH_3-C \xrightarrow{C}CH^- \xrightarrow{BrCH_2COOC_2H_5} \xrightarrow{H_3O^+, heat}$   
 $CH_3CH_2OC \xrightarrow{(0)} \xrightarrow{$ 

4

Hence, there are four  $Fe^{2+}$  and four  $O^{2-}$  ions in each unit cell.

(a): The bond angle 180° means *sp*-hybridisation 6. which has 50% s-character.





Which is why the MTG team created 40 Years Chapterwise Solutions. The most comprehensive 'real' question bank out there, complete with detailed solutions by experts. An invaluable aid in your quest for success in JEE. Visit www.mtg.in to order online. Or simply scan the QR code to check for current offers.

Note: 40 Years Chapterwise Solutions are also available for each subject separately.



Scan now with your smartphone or tablet

Application to read QR codes required

Available at all leading book shops throughout India. To buy online visit www.mtg.in.

For more information or for help in placing your order, call 0124-6601200 or e-mail info@mtg.in

- (d): Complimentary DNA sequence : TTAGTCAA Complimentary RNA sequence : UUAGUCAA in DNA bases are bonded in complementary pair as A=T and G≡C while in RNA as A=U and G≡C.
- (b,d) : Amylase hydrolyses α-linkage. It exhibits mutarotation, since in β-D-glucose C-1 (OH) group is hemiacetalic.
- 9. (a, c) :

(a) The oxidation of 3° amine to amine oxide followed by Cope reaction on heating gives Hoffmann alkene (less substituted).



(b) will not give Hoffmann alkene. Benzylic H atom is more acidic due to (-I) effect of Ph.

- (c) will give Hoffmann alkene  $CH_2 = CH_2$ .
- (d) will not give Hoffmann alkene (Cope reaction).

#### 10. (a, b, c)



 (b, c, d): In acidic medium, KMnO<sub>4</sub> gives 5 oxygen while acidic K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub> gives 3 oxygen and secondary alcohols are oxidised to ketones.

Tertiary alcohols are resistance to oxidation in neutral or alkaline  $KMnO_4$  solution but are readily oxidised in acidic solution ( $KMnO_4/H_2SO_4$ ) to a mixture of a ketone and an acid each contains lesser number of carbon atoms than the original alcohol.

**13.** (a, c, d) : Complex ions in which ligands can be interchanged rapidly are said to be labile.

15. (d) : 
$$E_{cell}^{\circ} = E_{R}^{\circ} - E_{L}^{\circ} = 0.8 - 0.05 = 0.75 \text{ V}$$
  
 $\Delta G_{cell}^{\circ} = -2FE_{cell}^{\circ} = -RT\ln K$   
 $\ln K = \frac{2F}{RT} \times 0.75 = 2 \times 38.9 \times 0.75 = 58.35$ 

**16.** (a) : Because  $E_{Ag^+|Ag}^{\circ} > E_{[Ag(NH_3)_2]^+|Ag}^{\circ}$ 

17. (a): - OTs is a good leaving group *i.e.* better than -OH group.



. . . . . . 🗞 🚸

# Your favourite MTG Books/Magazines available in **PUNJAB at**

- Sunder Book Depot Amritsar
  - Ph: 0183-2544491, 2224884, 5059033; Mob: 9814074241
- Malhotra Book Depot Amritsar Mob: 8000300086, 9646537157, 9888733844
- Navchatan Book Depot Barnala
   Mob: 97790220692, 9417906880, 9779090135, 9876147263, 9779050692
- Mehta Book Centre Bathinda Mob: 9876029048, 9464158497
- Goyal Traders Book Sellers Bathinda
   Ph: 0164-2239632: Mob: 9417924911 9814485520
- Aggarwal Book Centre Bathinda Ph: 0164-2236042; Mob: 9417816439
- S M Enterprises Bathinda Ph: 0164-2240450; Mob: 7508622881, 9417363362
- Hans Raj Dogra & Sons Gurdaspur Mob: 9872032357
- Gupta Technical & Computer Book Shop Jalandhar
- Ph: 0181-2200397; Mob: 9915001916, 9779981081
- Cheap Book Store Jalandhar Mob: 9872223458
- City Book Shop Jalandhar Ph: 0181-2620800; Mob: 9417440753
- Deepak Book Distributiors Jalandhar
- Ph: 0181-2222131; Mob: 8528391133, 9872328131 Kiran Book Shop - Jalandhar Mob: 9876631526, 9779223883, 9872377808
- Amit Book Depot Ludhiana
- Ph: 0161-5022930, 5022930; Mob: 9815323429, 9815807871
- Bhatia Book Centre Ludhiana Ph: 0161-2747713; Mob: 9815277131
- Chabra Book Depot Ludhiana Ph: 0161-6900900, 2405427; Mob: 9501080070
- Gupta Book World Ludhiana
- Ph: 0161-2446870, 2409097, 3942992; Mob: 9463027555
- Khanna Book Depot Nabha Ph: 01765-220095; Mob: 9814093193, 9814309320
   Goel & Sons Book Depot Patiala Ph: 0172-2213643, 2202045; Mob: 9914096991
- Adarsh Pustak Bhandar Patiala
- Ph: 0175-2311430; Mob: 9814347613, 9815651737
- Pepsu Book Depot Patiala Ph: 0175-2214696; Mob: 9814041623, 9914903305

Visit **"MTG IN YOUR CITY"** on www.mtg.in to locate nearest book seller OR write to **info@mtg.in** OR call **0124-6601200** for further assistance.



# PRACTICE PAPER

- 1. Which of the following statements is incorrect?
  - (a) The trivial names of organic compounds are called common names.
  - (b) The systematic names of organic compounds are obtained from the IUPAC system.
  - (c) The systematic names of alkanes are based on the number of C atoms in the longest continuous chain of C atoms.
  - (d) The IUPAC name of  $(CH_3)_2CHCH(CH_3)_2$  is 1,3-dimethylhexane.
- 2. If a compound, on analysis was found to contain C = 18.5%, H = 1.55%, Cl = 55.04% and O = 24.81%, then its empirical formula is
  - (a)  $C_2H_2OCl$ (b) CH<sub>2</sub>ClO<sub>2</sub>

(c) CHClO (d)  $ClCH_2O$ 

3. Identify x, y and z for the following metallurgical process :

Metal sulphide  $\xrightarrow{x}$  Metal oxide  $\xrightarrow{y}$ Impure metal  $\xrightarrow{z}$  Pure metal.

*x*, *y* and *z* are respectively

- (a) roasting, smelting, electrolysis
- (b) roasting, calcination, smelting
- (c) calcination, auto-reduction, bassemerisation
- (d) none of the above is correct.
- 4.  $Ca^{2+}$  and  $F^{-}$  ions are located in  $CaF_2$  crystal respectively at face centred cubic lattice points and in
  - (a) tetrahedral voids
  - (b) half of tetrahedral voids
  - (c) octahedral voids
  - (d) half of octahedral voids.
- 5. Which one of the following statements regarding photochemical smog is not correct?
  - (a) Photochemical smog is formed by the combination of smoke, dust and fog containing sulphur dioxide from polluted air.

(b) Photochemical smog causes irritation in eyes and throat.

:xam on Mav 2018

- (c) Carbon monoxide does not play any role in photochemical smog formation.
- (d) Photochemical smog is oxidising in nature.
- 6. In ice, oxygen atom is surrounded
  - (a) tetrahedrally by 4 hydrogen atoms
  - (b) octahedrally by 2 oxygen and 4 hydrogen atoms
  - (c) tetrahedrally by 2 hydrogen and 2 oxygen atoms
  - (d) octahedrally by 6 hydrogen atoms.
- Freshly obtained SnO<sub>2</sub> is added to water containing 7. a little HCl. The sol obtained would be \_\_\_\_\_ due to preferential adsorption of \_ ions.
  - (a) positively charged, H<sup>+</sup>
  - (b) negatively charged,  $SnO_3^{2-}$
  - (c) positively charged, Sn<sup>4+</sup>
  - (d) positively charged,  $Sn^{2+}$
- Non-stick cookwares generally have a coating of a 8. polymer, whose monomer is
  - (a)  $CH_2 = CH_2$ (b)  $CH_2 = CHCN$
  - (d)  $CF_2 = CF_2$ (c)  $CH_2 = CHCl$
- The correct order of electron gain enthalpy values  $(\Delta_{eg}H)$  of the halogen atoms is

(a) 
$$F < Cl < Br < I$$
 (b)  $I < Br < F < Cl$ 

(c) 
$$I < Br < Cl < F$$
 (d)  $Cl < Br < I < F$ 

10. 
$$H_3 \overset{+}{N}_{Z} \overset{-}{\underset{COOH}{\bigvee}} \overset{-}{N} \overset{-}{H}_3$$

Arrange X, Y and Z in order of decreasing acidic strength.

- (a) X > Z > Y(b) Z < X < Y(c) X > Y > Z(d) Z > X > Y
- 11. If  $K_{sp(AgCNS)} = 1 \times 10^{-12}$  and  $K_{sp(AgBr)} = 5 \times 10^{-13}$ , then the values of simultaneous solubility of AgCNS and AgBr in a solution of water will be

- (a)  $8.16 \times 10^{-7}$ ,  $4.08 \times 10^{-7}$ (b)  $4.08 \times 10^{-7}$ ,  $8.16 \times 10^{-7}$

- (c) 8.16, 4.08 (d)  $1 \times 10^{-12}$ ,  $5 \times 10^{-13}$
- 12. Arrange the following compounds in order of increasing dipole moment :
  - II. *m*-Dichlorobenzene I. Toluene
  - III. o-Dichlorobenzene IV. p-Dichlorobenzene
  - (a) I < IV < II < III(b) IV < I < II < III(c) IV < I < III < II(d) IV < II < III < I
- **13.** The repeating unit in silicone is

(a) 
$$\operatorname{SiO}_2$$
 (b)  $-\operatorname{Si}_{i} - O - \operatorname{R}_{i}$   
(c)  $\operatorname{O}_{i} - \operatorname{Si}_{i} - O - \operatorname{O}_{i}$  (d)  $-\operatorname{Si}_{i} - O - O - R$   
 $R$ 

- 14. When nitrobenzene is treated with  $Br_2$  in presence of FeBr<sub>3</sub>, the major product is monobromonitrobenzene. Which of the following statements is related to the process?
  - (a) Electron density on *meta*-carbon is more than that on ortho- or para-carbons.
  - (b) Loss of aromaticity occurs, when Br<sup>+</sup> attacks at the ortho- or para-positions and not at *meta*-position.
  - (c) Intermediate carbocation formed after initial attack of Br<sup>+</sup> at the *meta*-position is least stabilised.
  - (d) Easier loss of  $H^+$  from *meta*-position than from ortho- and para -position to regain aromaticity.
- 15. Which of the following plots correctly represents variation of equivalent conductance  $(\Lambda_{eq})$  with dilution for a strong electrolyte?



16. Which of the following does not represent the mathematical expression for the Heisenberg uncertainty principle?

(a) 
$$\Delta x.\Delta p \ge \frac{h}{4\pi}$$
 (b)  $\Delta x.\Delta v \ge \frac{h}{4\pi m}$   
(c)  $\Delta E.\Delta t \ge \frac{h}{4\pi}$  (d)  $\Delta E.\Delta x \ge \frac{h}{4\pi}$ 

17. Which of the following has maximum energy?



18. In the following reactions sequence,

$$Ph \xrightarrow{O} OH \xrightarrow{Heat} A \xrightarrow{I_2} B + C$$

(\* implies <sup>13</sup>C labelled carbon)

the correct structures of A, B and C are

(a) A = Ph  $CH_3$ , B = Ph ONa,  $C = CHI_3$ (b)  $A = Ph CH_3, B = Ph ONa$ ,  $C = CHI_3$ (c)  $A = \bigwedge_{*}^{O} B = \bigwedge_{-+}^{O} C C = CHI_3$ 

(d) 
$$A = \underbrace{Ph}_{CH_3} \underbrace{Ph}_{CH_3} B = \underbrace{Ph}_{Ph} \underbrace{ONa}_{CH_3} CH_3 I$$

- **19.** Extraction of zinc from zinc blende is achieved by
  - (a) electrolytic reduction
  - (b) roasting followed by reduction with carbon
  - (c) roasting followed by reduction with another metal
  - (d) roasting followed by self-reduction.
- 20. The pair of amphoteric hydroxides is
  - (a) Al(OH)<sub>3</sub>, LiOH
  - (b)  $Be(OH)_2$ ,  $Mg(OH)_2$
  - (c)  $B(OH)_3$ ,  $Be(OH)_2$
  - (d)  $Be(OH)_2$ ,  $Zn(OH)_2$ .



- 21. A solution containing 2.675 g of CoCl<sub>3</sub>.6NH<sub>3</sub> (molar mass =  $267.5 \text{ g mol}^{-1}$ ) is passed through a cation exchanger. The chloride ions obtained in solution were treated with excess of AgNO<sub>3</sub> to give 4.75 g of AgCl (molar mass =  $143.5 \text{ g mol}^{-1}$ ). The formula of the complex is
  - (a)  $[CoCl_3(NH_3)_3]$ (b)  $[CoCl(NH_3)_5]Cl_2$

(c) 
$$[Co(NH_3)_6]Cl_3$$
 (d)  $[CoCl_2(NH_3)_4]Cl_3$ 

- 22. To prepare a solution of concentration of 0.03 g/mL of AgNO<sub>3</sub>, what amount of AgNO<sub>3</sub> should be added in 60 mL of solution?
  - (a) 1.8 g (b) 0.8 g
  - (c) 0.18 g (d) None of these
- 23. A solution is 0.1 M with respect to  $Ag^+$ ,  $Ca^{2+}$ ,  $Mg^{2+}$  and  $Al^{3+}$ . Which will precipitate at lowest concentration of  $PO_4^{3-}$  ion when  $Na_3PO_4$  solution is added?
  - (a)  $Ag_3PO_4(K_{sp} = 1 \times 10^{-6})$
  - (a)  $\operatorname{Mg3}(\operatorname{PO}_4)_2(K_{sp} = 1 \times 10^{-33})$ (b)  $\operatorname{Ca}_3(\operatorname{PO}_4)_2(K_{sp} = 1 \times 10^{-33})$ (c)  $\operatorname{Mg3}(\operatorname{PO}_4)_2(K_{sp} = 1 \times 10^{-24})$ (d)  $\operatorname{AlPO}_4(K_{sp} = 1 \times 10^{-20})$
- 24. Which of the following is the correct representation of relative lowering of vapour pressure?

(a) 
$$\frac{P^{o}}{\Delta P} = \frac{P^{o} - P}{P^{o}}$$
 (b)  $\frac{\Delta P}{P^{o}} = \frac{P - P^{o}}{P^{o}}$   
(c)  $\frac{P^{o}}{\Delta P} = \frac{P^{o}}{P^{o} - P}$  (d)  $\frac{\Delta P}{P^{o}} = \frac{P^{o} - P}{P^{o}}$ 

**25.** The reaction of cyanamide  $(NH_2CN)$  with dioxygen was carried out in a bomb calorimeter, and  $\Delta U$  was found to be –742.7 kJ mol<sup>-1</sup> at 298 K. Calculate the enthalpy change for the reaction at 298 K.

$$NH_{2}CN_{(s)} + \frac{3}{2}O_{2(g)} \longrightarrow N_{2(g)} + CO_{2(g)} + H_{2}O_{(l)}$$
  
(a) -741.46 kJ mol<sup>-1</sup> (b) -743.9 kJ mol<sup>-1</sup>  
(c) +741.46 kJ mol<sup>-1</sup> (d) +743.9 kJ mol<sup>-1</sup>

26. Among the following which is least acidic?

- (a) Phenol (b) *o*-Cresol
- (d) *p*-Chlorophenol (c) *p*-Nitrophenol
- 27. Which curve corresponds to the temperature dependance of the rate constant (k) of a simple one step reaction?





**28.** It is desired to increase the volume of 80  $\text{cm}^3$  of a gas by 20% without changing pressure. To what temperature the gas should be heated if its initial temperature is 25 °C?

(d) 35 °C (c) 630.6 °C

29. 
$$CCl_3CH = CH_2 \xrightarrow{Cl_2 + H_2O} A$$
; A is  
(a)  $CCl_3CHCH_2Cl$  (b)  $CCl_3CHCH_2OH$ 

$$\begin{array}{ccc} & & & | \\ & OH & & Cl \\ (c) & CCl_3CHCH_2 & (d) & CCl_3CHCH_2 \\ & | & | & | \\ & Cl & Cl & OHOH \end{array}$$

**30.** Consider the following reaction,

$$CH_{2}-CH_{2}-CH-CH-CH_{3}-C$$

The correct explanation is

- (a) the product is formed due to nucleophilic substitution
- (b) the product is formed according to Saytzeff's rule
- (c) conjugated double bond product is formed due
- to higher stability by resonance (d)  $(CH_3)_3CO^-$  is a better leaving group.
- 31. Which of the following cannot be made by using Williamson's synthesis?
  - (a) Di-tert-butyl ether (b) Methoxybenzene
  - (c) Benzyl *p*-nitrophenyl ether
  - (d) Methyl tert-butyl ether
- **32.** Metal carbide on reaction with water forms  $CH_4$ . Carbide can be

(a)  $CaC_2$  (b)  $Mg_3C_2$  (c)  $Be_2C$ (d) SiC

- 33. Time required to deposit one millimole of aluminium metal by the passage of 9.65 amperes through aqueous solution of aluminium ion is
  - (a) 30 s (b) 10 s
  - (c) 30,000 s (d) 10,000 s.



- 34. The correct order of dipole moment is
  - (a)  $CH_4 < NF_3 < NH_3 < H_2O$
  - (b)  $NF_3 < CH_4 < NH_3 < H_2O$
  - (c)  $NH_3 < NF_3 < CH_4 < H_2O$
  - (d)  $H_2O < NH_3 < NF_3 < CH_4$
- 35. The major product of the following reaction,



36. Which of the following electrolytes will have maximum flocculation value for Fe(OH)<sub>3</sub> sol? (b) Na<sub>2</sub>S (a) NaCl

(d) K<sub>2</sub>SO<sub>4</sub> (c)  $(NH_4)_3PO_4$ 

37. Which of the following fluorides has the lowest melting point?

(a) BaF<sub>2</sub> (b)  $SrF_2$ (c)  $CaF_2$ (d)  $BeF_2$ 

- 38. Which of the following pairs have both the elements showing highest oxidation states equal to
  - [ns + (n-1)d] electrons
  - (b) Cr and Co (a) Ti and Fe
  - (c) Cr and Mn (d) Co and Ni.
- 39. Which form of trans-1, 4-cyclohexanediol is most stable?





- **40.** A gas z is bubbled through a solution containing  $x^{-}$  and  $y^{-}$  ions. If the reduction potential are in the order x > y > z, then
  - (a) y will oxidise z and not x
  - (b) y will oxidise x and not z
  - (c) y will oxidise both x and z
  - (d) y will reduce both x and z.

#### **ASSERTION AND REASON**

**Directions :** In the following guestions (41-60), a statement of assertion is followed by a statement of reason. Mark the correct choice as :

- (a) If both assertion and reason are true and reason is the correct explanation of assertion.
- (b) If both assertion and reason are true but reason is not the correct explanation of assertion.
- (c) If assertion is true but reason is false.
- (d) If both assertion and reason are false.
- **41.** Assertion : Photochemical smog is produced by nitrogen oxides. Reason : Vehicular pollution is a major source of nitrogen oxides.
- 42. Assertion : The reaction,  $2NO_{(g)} + O_{2(g)} \rightleftharpoons 2NO_{2(g)}$  is favoured in the forward direction with increase of pressure. **Reason :** The reaction is exothermic.
- **43.** Assertion : A catalyst enhances the rate of a reaction. Reason : A catalyst remains unchanged in mass and in chemical composition at the end of the reaction.
- 44. Assertion : The micelle formed by sodium stearate in water has – COO<sup>-</sup> groups at the surface. Reason : Surface tension of water is reduced by the addition of stearate.
- 45. Assertion : Kinetic gas equation is modified for real gases.

Reason : This modification is carried out with respect to actual volume of molecules and attractive forces between the gaseous molecules.

- 46. Assertion : The total number of stereoisomers shown by  $[Co(en)_2Cl_2]^+$  complex ion is three. **Reason :**  $[Co(en)_2Cl_2]^+$  complex ion has octahedral geometry.
- 47. Assertion: 0.1 M aqueous solution of glucose has same depression in the freezing point as 0.1 M aqueous solution of urea.

**Reason:**  $K_f$  value for both the solutions is same.

**48.** Assertion : Diamond and graphite do not have the same crystal structures.

Reason : Diamond is crystalline while graphite is amorphous.

24 📃

# The most Reliable and Featured books for **AIIMS** in the market



- 49. Assertion : Helium has the highest value of ionisation energy among all the elements known.Reason : Helium has the highest value of electron affinity among all the elements known.
- **50. Assertion :** In H<sub>2</sub>SO<sub>5</sub>, oxidation number of S is +6. **Reason :** There is one peroxide linkage in H<sub>2</sub>SO<sub>5</sub>.
- 51. Assertion : Absorption spectrum consists of some bright lines separated by dark spaces.Reason : Emission spectrum consists of dark lines.
- **52.** Assertion :  $S_N 2$  reaction of an optically active alkyl halide with an aqueous solution of KOH always gives an alcohol with opposite sign of rotation. **Reason** :  $S_N 2$  reactions proceed with inversion of configuration.
- **53.** Assertion : Benzene on heating with conc.  $H_2SO_4$  gives benzenesulphonic acid which when heated with superheated steam under pressure gives benzene.

Reason : Sulphonation is a reversible process.

- **54.** Assertion : Sucrose is a non-reducing sugar. Reason : It has glycosidic linkage.
- 55. Assertion : Calgon is not suitable for removing Ca<sup>2+</sup> and Mg<sup>2+</sup> ions from hard water.
  Reason : Calgon forms insoluble complex with Ca<sup>2+</sup> and Mg<sup>2+</sup> ions.
- **56. Assertion :** Anilinium chloride is more acidic than ammonium chloride.

**Reason :** Anilinium ion is resonance stabilized.

57. Assertion : Essential oils are purified by steam distillation.Reason : Essential oils are insoluble in water and

are non-volatile in steam.

- 58. Assertion : Aqueous solution of CoCl<sub>2</sub> is pink in colour. It turns blue in presence of conc. HCl.
   Reason : It is due to the formation of [CoCl<sub>4</sub>]<sup>2-</sup>.
- **59. Assertion :** Mercury is liquid at room temperature. **Reason :** In mercury, there is no unpaired *d*-electron and thus metallic bonding is weakest.
- **60.** Assertion : The bond order of helium is always zero. **Reason** : The number of electrons in bonding molecular orbital and antibonding molecular orbital is equal.

## 



Element	%	Atomic mass	Relative number of atoms	Simplest ratio of atoms
С	18.5	12	$\frac{18.5}{12} = 1.542$	1
Н	1.55	1	$\frac{1.55}{1} = 1.55$	1
Cl	55.04	35.5	$\frac{55.04}{35.5} = 1.55$	1
О	24.81	16	$\frac{24.81}{16} = 1.55$	1

Therefore, empirical formula of the compound is CHClO.

3. (a) : The conversion of metal sulphide to metal oxide involves the process of roasting *i.e.*, x is roasting. The metal oxides can then be converted to impure metal by reduction *i.e.*, 'y' is smelting.

The conversion of impure metal to pure metal involves a process of purification. Thus, 'z' is electrolysis.

4. (a) : In  $CaF_2$  crystal,  $Ca^{2+}$  ions are present at corners and at the centre of each face of the cube while  $F^-$  ions occupy all the tetrahedral voids.

5. (a) : Classical smog is formed by the combination of smoke, dust and fog containing sulphur dioxide from polluted air.

6. (a) : X-ray studies have shown that in ice, four hydrogen atoms tetrahedrally surround each oxygen atom.



7. (c) :  $SnO_2 + 4HCl \longrightarrow Sn^{4+} + 4Cl^- + 2H_2O$ Sn<sup>4+</sup> ions are preferentially adsorbed by SnO<sub>2</sub> particles.

8. (d) :  $nCF_2 = CF_2 \longrightarrow + CF_2 - CF_2 + \frac{1}{n}$ Teflon

Teflon is used for non-stick cookwares.



**9.** (b) : As we move from Cl to I, the electron gain enthalpy becomes less and less negative due to increase in the atomic radius. Due to small size of F, the electron-electron repulsions in the relatively compact 2p-subshell are comparatively large and hence, the incoming electron is not accepted with the same ease as in case of Cl. Consequently, the electron gain enthalpy of F is less negative than that of Cl.

10. (a) : Carboxylic acid is stronger acid than  $\overline{NH}_3$ , therefore, X is the strongest acid. Since, — COOH has -I-effect which decreases with distance, therefore, -I-effect is more pronounced on Z than on Y. As a result, Z is more acidic than Y. Thus, overall order of decreasing acidic strength is X > Z > Y.

11. (a) : AgCNS 
$$\implies Ag^+ + CNS^-$$
  
 $x \qquad x$   
AgBr  $\implies Ag^+ + Br^-$   
 $y \qquad y'$   
 $\therefore \quad [Ag^+] = (x + y), [CNS^-] = x, [Br^-] = y$   
 $K_{sp[AgCNS]} = [Ag^+] [CNS^-] = x (x + y)$   
 $\Rightarrow \quad 1 \times 10^{-12} = x(x + y)$  ...(i)  
and,  $K_{sp[AgBr]} = [Ag^+] [Br^-] = y(x + y)$   
 $\Rightarrow \quad 5 \times 10^{-13} = y(x + y)$  ...(ii)  
On solving eqn (i) and (ii), we get  
 $x = 8.16 \times 10^{-7} \text{ mol/L}, y = 4.08 \times 10^{-7} \text{ mol/L}$ 

**12.** (b) : Dipole moment of *p*-dichlorobenzene is zero whereas *o*- and *m*-dichlorobenzene have higher dipole moment than toluene. Again *o*-dichlorobenzene has higher dipole moment than *m*-dichlorobenzene. Hence, the order is IV < I < II < III.

**13.** (b) :  $R_2$ SiO – is the repeating unit in silicone.



This reaction is aromatic electrophilic substitution reaction.  $-NO_2$  group being a powerful electron withdrawing group deactivates the benzene ring and provides the electron density at *meta*-position which can be shown by resonating structures.



Hence, reaction occurs at *meta*-position not at *ortho*- or *para*-position.

**15.** (b) : Equivalent conductance increases with dilution and ultimately becomes constant.

**16.** (d) : Heisenberg uncertainty principle can be written as,

$$\Delta x.\Delta p \ge \frac{h}{4\pi} \text{ or } \Delta x.m\Delta v \ge \frac{h}{4\pi} \implies \Delta x.\Delta v \ge \frac{h}{4\pi m}$$
  
Also,  $\Delta p.\Delta x = \Delta E.\Delta t \Rightarrow \Delta E.\Delta t \ge \frac{h}{4\pi}$ 

**17.** (c) : The order of increasing energy of the subatomic orbitals is s . The energy in excited state is more than that in the ground state. Since option (c) is an excited state with maximum number of unpaired electrons, therefore it has maximum energy.



#### 19. (b)

**20.** (d) : Both  $Be(OH)_2$  and  $Zn(OH)_2$  are amphoteric in nature.

21. (c) : Moles of the complex =  $\frac{2.675}{267.5} = 0.01$ Moles of AgCl precipitated =  $\frac{4.75}{143.5} = 0.033$ Thus, 1 mole of the complex will precipitate =  $\frac{0.033}{0.01} = 3$  moles of AgCl

CHEMISTRY TODAY | MARCH '18

27

This means that 1 molecule of the complex contains 3 ionisable Cl. Hence, the formula is  $[Co(NH_3)_6]Cl_3$ .

22. (a) : Concentration in terms of g/mL is given as  $0.03 = \frac{\text{mass of AgNO}_3}{60}$ Mass of AgNO<sub>3</sub> =  $60 \times 0.03 = 1.8$  g **23.** (d) : (a)  $Ag_3PO_4 \Longrightarrow 3Ag^+ + PO_4^{3-}$  $K_{sp} = [Ag^+]^3 [PO_4^{3-}]$  $[PO_4^{3-}] = \frac{K_{sp}}{(0.1)^3} = \frac{1 \times 10^{-6}}{10^{-3}} = 10^{-3} M$ (b)  $Ca_{3}(PO_{4})_{2} \Longrightarrow 3Ca^{2+} + 2PO_{4}^{3-}$  $K_{sp} = [Ca^{2+}]^{3} [PO_{4}^{3-}]^{2}$  $10^{-33} = (0.1)^3 [PO_4^{3-}]^2$  $[PO_4^{3-}]^2 = \frac{10^{-33}}{10^{-3}} = 10^{-30}$  $[PO_4^{3-}] = 10^{-15} M$ (c)  $Mg_3(PO_4)_2 \Longrightarrow 3Mg^{2+} + 2PO_4^{3-}$  $[PO_4^{3-}]^2 = \frac{10^{-24}}{10^{-3}} = 10^{-21}$  $[PO_4^{3-}] = 3.16 \times 10^{-11} \text{ M}$ (d)  $AlPO_4 \Longrightarrow Al^{3+}_{0,1} + PO_4^{3-}$  $\left[PO_4^{3-}\right] = \frac{K_{sp}}{0.1} = \frac{10^{-20}}{0.1} = 10^{-19} \text{ M}$ 24. (d) 25. (a) : NH<sub>2</sub>CN<sub>(s)</sub> +  $\frac{3}{2}$ O<sub>2(g)</sub>  $\longrightarrow$  N<sub>2(g)</sub> + CO<sub>2(g)</sub>  $+ H_2O_{(1)}$  $\Delta n_g = n_p - n_r = 2 - \frac{3}{2} = \frac{1}{2} = 0.5 \text{ mol}$  $\Delta H = \Delta U + \Delta n_g RT$   $\Delta H = -742.7 \text{ kJ mol}^{-1} + (0.5 \text{ mol} \times 8.314 \times 10^{-3} \text{ kJ mol}^{-1} \text{ K}^{-1}$ × 298 K)

 $\Delta H = (-742.7 \text{ kJ} + 1238.786 \times 10^{-3} \text{ kJ}) \text{ mol}^{-1}$ = -741.46 kJ mol<sup>-1</sup>

**26.** (b) : When an electron withdrawing group (like  $-NO_2$ , -Cl) is attached to the phenol ring, it stabilises the negative charge on the oxygen of phenoxide ion. Due to this reason, acidic character of

28 CHEMISTRY TODAY | MARCH '18

phenol increases. But when an electron donating group (like  $-CH_3$ ) is attached to the phenol ring, it destabilises the ring and hence, acidic character of phenol decreases. Thus, the correct order of acidic character is *p*-nitrophenol > *p*-chlorophenol > *p*henol > *o*-cresol.

**27.** (c) : Rate constant *k* increases exponentially as *T* increases.

$$k = Ae^{-E_a/RT} = \frac{A}{e^{E_a/RT}}$$

**28.** (b) : The desired increase in the volume of gas is

20% of 80 cm<sup>3</sup> = 
$$\frac{80}{100} \times 20 = 16 \text{ cm}^3$$

**30.** (c) :  $(CH_3)_3CO^-$  is a better base than a nucleophile. Hence, elimination occurs. The product formed is resonance stabilised.

**31.** (a) : Di-*tert*-butyl ether cannot be made by Williamson's synthesis, since *tert*-alkyl halides prefer to undergo elimination rather than substitution, *i.e.*,

$$CH_{3} - CH_{3} - C$$

32. (c) : 
$$Be_2C + 4H_2O \longrightarrow 2Be(OH)_2 + CH_4$$
  
33. (a) :  $Al^{3+} + 3e^- \rightarrow Al$ 

For deposition of 1 mole of  $Al^{3+}$ , charge required

 $\equiv 3 \times 96500 \text{ C}$ 

For deposition of 1 millimole of Al<sup>3+</sup>, charge required  $\equiv 3 \times 96.500 \text{ C}$ 

As 
$$Q = It \implies t = \frac{Q}{I} = 3 \times \frac{96.5}{9.65} = 30$$
 s

34. (a) : In  $H_2O$ , electronegativity difference is highest. So, dipole moment is highest in H<sub>2</sub>O. CH<sub>4</sub> has a symmetrical tetrahedral structure and its dipole moment is zero. Net dipole moment of NF<sub>3</sub> is less than that of NH<sub>3</sub>.

35. (a) : This reaction will not proceed via  $S_N 1$ mechanism as the carbocation formed will be destabilised by the -I effect of flourine and -R effect of -NO<sub>2</sub> group. Benzyl halides are more reactive than aryl halides in S<sub>N</sub>2 reactions. Therefore, the reaction occurs in the side chain with inversion of configuration at the chiral centre.



36. (a) : Flocculation value  $\propto \frac{1}{\text{Coagulating power}}$ 

Greater the coagulating power of effective ion, lower will be the flocculation value of the electrolyte.

Fe(OH)<sub>3</sub> is a positively charged sol. According to Hardy Schulze rule, greater the charge on the oppositively charged ion of the electrolyte added, more effective it is in coagulation.

Among the four electrolytes, NaCl has the lowest coagulating power, so its flocculation value will be maximum.

**37.** (d) : Being covalent,  $BeF_2$  has the lowest melting point (554°C) while other fluorides melt at around 1400°C.

**38.** (c) : Cr and Mn show the highest oxidation states +6 and +7 respectively.

**39.** (a) : When intramolecular hydrogen bonding is possible between groups at 1 and 4 positions the molecule assumes a boat conformation rather than the chair conformation in which hydrogen bonding is not possible.

40. (a) : More the value of reduction potential, more is the tendency to accept electrons or get reduced. As gas z has lowest reduction potential, therefore, z will be oxidised easily.

**41.** (b) :  $NO_2 + U.V.$  light  $\rightarrow NO + O$ 

Atomic oxygen produced as above reacts with hydrocarbons to give a variety of free hydrocarbon radicals, aldehydes, ketones, O3, peroxyacetyl nitrate (PAN), etc. PAN mixes with the fog and gets condensed on smoke or dust particles in the air to form a smog which is called photochemical smog.

42. (b) : According to Le-Chatelier's principle, with increase of pressure, equilibrium shifts in that direction in which lesser number of moles of gaseous products are produced.

43. (b)

44. (b) : Micelle is formed if molecules with polar and non-polar ends assemble in bulk to give non-polar interior and polar exterior.

45. (a) : The kinetic gas equation for one mole of a real gas is  $\left(p + \frac{a}{V_m^2}\right)(V_m - b) = RT$ 

This is known as van der Waals' equation.

**46.** (b):  $[Co(en)_2Cl_2]^+$  exist in two geometrical isomers cis and trans. Trans isomer does not show optical isomerism since it is symmetrical while cis-isomer shows optical isomerism and exists as *d*- and *l*-form. So, total three stereoisomers are possible.

47. (b) : Depression in freezing point is a colligative property which depends on the number of particles present in the solution. As both 0.1 M solution of glucose and 0.1 M solution of urea contain same number of particles therefore, both will have same depression in freezing point.

**48.** (c) : Diamond and graphite both are crystalline. Crystalline structure is different as in diamond, C-atoms are  $sp^3$ -hybridised while in graphite, they are  $sp^2$ hybridised.



**49.** (c) : Helium contains fully filled  $1s^2$  orbital which has more penetrating effect and is very close to the nucleus and hence, has higher value of ionisation energy. The electron affinity of He is the lowest of all the noble gases, due to the smallest size of He.

50. (a)

**51.** (d) : Absorption spectrum is produced when white light is passed through a substance and transmitted light is analysed by a spectrograph. The dark spaces corresponds to the light radiation absorbed by the substance. An emission spectrum is produced by analysing the radiant energy emitted by an excited substance by a spectrograph. Thus, discontinuous spectra consisting of a series of sharp lines and separated by dark bands are obtained.

52. (a)

53. (a) : 
$$\langle O \rangle$$
 + HOSO<sub>3</sub>H  $\langle O \rangle$  + H<sub>2</sub>O

**54.** (a) : Sucrose is a non-reducing sugar as it does not reduce Tollens' or Fehling's reagent, due to absence of free aldehyde or ketone group. It contains stable acetal

or ketal structure which cannot be opened into a free carboxyl group.

Sugar is composed of  $\alpha$ -*D*-glucopyranose unit and  $\beta$ -*D*-fructofuranose unit. These units are joined by  $\alpha$ ,  $\beta$ -glycosidic linkage between C-1 of the glucose unit and C-2 of the fructose unit.

55. (d) : Calgon is used to treat hard water by removing  $Ca^{2+}$  and  $Mg^{2+}$  ions in the form of soluble complexes.

**56.** (c) : Anilinium ion is not resonance stabilised since nitrogen does not have a lone pair of electrons.

**57.** (c) : Essential oils are insoluble in water and are volatile in steam.

58. (a) : 
$$[Co(H_2O)_6]^{2+}_{(aq.)} + 4Cl^-_{(aq)} \rightleftharpoons [CoCl_4]^{2-}_{(aq.)} + 6H_2O_{(l)}$$

According to Le Chatelier's principle, on adding conc. HCl, the equilibrium shifts in forward direction giving blue colour.

ی ک



CO II

CHEMISTRY TODAY | MARCH '18

30

**GEAR UP** FOR JEE MAIN **EXAMON** 8<sup>th</sup> April (offline) 15<sup>th</sup> & 16<sup>th</sup> April (online) 2018

- 1. Which of the following thermodynamic conditions at constant pressure and temperature is necessary for the spontaneity of a process?
  - (a) d(U TS + PV) > 0
  - (b) d(U TS + PV) < 0
  - (c) d(U TS + PV) = 0(d) d(U + TS + PV) < 0
- The same dust (III) of the fallowing rest time second



- **3.** In the manufacture of H<sub>2</sub>SO<sub>4</sub>, the nitrated acid from the Gay-Lussac's tower is chemically
  - (a)  $NO_2 \cdot H_2 SO_4$
  - (b) NO·H<sub>2</sub>SO<sub>4</sub>
  - (c)  $NO \cdot 2H_2SO_4$
  - (d) NO·HSO<sub>4</sub>
- 4. Find the equilibrium constant for the reaction,  $Cu^{2+} + In^{2+} \rightleftharpoons Cu^{+} + In^{3+}$ Given that,  $E_{Cu^{2+}|Cu^{+}}^{\circ} = 0.15$  V,  $E_{In^{2+}|In^{+}}^{\circ} = -0.4$  V,  $E_{In}^{\circ}{}^{3+}|_{In^{+}} = -0.42$  V (a)  $10^{10}$  (b)  $10^{15}$ 
  - (c)  $10^{20}$  (d)  $10^{18}$
- 5. The reaction of cyclooctyne with HgSO<sub>4</sub> in the presence of aqueous H<sub>2</sub>SO<sub>4</sub> gives



6. For the two gaseous reactions, following data is given :

$$A \to B; \ k_1 = 10^{10} \ e^{-20000/T}$$

$$C \rightarrow D; \ k_2 = 10^{12} \ e^{-24606/T}$$

the temperature at which  $k_1$  becomes equal to  $k_2$  is (a) 400 K (b) 1000 K

- (c) 800 K (d) 1500 K
- **7.** The number of hexagonal faces present in a truncated octahedron is
  - (a) 6 (b) 8
  - (c) 4 (d) 16
- 8. Identify the product for the given reaction.

CH<sub>3</sub>−CH<sub>2</sub>−C−OC<sub>2</sub>H<sub>5</sub>−
$$\frac{Na\ddot{O}H}{H_2O}$$
 Product  
(a) CH<sub>3</sub>−CH<sub>2</sub>−C− $\overset{O}{C}$ − $\overset{O}{O}$   
(b) CH<sub>3</sub>−CH<sub>2</sub>−C−O<sup>-</sup>  
(c) CH<sub>3</sub>CH<sub>2</sub>− $\overset{*}{O}$ −H

- (d) Both (a) and (b)
- 9. In the given Ellingham diagram, *X*, *Y* and *Z* represent graph for metal oxides. At temperature below 983 K



- (a) Y will reduce oxide Z
- (b) *Y* will reduce oxide *X*
- (c) Z will reduce oxide X
- (d) Z will reduce oxide Y.



- **10.** Which one of the following represents the correct increasing order of bond angles in the given molecules?
  - (a)  $H_2O < OF_2 < OCl_2 < ClO_2$
  - (b)  $OCl_2 < ClO_2 < H_2O < OF_2$
  - (c)  $OF_2 < H_2O < OCl_2 < ClO_2$
  - (d)  $ClO_2 < OF_2 < OCl_2 < H_2O$
- **11.** Predict the order of  $\Delta_o$  for the following compounds :
  - I.  $[Fe(H_2O)_6]^{2+}$
  - II.  $[Fe(CN)_2(H_2O)_4]$
  - III.  $[Fe(CN)_4(H_2O)_2]^{2-1}$
  - (a) (I) < (II) < (III)
  - (b) (II) < (I) < (III)
  - (c) (III) < (II) < (I)
  - (d) (II) < (III) < (I)
- **12.** Reaction of cyclohexanone with dimethylamine in the presence of catalytic amount of an acid forms a compound if water during the reaction is continuously removed. The compound formed is generally known as
  - (a) an enamine (b) a Schiff's base
  - (c) an amine (d) an imine.
- 13. Which of the following is aromatic?

(a)	[10]-Annulene	(b) [14]-Annulene
(c)	[16]-Annulene	(d) [18] -Annulene

**14.** The electronegativities of H and Cl are 2.1 and 3.0 respectively. The correct statement about the nature of HCl is

(a)	17% ionic	(b) 83% ionic
( )	FOO/ inmin	(1) 1000/ is a is

- (c) 50% ionic (d) 100% ionic.
- **15.** Identify I, II and III for the given reactions sequence.  $FeCr_2O_4 \xrightarrow{I} Na_2CrO_4 \xrightarrow{II} Na_2Cr_2O_7 \xrightarrow{III}$

		Na <sub>2</sub> Cr <sub>2</sub> C
Ι	II	III
(a) $Na_2CO_3/air$ ,	$\Delta$ H <sub>2</sub> SO <sub>4</sub>	С
(b) NaOH/air, $\Delta$	C, Δ	C, Δ
(c) Na <sub>2</sub> CO <sub>3</sub> /air,	$\Delta$ C, $\Delta$	C, Δ
(d) NaOH/air, $\Delta$	Al, $\Delta$	C, Δ

16. 1.325 g sample of fertilizer is heated with  $H_2SO_4$ and then treated with alkali. The gas evolved is passed into 50.0 mL of 0.2030 N  $H_2SO_4$ . 25.32 mL of 0.1980 N NaOH are required for the titration of unused acid. The percentage of nitrogen in the fertilizer is

(a)	5.30%	(b) 5.43%
(c)	4.99%	(d) 6.01%

17. A tripeptide (X) on partial hydrolysis gave two dipeptides *Cys-Gly* and *Glu-Cys*, *i.e.*,

Identify the tripeptide.

(a) *Glu-Cys-Gly* (b) *Gly-Glu-Cys* 

Cys-Gly

- (c) Cys-Gly-Glu (d) Cys-Glu-Gly
- 18. Which is not correct?
  - (a)  $Ge(OH)_2$  is amphoteric.
  - (b)  $GeCl_2$  is more stable than  $GeCl_4$ .
  - (c)  $GeO_2$  is weakly acidic.
  - (d)  $\operatorname{GeCl}_4$  in HCl forms  $[\operatorname{GeCl}_6]^{2-}$  ion.
- 19. The enthalpy change involved in the oxidation of glucose is -2880 kJ mol<sup>-1</sup>. 25% of this energy is available for muscular work. If 100 kJ of muscular work is needed to walk one km, what is the maximum distance that a person will be able to walk after eating 120 g of glucose?
  - (a) 4.80 km (b) 5.25 km
  - (c) 3.80 km (d) 5.75 km
- **20.** Elastol is a polymer used to cleanup oil spill. It is a non-toxic, non-dispersant chemical. One gallon can remove 150 gallons of heavy oil. The monomer of elastol is

(a) 
$$CH_2 = CH - CN$$
 (b)  $CH_2 = C - CH_3$   
(c)  $CH_2 = C - COOCH_3$ 

(d) 
$$CH_2 = CH - Ph$$

ĊН-

- **21.** Which of the following reagents reacts differently with HCHO, CH<sub>3</sub>CHO and CH<sub>3</sub>COCH<sub>3</sub>?
  - (a) HCN (b) NH<sub>2</sub>NH<sub>2</sub>
  - (c)  $NH_2OH$  (d)  $NH_3$



## iranchembook.ir/edu

- **22.** *Fac-mer* isomerism is associated with which one of the following complexes?
  - (a)  $[M(AA)_2]$  (b)  $[MA_3B_3]$

(c)  $[M(AA)_3]$  (d) [MABCD]

**23.** Which of the following ions, derive from cyclooctatetraene (COT) by adding or removing a suitable number of  $\pi$ -electrons, are aromatic?

$$\begin{array}{ccccccc} C_8 H_8^+ & C_8 H_8^{2+} & C_8 H_8^- & C_8 H_8^{2-} \\ I & II & III & IV \\ \mbox{(a) I, II} & \mbox{(b) II, IV} & \mbox{(c) III, IV} & \mbox{(d) II, III} \end{array}$$

- **24.** Ammonia forms complexes with Ag<sup>+</sup> ion according to the following reactions :
  - (i)  $[\operatorname{Ag}(\operatorname{H}_{2}\operatorname{O})_{2}]_{(aq)}^{+} + \operatorname{NH}_{3(aq)} \rightleftharpoons$  $[\operatorname{Ag}(\operatorname{NH}_{3})(\operatorname{H}_{2}\operatorname{O})]_{(aq)}^{+} + \operatorname{H}_{2}\operatorname{O}_{(l)}$ (ii)  $[\operatorname{Ag}(\operatorname{NH}_{3})(\operatorname{H}_{2}\operatorname{O})]_{(aq)}^{+} + \operatorname{NH}_{3(aq)} \rightleftharpoons$  $[\operatorname{Ag}(\operatorname{NH}_{3})_{2}]_{(aq)}^{+} + \operatorname{H}_{2}\operatorname{O}_{(l)}$

The equilibrium constants for the reaction (i) and (ii) are  $2.0 \times 10^3$  and  $8.3 \times 10^3$  respectively, then the equilibrium constant of the following reaction is  $[Ag(H_2O)_2]^+_{(aq)} + 2NH_{3(aq)} \rightleftharpoons$ 

- (a)  $4.15 \times 10^{6}$ (b)  $2.0 \times 10^{3}$ (c)  $8.3 \times 10^{3}$ (d)  $16.6 \times 10^{6}$
- **25.** In spite of being an odd-electron molecule,  $ClO_2$  does not dimerise because
  - (a) the odd electron is delocalised
  - (b) the odd electron is localised on the chlorine atom
  - (c) the two Cl O bonds do not have the same length
  - (d) of  $p_x p_x$  bonding in the chlorine atom.
- **26.** During the preparation of arenediazonium salts, the excess of nitrous acid is destroyed by adding
  - (a) aq. NaOH (b) aq.  $Na_2CO_3$
  - (c) aq.  $NH_2CONH_2$  (d) aq. KI.
- **27.** Which of the following is a non-hypnotic tranquilizer?
  - (a) Chlordiazepoxide (b) Meprobamate
  - (c) Equanil (d) All of these
- **28.** The Schrodinger wave equation for hydrogen atom is

$$\Psi_{2s} = \frac{1}{4(2\pi)^{1/2}} \left(\frac{1}{a_0}\right)^{5/2} \left(2 - \frac{r}{a_0}\right) e^{-r/2a_0}$$

where,  $a_0$  is Bohr's radius. Let the radial node in 2*s* be at  $r_0$  then the value of *r* in terms of  $a_0$  is

(a) 
$$r = a_0/2$$
 (b)  $2r = a_0$   
(c)  $r = 2a_0$  (d)  $r = a_0$ 

**29.** Identify the product (*A*) in the given reaction,  $CH_3 \swarrow \stackrel{CH_2}{\longrightarrow} \stackrel{LiAlH_4}{\longrightarrow} A.$ 

CH<sub>3</sub> 
$$O$$
 EOH  
(a) (CH<sub>3</sub>)<sub>2</sub>CHCH<sub>2</sub>OH (b) CH<sub>3</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>OH  
(c) (CH<sub>3</sub>)<sub>3</sub>COH (d) CH<sub>3</sub>CHOHCH<sub>2</sub>CH<sub>3</sub>

(a)  $HMnO_4$  (b)  $HMnO_5$ (c)  $H_2MnO_4$  (d)  $H_2MnO_3$ 

SOLUTIONS

1. (b)  
2. (b):  

$$\int_{c} \overset{c}{C} \equiv CH \xrightarrow{O_{3}} \underbrace{\int_{c} \overset{c}{COOH}}_{COOH + 2HCOOH} \xrightarrow{COOH}_{c} \overset{c}{COOH}_{COOH + 2HCOOH}$$

$$\int_{c} \overset{H_{3}O^{+}}{Ph} - \overset{c}{C} \underbrace{O}_{OMgBr} \overset{PhMgBr}{CO_{2}} \overset{c}{CO}_{(II)} \xrightarrow{A} \underbrace{COOH}_{c} \xrightarrow{PhMgBr} \overset{c}{C}_{(II)} \xrightarrow{C}_{c} \xrightarrow{COOH}_{c}$$
PhČOOH  
(III)  
3. (d): 2H\_{2}SO\_{4} + NO + NO\_{2} \rightarrow 2NO \cdot HSO\_{4} + H\_{2}O
4. (a):  $Cu^{2+} + e^{-} \rightarrow Cu^{+}; \qquad \Delta G_{1}^{\circ} = -0.15 \text{ F}_{1n}^{2+} + e^{-} \rightarrow In^{+}; \qquad \Delta G_{2}^{\circ} = +0.40 \text{ F}_{1n}^{2+} + e^{-} \rightarrow In^{+}; \qquad \Delta G_{2}^{\circ} = +0.40 \text{ F}_{1n}^{2+} + e^{-} \rightarrow In^{+}; \qquad \Delta G_{2}^{\circ} = -0.54 \text{ F}_{1n}^{2+} + e^{-} \rightarrow Cu^{+} + In^{3+}; \qquad \Delta G^{\circ} = -0.59 \text{ F}_{1n}^{2+} - 1n^{2+} \rightarrow Cu^{+} + In^{3+}; \qquad \Delta G^{\circ} = -0.59 \text{ F}_{1n}^{2+} - 1n^{2+} \rightarrow Cu^{+} + In^{3+}; \qquad \Delta G^{\circ} = -0.59 \text{ F}_{1n}^{2+} - 1n^{2+} \rightarrow Cu^{+} + In^{3+}; \qquad \Delta G^{\circ} = -0.59 \text{ F}_{1n}^{2+} - 1n^{2+} \rightarrow Cu^{+} + In^{3+}; \qquad \Delta G^{\circ} = -0.59 \text{ F}_{1n}^{2+} - 1n^{2+} \rightarrow Cu^{+} + In^{3+}; \qquad \Delta G^{\circ} = -0.59 \text{ F}_{1n}^{2+} - 1n^{2+} \rightarrow Cu^{+} + In^{3+}; \qquad \Delta G^{\circ} = -0.59 \text{ F}_{1n}^{2+} - 1n^{2+} \rightarrow Cu^{+} + In^{3+}; \qquad \Delta G^{\circ} = -0.59 \text{ F}_{1n}^{2+} - 1n^{2+} \rightarrow Cu^{+} + In^{3+}; \qquad \Delta G^{\circ} = -0.59 \text{ F}_{1n}^{2+} - 1n^{2+} \rightarrow Cu^{+} + In^{3+}; \qquad \Delta G^{\circ} = -0.59 \text{ F}_{1n}^{2+} - 1n^{2+} \rightarrow Cu^{+} + In^{3+}; \qquad \Delta G^{\circ} = -0.59 \text{ F}_{1n}^{2+} - 1n^{2+} \rightarrow Cu^{+} + In^{3+}; \qquad \Delta G^{\circ} = -0.59 \text{ F}_{1n}^{2+} - 1n^{2+} \rightarrow Cu^{+} + In^{3+}; \qquad \Delta G^{\circ} = -0.59 \text{ F}_{1n}^{2+} - 1n^{2+} \rightarrow Cu^{+} + 1n^{3+}; \qquad \Delta G^{\circ} = -0.59 \text{ F}_{1n}^{2+} - 1n^{2+} \rightarrow Cu^{+} + 1n^{2+} \rightarrow Cu^$ 

5. (d): 
$$H_2O \xrightarrow{H_2SO_4/HgSO_4}$$
  
Cyclooctyne
$$\left[ \underbrace{\bigcirc H}_H \right] \xrightarrow{\text{Tautomerises}}_{\text{Cyclooctanone}} \underbrace{\bigcirc O}_{\text{Cyclooctanone}}$$



6. (b): 
$$A \to B$$
;  $k_1 = 10^{10}e^{-20000/T}$   
 $C \to D$ ;  $k_2 = 10^{12}e^{-24606/T}$   
When,  $k_1 = k_2$   
 $10^{10}e^{-20000/T} = 10^{12}e^{-24606/T}$   
 $e^{4606/T} = 100$   
 $\frac{4606}{T} = 2.303 \log 100 = 2.303 \times 2$   
 $\therefore T = \frac{4606}{2.303 \times 2} = 1000 \text{ K}$ 

7. (b): Truncated octahedron has 14 faces, 8 regular hexagonals and 6 squares.



- 9. (a) :  $\Delta G^{\circ}$  of *Y* is less than *Z* and hence, it will reduce oxide of *Z*.
- 10. (c):  $O_{103^{\circ}} < O_{1045^{\circ}} < O_{111^{\circ}} < O_{1$
- 11. (a) : The value of  $\Delta_o$  for mixed ligands depends on the additive contributions of the ligand strengths. Since,  $CN^-$  has greater ligand strength than H<sub>2</sub>O, the strength increases as the number of  $CN^-$  ions increases. Hence, the correct order of  $\Delta_o$  is III > II > I.
- 12. (a) :



34

CHEMISTRY TODAY | MARCH '18

13. (d): [18]-Annulene is aromatic since it is planar and contains (4n + 2) π-electrons. Although annulenes [10] and [14] also contain (4n + 2) π-electrons but the crowding of hydrogens inside the ring prevents planarity and hence are not aromatic.

**14.** (a) : % ionic character = 
$$16(\chi_A - \chi_B) + 3.5(\chi_A - \chi_B)^2$$
  
=  $16(3.0 - 2.1) + 3.5(3.0 - 2.1)^2$   
=  $14.4 + 2.835 = 17.235 ≈ 17\%$ 

15. (a):

$$4\text{FeCr}_{2}\text{O}_{4} + \underbrace{8\text{Na}_{2}\text{CO}_{3} + 7\text{O}_{2}}_{(I)} \xrightarrow{\Delta}$$

$$8\text{Na}_{2}\text{CrO}_{4} + 2\text{Fe}_{2}\text{O}_{3} + 8\text{CO}_{2}$$

$$2\text{Na}_{2}\text{CrO}_{4} \xrightarrow{\text{H}_{2}\text{SO}_{4}(\text{II})} \text{Na}_{2}\text{Cr}_{2}\text{O}_{7} + \text{Na}_{2}\text{SO}_{4} + \text{H}_{2}\text{O}_{3}\text{CO} + \text{Na}_{2}\text{Cr}_{2}\text{O}_{4} \xleftarrow{3\text{C(III)}}$$

- 16. (b) : Weight of sample (W) = 1.325 g Volume of acid  $(H_2SO_4)$  used  $(V_1) = 50$  mL Normality of acid  $(N_1) = 0.2030$  N Volume of alkali required  $(V_2) = 25.32$  mL Normality of alkali  $(N_2) = 0.1980$  N Milliequivalents of  $H_2SO_4$  left after reaction with NH<sub>3</sub> = Milliequivalents of alkali used for neutralisation of rest  $H_2SO_4 = N_2V_2 = 0.1980 \times 25.32 = 5.013$ Milliequivalents of  $H_2SO_4$  taken to absorb NH<sub>3</sub> =  $N_1V_1$ =  $0.2030 \times 50 = 10.15$ 
  - :. Milliequivalents of  $H_2SO_4$  which has reacted with  $NH_3(x) =$  Milliequivalents of acid taken

- Milliequivalents of acid left

$$= 10.15 - 5.013 = 5.137$$

Now, % N = 
$$\frac{1.4x}{W} = \frac{1.4 \times 5.137}{1.325} = 5.43\%$$

17. (a): Since the tripeptide on hydrolysis gave two dipeptides *Glu-Cys* and *Cys-Gly*. Hence, cystine must be in between glutamic acid and glycine.

**18.** (b):  $Ge^{4+}$  is more stable than  $Ge^{2+}$ , thus  $GeCl_4$  is more stable than  $GeCl_2$ .

**19.** (a) : Molar mass of glucose  $(C_6H_{12}O_6) = 180 \text{ g mol}^{-1}$ Combustion reaction of glucose can be written as 
$$\begin{split} \mathrm{C_6H_{12}O_{6(s)}+6O_{2(g)}} &\to \mathrm{6CO_{2(g)}+6H_2O_{(l)};} \\ \Delta H &= -2880 \ \mathrm{kJ \ mol^{-1}} \end{split}$$

Number of moles of 120 g of glucose

$$= \frac{120 \text{ g}}{180 \text{ g mol}^{-1}} = \frac{2}{3} \text{ mol}$$

Enthalpy available from 120 g of glucose

$$=\frac{2}{3} \times 2880 = 1920 \text{ kJ}$$

Enthalpy available for muscular work =  $1920 \times \frac{25}{100}$  $= 480 \, \text{kJ}$ 

Distance to which a person can move

$$= \left(\frac{1\,\mathrm{km}}{100\,\mathrm{kJ}}\right) \times 480\,\mathrm{kJ} = 4.80\,\mathrm{km}$$

- **20.** (b): Elastol is a polymer of 2-methylpropene.
- 21. (d): NH<sub>3</sub> reacts differently. With HCHO, it gives urotropine, with acetaldehyde, it gives  $CH_3CH = NH$ and with acetone, it gives diacetone amine.

22. (b)

K

23. (b): Removal of two electrons from COT (cyclooctatetraene), gives cyclic species II ( $C_8H_8^{2+}$ ) which has 6  $\pi$ -electrons and hence is aromatic. Similarly, addition of two electrons to COT gives a cyclic species IV ( $C_8H_8^{2-}$ ) which has 10  $\pi$ -electrons and hence is also aromatic.

24. (d): 
$$[Ag(H_2O)_2]^+_{(aq)} + NH_{3(aq)} \rightleftharpoons$$
  
 $[Ag(NH_3)(H_2O)]^+_{(aq)} + H_2O_{(l)} \dots (1)$   
 $K_1 = 2.0 \times 10^3$ 

$$K_1 = \frac{[\text{Ag}(\text{NH}_3)(\text{H}_2\text{O})]^+_{(aq)}}{[\text{Ag}(\text{H}_2\text{O})_2]^+_{(aq)}[\text{NH}_{3(aq)}]} \qquad \dots (i)$$

$$[Ag(NH_{3})(H_{2}O)]^{+}_{(aq)} + NH_{3(aq)} \rightleftharpoons [Ag(NH_{3})_{2})]^{+}_{(aq)} + H_{2}O_{(l)} \dots (2)$$
  
$$K_{2} = 8.3 \times 10^{3}$$

$$_{2} = \frac{[Ag(NH_{3})_{2}]^{+}_{(aq)}}{\dots(ii)}$$

$$\begin{bmatrix} Ag(NH_3)(H_2O) \end{bmatrix}_{(aq)}^+ \begin{bmatrix} NH_{3(aq)} \end{bmatrix}$$

$$[\operatorname{Ag}(\operatorname{H}_{2}\operatorname{O})_{2}]^{+}_{(aq)} + 2\operatorname{NH}_{3(aq)} \rightleftharpoons [\operatorname{Ag}(\operatorname{NH}_{3})_{2}]^{+}_{(aq)} + 2\operatorname{H}_{2}\operatorname{O}_{(l)}$$

$$K = \frac{[\mathrm{Ag}(\mathrm{NH}_{3})_{2}]_{(aq)}^{+}}{[\mathrm{Ag}(\mathrm{H}_{2}\mathrm{O})_{2}]_{(aq)}^{+}[\mathrm{NH}_{3(aq)}]^{2}} \qquad \dots (\mathrm{i}\mathrm{i}\mathrm{i}\mathrm{i})$$

From equations (i), (ii) and (iii), it is clear that  $K = K_1 K_2 = 2 \times 10^3 \times 8.3 \times 10^3 = 16.6 \times 10^6$ 

**25.** (a): The odd electron of  $Cl_2$  is delocalised thus, ClO<sub>2</sub> does not dimerise.



- **26.** (c) : Urea decomposes  $HNO_2$  to  $N_2$ ,  $CO_2$  and  $H_2O$ .  $NH_2CONH_2 + 2HONO \rightarrow 2N_2^{\uparrow} + CO_2^{\uparrow} + 3H_2O$
- 27. (d)
- **28.** (c) :  $\psi_{2s}^2$  = probability of finding electrons within 2*s* sphere.

 $\psi_{at node}^2 = 0$  (probability of finding an electron is zero at node)

So, 
$$\psi^2 = 0 = \frac{1}{32\pi} \left[ \frac{1}{a_0} \right]^3 \left[ 2 - \frac{r}{a_0} \right]^2 \times e^{-\frac{r}{a_0}}$$
  
or  $\left[ 2 - \frac{r}{a_0} \right] = 0 \implies 2 = \frac{r}{a_0}$  or  $r = 2a_0$ 

29. (c) : In the basic medium, the reaction occurs by  $S_N2$  mechanism and the nucleophile, *i.e.*, H<sup>-</sup> ion (from LiAlH<sub>4</sub>) attacks the smaller alkyl group.



30. (a)

...(3)

Your favourite MTG Books/Magazines available in ORISSA at

Padmalaya Book Seller - Bhubaneswar Ph: 0674-2396922, 3039692; Mob: 9437026922 Pragnya - Bhubaneswar Ph: 0674-2395757, 30396922; Mob: 9861135495 Sri Ram Books - Bhubaneswar Ph: 0674-2391385; Mob: 9439921093 Sagar Book Store - Bhubaneswar Ph: 0674-2506040, 2516040; Mob: 8763406040 Sri Madhab Book Store - Cuttack Ph: 0671-2431158, 2431148; Mob: 9437024244 A.K. Mishra Agency Pvt. Ltd. - Cuttack Ph: 0671-2322244, 2322255, 2322266, 2332233; Mob: 9437025991 Kitab Mahal - Cuttack Ph: 0671-2648333, 2648301; Mob: 9861388145 Visit "MTG IN YOUR CITY" on www.mtg.in to locate nearest book seller OR write to info@mtg.in OR call 0124-6601200 for further assistance.



# **PRACTICE PAPER**



- Compressibility factor (Z) for N<sub>2</sub> at -50 °C and 800 atm pressure is 1.95. Calculate the number of moles of N<sub>2</sub> gas required to fill a gas cylinder of 100 mL capacity under the given conditions.
  - (a) 2.24 (b) 1.12
  - (c) 6.10 (d) 2.90
- Benzaldehyde reacts with ammonia to form

   (a) hydrobenzamide
   (b) benzamide
  - (c) aniline (d) phenyl cyanide.
- **3.** Which of the following is not a property of hydrophilic sols?
  - (a) High concentration of dispersed phase can be easily attained.
  - (b) Coagulation is reversible.
  - (c) Viscosity and surface tension are nearly same as that of water.
  - (d) The charge of the particle depends on the pH value of the medium; it may be positive, negative or even zero.
- **4.** Which of the following alcohols is most reactive with HCl in the presence of ZnCl<sub>2</sub>?

(a) 
$$CH_3$$
  
 $\downarrow$   
 $CH_3$   
 $CH_3$   
 $CH_3$   
 $CH_3$   
(b)  $CH_3$   
 $CH_3$   
 $CH_3$   
 $CH_3$   
 $CH_3$   
(c)  $CH_3$   
 $CH_$ 

- 5. A ball of mass 200 g is moving with a velocity of 10 m sec<sup>-1</sup>. If the error in measurement of velocity is 0.1%, the uncertainty in its position is
  - (a)  $3.32 \times 10^{-31}$  m
  - (b)  $3.34 \times 10^{-27}$  m
  - (c)  $5.32 \times 10^{-25}$  m
  - (d)  $2.64 \times 10^{-32}$  m

When neopentyl bromide is subjected to Wurtz

Exam on 6<sup>th</sup> May 2018

- reaction, the product formed is (a) 2, 2, 4, 4-tetramethylhexane
- (b) 2, 2, 4, 4-tetramethylpentane
- (b) 2, 2, 4, 4-tetrainethylpentane
- (c) 2, 2, 5, 5-tetramethylhexane
  (d) 2, 2, 3, 3-tetramethylhexane.
- 7. Which of the following statements about primary amines is false?
  - (a) Aryl amines react with nitrous acid to produce nitrophenols.
  - (b) Alkyl amines are stronger bases than ammonia.
  - (c) Alkyl amines are stronger bases than aryl amines.
  - (d) Alkyl amines react with nitrous acid to produce alcohols.
- 8. NaNO<sub>3</sub> when decomposes above 800 °C does not give

(a) 
$$N_2$$
 (b)  $O_2$  (c)  $NO_2$  (d)  $Na_2O$ 

**9.** Which of the following is a free radical substitution reaction?

(a) 
$$CH_3 + Cl_2 \xrightarrow{Boil} CH_2Cl$$
  
(b)  $+ CH_3Cl \xrightarrow{Anhyd. AlCl_3} CH_3$   
(c)  $CH_2Cl$   
 $+ AgNO_2 \longrightarrow CH_2NO_2$ 

(d)  $CH_3CHO + HCN \longrightarrow CH_3CH(OH)CN$ 

- 10. The density of sodium borohydride is  $1.074 \text{ g/cm}^3$ . 3.91 g of sodium borohydride contains  $2.50 \times 10^{23}$  atoms of H. The number of moles of H atoms present in 28.0 cm<sup>3</sup> of sodium borohydride is
  - (a) 3.192 (b) 2.03
  - (c) 1.67 (d) 1.92


- 11. The resistance of 0.5 N solution of an electrolyte in a conductivity cell was found to be 25 ohm. Calculate the equivalent conductivity of the solution if the electrodes in the cell are 1.6 cm apart and have an area of  $3.2 \text{ cm}^2$ .
  - (a)  $10 \text{ S cm}^2$  equiv (b)  $15 \text{ S cm}^2$  equiv
  - (c)  $20 \text{ S cm}^2$  equiv (d)  $40 \text{ S cm}^2$  equiv
- **12.** The volume strength of 1.5 N H<sub>2</sub>O<sub>2</sub> solution is (a) 4.8 (b) 8.4 (c) 3.0 (d) 8.0
- **13.** Aluminium crystallizes in a cubic close packed structure. Its metallic radius is 125 pm. What is the length of the side of unit cell?
  - (a) 145 pm (b) 353.5 pm
  - (c) 125 pm (d) 250 pm
- 14. The gases that give rise to photochemical smog are
  - (a) oxides of sulphur (b) oxides of nitrogen
  - (c) oxides of carbon (d) oxygen.
- **15.** Identify the final product (*Z*) in the following sequence of reactions :
  - $(CH_3)_2CO + HCN \longrightarrow X \xrightarrow{H_3O^+} Y \xrightarrow{H_2SO_4}_{Heat} Z$ (a) (CH\_3)\_2C(OH)COOH (b) CH\_2=C(CH\_3)COOH (c) HOCH\_2CH(CH\_3)COOH (d) CH\_3CH=CHCOOH
- 16. Calculate the longest wavelength (in Å) which can remove the electron from first Bohr's orbit. (Given : *E*<sub>1</sub> = 13.6 eV)
  (a) 303.81 (b) 912.24 (c) 1095.12 (d) 1215.67
- 17. The product of acid catalysed hydration of 2-phenyl propene is
  - (a) 3-phenyl-2-propanol (b) 1-phenyl-2-propanol
  - (c) 2-phenyl-2-propanol (d) 2-phenyl-1-propanol.
- **18.** Which of the following statements is not true about glucose?
  - (a) It is an aldohexose.
  - (b) On heating with HI it forms *n*-hexane.
  - (c) It is present in furanose form.
  - (d) It does not give 2,4-DNP test.
- **19.** In a system :  $A_{(s)} \rightleftharpoons 2B_{(g)} + 3C_{(g)}$ , if the concentration of *C* at equilibrium is increased by a factor 2, it will cause the equilibrium concentration of *B* to change by
  - (a) two times of its original value
  - (b) one half of its original value
  - (c)  $2\sqrt{2}$  times of its original value
  - (d)  $\frac{1}{2\sqrt{2}}$  time of its original value.

- **20.** Which of the following statements is not true?
  - (a) The Ellingham diagram shows the plots of  $\Delta G$  vs *T*.
  - (b) In froth floatation process, depressants are added to enhance the formation of froth.
  - (c) Extraction of zinc oxide is done by coke.
  - (d) CO is more effective reducing agent below 983 K.
- **21.** In a mixture of *A* and *B*, components show -ve deviations as
  - (a)  $\Delta V_{\text{mix}}$  is +ve
  - (b) *A*–*B* interactions are weaker than *A*–*A* and *B*–*B* interactions
  - (c)  $\Delta H_{\text{mix}}$  is +ve
  - (d) *A*–*B* interactions are stronger than *A*–*A* and *B*–*B* interactions.
- 22. Out of vanadium (V), chromium (Cr), manganese (Mn) and iron (Fe), which one is expected to have the highest second ionisation enthalpy?
  (a) V (b) Cr (c) Mn (d) Fe
- **23.** In a first order reaction, the initial amount of a substance becomes 1/3 in 100 seconds. How much time will be taken to reduce the concentration to 1/9 of the initial concentration?
  - (a) 200 sec (b) 100 sec (c) 50 sec (d) 400 sec
- 24. Among the following halides :

 1. BCl<sub>3</sub>
 2. AlCl<sub>3</sub>
 3. GaCl<sub>3</sub>
 4. InCl<sub>3</sub>

 the order of decreasing Lewis acid character is
 (a)
 1, 2, 3, 4
 (b)
 4, 3, 2, 1

 (c)
 3, 4, 2, 1
 (d)
 2, 3, 4, 1.

- **25.** Which of the following statements is not true about low density polythene?
  - (a) Obtained through free radical addition
  - (b) Chemically inert and tough
  - (c) Good conductor of electricity
  - (d) Highly branched structure
- **26.** Identify a reagent from the following which can easily distinguish between but-1-yne and but-2-yne.
  - (a) Bromine, CCl<sub>4</sub>
  - (b) H<sub>2</sub>, Lindlar's catalyst
  - (c) Dilute  $H_2SO_4$ ,  $HgSO_4$
  - (d) Ammoniacal Cu<sub>2</sub>Cl<sub>2</sub> solution
- 27. Which of the following is paramagnetic in nature?
  - (a)  $[Cr(CO)_6]$  (b)  $[Fe(CO)_5]$ (c)  $[Fe(CN)_6]^{4-}$  (d)  $[Cr(NH_3)_6]^{3+}$
- **28.** Which of the following contains maximum number of lone pairs of electrons on the central atom?
  - (a)  $ClO_3^-$  (b)  $XeF_4$
  - (c)  $SF_4$  (d)  $I_3^-$ 
    - CHEMISTRY TODAY | MARCH '18



- 29. Match List I with List II and select the correct option.
  - List I List II (I) Iodoform (A) Anaesthetic (II) Methyl salicylate (B) Antiseptic
  - (III) Diethyl ether (C) Insecticide
  - (IV) Hexachlorocyclohexane (D) Detergent (E) Pain balm
  - (a) I B, II E, III C, IV D
  - (b) I D, II B, III A, IV C
  - (c) I B, II E, III A, IV C
  - (d) I C, II A, III D, IV B
- 30. In group 14, the inert-pair effect is more prominent in
  - (b) carbon and silicon (a) tin and lead
  - (c) carbon and lead (d) none of these.
- 31. Which is the most suitable reagent for the following conversion?  $\cap$

$$CH_{3}-CH=CH-CH_{2}-CH_{3}\longrightarrow O$$

$$CH_{3}-CH=CH-CH_{2}-CH_{3}-OH$$

- (a) Tollens' reagent
- (b) Benzoyl peroxide
- (c) I<sub>2</sub> and NaOH solution
- (d) LiAlH<sub>4</sub>/C<sub>2</sub>H<sub>5</sub>OH
- **32.** The values of  $T_c$  for few gases are given below : H<sub>2</sub>: 33.2 K, O<sub>2</sub>: 154.3 K, He: 5.3 K and CO<sub>2</sub>: 304.10 K. What is the correct increasing order of liquefaction of the above gases?
  - (a)  $\text{He} < \text{O}_2 < \text{H}_2 < \text{CO}_2$
  - (b)  $\text{He} < \text{H}_2 < \text{O}_2 < \text{CO}_2$
  - (c)  $CO_2 < O_2 < H_2 < He$
  - (d)  $O_2 < CO_2 < H_2 < He$
- 33. The basic character of the transition metal monoxides follows the order
  - (a) CrO > VO > FeO > TiO
  - (b) TiO > FeO > VO > CrO
  - (c) TiO > VO > CrO > FeO
  - (d) VO > CrO > TiO > FeO
- 34. 0.316 g of an organic compound, after heating with fuming nitric acid and barium nitrate crystals in a sealed tube gave 0.466 g of the precipitate of barium sulphate. The percentage of sulphur in the compound is

(c) 15.85 (d) 30.15

- 35. Nitrogen oxide that does not contain N—N bond is (a)  $N_2O$ (b)  $N_2O_3$  (c)  $N_2O_4$  (d)  $N_2O_5$
- **36.** Compare x and y bond angles for the given molecule :



- 37. In context with the transition elements, which of the following statements is incorrect?
  - (a) In addition to the normal oxidation states, zero oxidation state is also shown by elements in complexes.
  - (b) In the highest oxidation states, transition elements show basic character and form cationic complexes.
  - (c) In the highest oxidation states of the first five transition elements (Sc to Mn), all the 4s and 3d electrons are used for bonding.
  - (d) Once the  $d^5$  configuration is exceeded, the tendency to involve all the 3*d* electrons in bonds decreases.
- **38.** Enantiomers have
  - (a) identical m.pt./b.pt. but different refractive indices
  - (b) identical m.pt./b.pt. and refractive indices but rotate plane polarised light in opposite directions but to the same extent
  - (c) different refractive indices and rotate plane polarised light in the same direction but to different extent
  - (d) different m.pt./b.pt. but rotate plane polarised light in different directions but to the same extent.
- 39. A unit cell of sodium chloride has four formula units. The edge length of the unit cell is 0.564 nm. Density of sodium chloride is
  - (a)  $1.08 \text{ g cm}^{-3}$ (b)  $2.16 \text{ g cm}^{-3}$
  - (c)  $3.24 \text{ g cm}^{-3}$ (d) none of these.
- **40.** Some properties of the two species,  $NO_3^-$  and  $H_3O^+$ are described below. Which one of them is correct?
  - (a) Dissimilar in hybridisation for the central atom with different structures.
  - (b) Isostructural with same hybridisation for the central atom.
  - (c) Isostructural with different hybridisation for the central atom.



### iranchembook.ir/edu

- (d) Similar in hybridisation for the central atom with different structures.
- **41.** The plot of  $\log_{10}K$  vs 1/T leads to a straight line having intercept equal to

(a) 
$$\Delta G^{\circ}$$
 (b)  $\frac{\Delta G^{\circ}}{2.303R}$   
(c)  $\frac{\Delta S^{\circ}}{2.303R}$  (d)  $\frac{\Delta H^{\circ}}{2.303R}$ 

42. Which of the following complexes has magnetic moment of 2.83 B.M.?

(a) 
$$[Ni(NH_3)_6]^{2+}$$
 (b)  $[Ni(CN)_4]^2$   
(c)  $TiCl_4$  (d)  $[CoCl_6]^{3-}$ 

43. The final product of the following sequence of reactions

$$CaO + C \xrightarrow{Heat} A \xrightarrow{H_2O} B$$
, is

- (a) ethanol
- (b) ethyl hydrogen sulphate
- (c) acetylene (d) ethylene glycol.
- 44. The same quantity of electricity that liberated 2.158 g of Ag was passed through a gold salt, and 1.314 g of gold was deposited. The equivalent mass of Ag is 107.9. Calculate oxidation state of Au in the salt. (At. mass of Au = 197)

(a) 
$$+2$$
 (b)  $+3$  (c)  $+1$  (d) 0

**45.** If  $E_{M^+/M}^{\circ} = -1.2 \text{ V}$ ,  $E_{X_2/X^-}^{\circ} = 1.1 \text{ V}$  and  $E_{O_2/H_2O}^{\circ} = 1.23 \text{ V}$ , then on electrolysis of aqueous solution of salt MX, the products obtained are (a

a) 
$$M, X_2$$
 (b)  $H_2, X_2$  (c)  $H_2, O_2$  (d)  $M, O_2$ 

### SOLUTIONS

1. (a) : We have, 
$$Z = \frac{PV}{nRT}$$
  
∴ Mole of N<sub>2</sub>(n) =  $\frac{PV}{ZRT} = \frac{800 \times 100}{1.95 \times 0.0821 \times 223 \times 1000}$   
= 2.24

2. (a): 
$$\begin{array}{c} C_{6}H_{5}-CH=O+H_{2} \\ C_{6}H_{5}-CH=O+H_{2} \\ \end{array} \\ \xrightarrow{\begin{array}{c} C_{6}H_{5}-CH=O+H_{2} \\ O+H_{2} \\ NH \end{array}} \\ \xrightarrow{\begin{array}{c} C_{6}H_{5}CH=N \\ C_{6}H_{5}CH=N \\ Hydrobenzamide \end{array}} \\ \xrightarrow{\begin{array}{c} C_{6}H_{5}CH=N \\ Hydrobenzamide \end{array}} \\ \xrightarrow{\begin{array}{c} C_{6}H_{5}CH=N \\ Hydrobenzamide \end{array}}$$

3. (c) : Hydrophilic sols have lower surface tension and higher viscosity than that of water.

4. (a) : Order of reactivity of alcohols towards Lucas reagent :  $3^{\circ} > 2^{\circ} > 1^{\circ}$ 

5. (d) : 
$$\Delta v = \frac{0.1}{100} \times 10 = 10^{-2} \text{ m sec}^{-1}$$
;  
Now,  $\Delta v \cdot \Delta x = \frac{h}{4\pi m}$   
 $\Delta x = \frac{6.625 \times 10^{-34}}{4 \times 10^{-2} \times 3.14 \times 200 \times 10^{-3}} = 2.64 \times 10^{-32} \text{ m}$   
6. (c) :  $2\text{CH}_3 - \text{C} - \text{CH}_2\text{Br} + 2\text{Na} \xrightarrow{\text{Dry ether}}_{\text{CH}_3}$   
Neopentyl bromide  
 $C\text{H}_3$   
Neopentyl bromide  
 $C\text{H}_3 - \text{C} - C\text{H}_2 - C\text{H}_2 - C\text{H}_3$   
 $C\text{H}_3 - C\text{H}_3 - C\text{H}_3 - C\text{H}_3$   
 $C\text{H}_3 - C\text{H}_3$   
 $C\text{H}_3 - C\text{H}_3 - C\text{H}_3$   
 $C\text{H}_3 - C\text{H}_3$   
 $C\text{H}_3 - C\text{H}_3 - C\text{H}_3$   
 $C\text{H}_3 - C\text{H}_3 - C\text{H}_3$   
 $C\text{H}_3 - C\text{H}_$ 

7. (a)

8. (c) : NaNO<sub>3</sub> 
$$\xrightarrow{500 \circ C}$$
 NaNO<sub>2</sub> +  $\frac{1}{2}$ O<sub>2</sub>  
2NaNO<sub>3</sub>  $\xrightarrow{>800 \circ C}$  Na<sub>2</sub>O + N<sub>2</sub> +  $\frac{5}{2}$ O<sub>2</sub>

9. (a) : Side chain chlorination takes place in the presence of heat or light by free radical substitution mechanism.

**10.** (a) : Weight of sodium borohydride in  $28.0 \text{ cm}^3$  $= 28 \times 1.074 = 30.072$  g : 3.91 g of sodium borohydride has moles of H atoms  $250 \times 10^{23}$ 

$$\frac{2.50 \times 10}{6.023 \times 10^{23}}$$

:. 30.072 g of sodium borohydride has moles of  $2.50 \times 10^{23}$  30.072 Uatoma

H atoms = 
$$\frac{1}{6.023 \times 10^{23}} \times \frac{1}{3.91}$$
  
= 3.192 moles of H atoms  
11. (d) :  $\rho = R \cdot \frac{a}{l} = \frac{25 \times 3.2}{1.6} = 50$   
 $\kappa = \frac{1}{\rho} = \frac{1}{50} = 0.02$   
 $\Lambda_{eq} = \kappa \times V = \kappa \times \frac{1000}{\text{Normality}} = \frac{0.02 \times 1000}{0.5}$   
= 40 S cm<sup>2</sup> equiv.

**12.** (b) : Volume strength =  $5.6 \times$  Normality  $= 5.6 \times 1.5 = 8.4$ 

13. (b): For a cubic close packed structure, length of the side of unit cell is related to radius as,

$$r = \frac{a}{2\sqrt{2}}$$
  
 $a = r \times 2\sqrt{2} = 125 \times 2 \times 1.414 \text{ pm} = 353.5 \text{ pm}$ 

CHEMISTRY TODAY | MARCH '18

14. (b)

15. (b) : 
$$(CH_3)_2CO + HCN \longrightarrow CH_3 \longrightarrow CH_3 \longrightarrow CH_3$$
  

$$HOOC - C = CH_2 \xleftarrow{H_2SO_4}{Heat} CH_3 \longrightarrow CH_3 \longrightarrow CH_3$$

$$CH_3 \longrightarrow CH_3 \longrightarrow COOH$$

$$(Z) \longrightarrow CH_3 \longrightarrow COOH$$

CH

**16.** (b) : The photon capable of removing electron from first Bohr's orbit must possess energy

= 13.6 eV = 13.6 × 1.602 × 
$$10^{-19}$$
 J  
= 21.787 ×  $10^{-19}$  J

$$\therefore \quad E = \frac{nc}{\lambda}$$
21.787 × 10<sup>-19</sup> =  $\frac{6.625 \times 10^{-34} \times 3.0 \times 10^8}{\lambda}$ 

 $\therefore \quad \lambda = 912.24 \times 10^{-10} \text{ m} = 912.24 \text{ Å}$ This is longest  $\lambda$  because a photon having  $\lambda$  higher than

this will possess energy lesser than required, as  $E \propto \frac{1}{\lambda}$ . 17. (c) : The reaction proceeds via carbocation formation.

$$C_{6}H_{5}-C \stackrel{\frown}{=} CH_{2} \stackrel{+H^{+}}{\longrightarrow} C_{6}H_{5} \stackrel{I}{-C} \stackrel{H_{3}}{\longrightarrow} \stackrel{H_{3}C}{\longrightarrow} \stackrel{H_{3}C$$

**18.** (c) : Glucose is present in pyranose form.

**19.** (**d**) :  $A_{(s)} \rightleftharpoons 2B_{(g)} + 3C_{(g)}$ ∴  $K_c = [C]^3 [B]^2;$ 

If [C] becomes twice, let conc. of B becomes B', then  $K_c = [2C]^3 [B']^2$  or  $[C]^3 [B]^2 = [2C]^3 [B']^2$ 

$$\therefore \quad \frac{[B']}{[B]} = \sqrt{\frac{1}{8}} = \frac{1}{2\sqrt{2}}$$

20. (b)

**21.** (d) : Option (d) is a required condition for negative deviation along with  $\Delta V_{\text{mix}} = -\text{ve}$  and  $\Delta H_{\text{mix}} = -\text{ve}$ .

**22.** (b) : The electronic configurations of these elements are

$$V (Z = 23) : 1s^2 2s^2 2p^6 3s^2 3p^6 3d^3 4s^2$$

CHEMISTRY TODAY | MARCH '18

Cr (Z = 24): 
$$1s^2 2s^2 2p^6 3s^2 3p^6 3d^5 4s^1$$
  
Mn (Z = 25):  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^5 4s^2$   
Fe (Z = 26):  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^6 4s^2$ 

In the case of chromium, the second electron has to be removed from the half-filled d-shell which is more stable.

23. (a) : For the first order reaction,

$$k = \frac{2.303}{t} \log_{10} \frac{a}{(a-x)}$$

Let the initial amount is  $a \mod L^{-1}$ , then

after 
$$t = 100$$
 seconds,  $(a - x) = \frac{a}{3} \mod L^{-1}$   
 $\therefore \quad k = \frac{2.303}{100} \log_{10} \frac{a}{a/3} = \frac{2.303}{100} \log_{10} 3$   
 $= 10.988 \times 10^{-3} \sec^{-1}$ 

Let the time required to reduce the concentration to a/9 is  $t_1$ , then

$$t_1 = \frac{2.303}{10.988 \times 10^{-3}} \log_{10} \frac{a}{a/9} = 200 \sec \theta$$

24. (a)

25. (c) : It is not a good conductor of electricity.

**26.** (d) : 1-Alkynes react with ammoniacal solution of  $Cu_2Cl_2$  to form red precipitate of the corresponding copper alkynides.

But-1-yne reacts with ammoniacal  $Cu_2Cl_2$  as follows :  $CH_3 - CH_2 - C \equiv CH + Cu(NH_3)_2Cl$  $\longrightarrow CH_3 - CH_2 - C \equiv C^-Cu^+$ 

But, but-2-yne does not react with this reagent.

**27.** (d) : CO and  $CN^-$  are strong field ligands which force the electrons to pair up and thus, complex is generally diamagnetic. NH<sub>3</sub> is a weak field ligand so that electrons remain unpaired and complex is generally paramagnetic.

<b>28.</b> (d) : $ClO_3^-$	: 1 lone pair
XeF <sub>4</sub>	: 2 lone pairs
$SF_4$	: 1 lone pair
$I_3^-$	: 3 lone pairs

29. (c) : Iodoform - Antiseptic

Methyl salicylate - Pain balm

Diethyl ether - Anaesthetic

Hexachlorocyclohexane - Insecticide

**30.** (a) : Inert pair effect increases as we move down the group.

31. (c)



# Last-minute check on your **NEET** readiness





MTG's NEET Explorer helps students self-assess their readiness for success in NEET. Attempting the tests put together by MTG's experienced team of editors and experts strictly on the NEET pattern and matching difficulty levels, students can easily measure their preparedness for success. Order now!

### HIGHLIGHTS:

- 10 Model Test Papers based on latest NEET syllabus
- Last 15 years' solved test papers of AIPMT / NEET
- Includes NEET 2017 solved paper
- Detailed solutions for self-assessment and to practice time management



Available at all leading book shops throughout India. For more information or for help in placing your order: Call 0124-6601200 or email: info@mtg.in

\*Application to read QR codes required



Scan now with your smartphone or tablet\*

Visit www.mtg.in for latest offers and to buy online!



**32.** (b) : Higher the value of  $T_c$ , more easily the gas can be liquified.

**33.** (c) : The size of given metals decreases whereas ionization enthalpy increases from Ti to Fe. Hence, the metallic character of the metals decreases and therefore, basicity of oxides decreases from Ti to Fe.

**34.** (b) : Mass of substance taken = 0.316 g Mass of BaSO<sub>4</sub> formed = 0.466 g From stoichiometry, BaSO<sub>4</sub> = S  $_{233}$   $_{32}$ 

(: molecular mass of  $BaSO_4 = 137 + 32 + 64 = 233 \text{ g mol}^{-1}$ )

Then, mass of S in 0.466 g of  $BaSO_4 = \frac{0.466 \times 32}{233} g$ 

Percentage of S in the compound =  $\frac{0.466 \times 32}{233} \times \frac{100}{0.316}$ = 20.25 %



36. (b)

37. (b) : In highest oxidation states, transition metals cannot form cationic complexes. Also, they show acidic character because in highest oxidation state, they can only accept the electrons and form anionic complexes.38. (b)

**39.** (b) : 
$$\rho_{\text{NaCl}} = \frac{Z \times M}{a^3 \times N_A}$$
  
 $\therefore Z = 4$ , formula mass (*M*) = 58.5,  $a = 5.64 \times 10^{-8}$  cm

$$\therefore \quad \rho = \frac{4 \times 58.5}{6.023 \times 10^{23} \times (5.64 \times 10^{-8})^3} = 2.16 \text{ g cm}^{-3}$$

**40.** (a) : No. of electron pairs at the central atom = no. of atoms bonded to it + 1/2[group number of central atom – valency of the central atom ± no. of electrons] No. of electron pairs at the central atom

in NO<sub>3</sub><sup>-</sup> = 3 + 
$$\frac{1}{2}$$
[5-6+1] = 3 (*sp*<sup>2</sup> hybridisation).

No. of electron pairs at the central atom in

in 
$$H_3O^+ = 3 + \frac{1}{2}[6 - 3 - 1] = 4$$
 (*sp*<sup>3</sup> hybridisation).

**41.** (c) :  $\Delta G^{\circ} = -2.303 RT \log_{10} K$   $\log_{10} K = -\frac{\Delta G^{\circ}}{2.303 RT} = -\frac{(\Delta H^{\circ} - T\Delta S^{\circ})}{2.303 RT}$  $= -\frac{\Delta H^{\circ}}{2.303 RT} + \frac{\Delta S^{\circ}}{2.303 R}$ 

Comparing it with straight line equation, y = mx + c

we get, slope 
$$(m) = \frac{-\Delta H^{\circ}}{2.303 R}$$
  
and intercept  $(c) = \frac{\Delta S^{\circ}}{2.303 R}$ 

**42.** (a) : 2.83 B.M. implies two unpaired electrons according to the expression,  $\mu = \sqrt{n(n+2)}$  B.M. The species Ni<sup>2+</sup>, Ni<sup>2+</sup>, Ti<sup>4+</sup> and Co<sup>3+</sup> in the given complexes have 3*d*<sup>8</sup>, 3*d*<sup>8</sup>, 3*d*<sup>0</sup>, and 3*d*<sup>6</sup> electronic configurations, respectively. CN being a strong field ligand causes pairing of electrons thus, [Ni(CN)<sub>4</sub>]<sup>2-</sup> has zero unpaired electrons with *dsp*<sup>2</sup> hybridisation, while NH<sub>3</sub> being a weak field ligand, does not cause pairing of electrons thus, [Ni(NH<sub>3</sub>)<sub>6</sub>]<sup>2+</sup> has two unpaired electrons and 2.83 B.M. magnetic moment.

43. (c) : CaO + 2C 
$$\xrightarrow{\text{Heat}}$$
 CaC<sub>2</sub>  $\xrightarrow{\text{H}_2\text{O}}$  CH  $\equiv$  CH  
(A) (B)

**44. (b) :** Number of equivalents of gold deposited = number of equivalents of silver deposited

*i.e.*, 
$$\frac{W_{\text{gold}}}{E_{\text{gold}}} = \frac{W_{\text{silver}}}{E_{\text{silver}}}$$
$$E_{\text{gold}} = \frac{E_{\text{silver}} \times W_{\text{gold}}}{W_{\text{silver}}} = \frac{107.9 \times 1.314}{2.158} = 65.7$$
Equivalent mass = Atomic mass

Oxidation no. of Au in salt

Thus, ox. no. of Au = 
$$\frac{\text{Atomic mass}}{E_{\text{gold}}} = \frac{197}{65.7} = 3$$
  
**45. (b)** :  $MX \longrightarrow M^+ + X^-$   
 $H_2O \longrightarrow H^+ + OH^-$ 

At cathode :  $H^+$  ions will get reduced as the standard reduction potential of  $M^+$  ions is negative (less than that of  $H^+$ ).

$$2H^+ + 2e^- \longrightarrow H_2$$

**At anode :** The species having low value of standard reduction potential are oxidised. Hence, the reaction at anode is

$$2X^- \longrightarrow X_2 + 2e^-$$

 $\therefore$  The products obtained are H<sub>2</sub> at cathode and X<sub>2</sub> at anode.

CHEMISTRY TODAY | MARCH '18



Dear students! Thank you all a lot for the feedback that you have given for the last article. This is the last part of Organometallic Compounds. Hopefully, this will fulfil the requirement of COMPETITIVE EXAMINATIONS you are going to face. Moreover, some more problems will be given in upcoming issues. Do read the basics of every chapter very carefully. Do take care, all the best!

\*Arunava Sarkar 🛓

### A BRIEF OVERVIEW OF ORGANOMETALLIC REAGENTS AND ORGANOMETALLIC CHEMISTRY

### **GRIGNARD REAGENTS**

### **O** Reaction with nitriles

Remember the basic idea here which is, whereever –CN group is connected, remove it from there and add –COCH<sub>3</sub>  $\begin{pmatrix} \text{or} - C - R \\ \parallel \\ O \end{pmatrix}$ .



[-CH<sub>3</sub> or -*R* comes from Grignard reagent]

### **O** Reaction with epoxide



If you use substituted epoxide, then you can also get 2° alcohol.



### **O** Reaction with acid anhydride



### **O** Reaction with alkyl halide

Reaction between Grignard reagent and an alkyl halide is often carried out very slowly as Grignard reagent is prepared from alkyl halide. So, the reaction between Grignard reagent and alkyl halide is carried out in presence of transition metal catalyst, specifically cobalt chloride (CoCl<sub>2</sub>). This kind of transition metal catalysed reaction is known as Kharasch reaction. For example,

CH<sub>3</sub>MgBr + CH<sub>3</sub>CH<sub>2</sub>Br 
$$\xrightarrow{\text{CoCl}_2}$$
 CH<sub>3</sub>CH<sub>2</sub>CH<sub>3</sub>  
(≈ 68%-70%)

It has been found that a reaction is possible in between Grignard reagent and  $\alpha$ -haloketone. For example,

$$C_6H_5COCH_2Cl + C_6H_5MgBr \xrightarrow{(1) dry ether, -28 °C} (2) ice cold HCl C_6H_5COCH_2C_6H_5$$

\*Institute of Chemistry (IOC)- Asansol, Durgapur, Dhanbad, Burdwan, Kolkata, Jamshedpur, Bokaro, Patna

CHEMISTRY TODAY | MARCH '18

### **ORGANOLITHIUM COMPOUNDS**

Obviously, organolithium compounds are characterised by a C – Li bond. Li is less electronegative than carbon and hence, the polarisation of carbon-lithium bond is shown as below :

$$\geq^{\underline{\delta}}_{C} - Li$$

Organolithium compounds are more reactive than Grignard reagents, however the nature of reaction remains the same in most of the cases and in some cases distinctly different.

Preparation of organolithium compounds

#### • From alkyl or aryl halides

 $R \rightarrow X + 2Li \xrightarrow{low temperature}_{organic solvent} R \rightarrow Li + LiX$ inert atmosphere (Ar)

Order of reactivity : R - I > R - Br > R - Cl

### Halogen-metal exchange

$$R - X + R'Li \longrightarrow R - Li + R' - X$$
  
e.g.,  
Br  
$$+ n - C_4 H_9 Li \longrightarrow O + n - C_4 H_9 Br$$

An important reaction to show the difference in reactivity between Grignard reagent and organolithium compound before getting into discussion on the reactions shown by organolithium compounds is :



So, organolithium compounds are more reactive than organomagnesium compounds.

#### Most useful reactions of organolithium compounds

# • Reaction with carbonyl compounds (with the focus on conjugated carbonyl compounds) Let us start the discussion with the similarities in the reactions with Grignard reagent.

Grignard reagent reacts with formaldehyde to give 1° alcohol (after hydrolysis), acetaldehyde to give 2° alcohol (after hydrolysis), acetone to give 3° alcohol (after hydrolysis). Same goes with organolithium compounds.



For the third case, there is a well known advantage of using organolithium compounds over Grignard reagents that is Grignard reagents do not react with hindered ketones but organolithium compounds do. Here is the best and most used example :

$$(CH_{3})_{3}C - C - C(CH_{3})_{3} \xrightarrow{(CH_{3})_{3}CLi} [(CH_{3})_{3}C]_{3}COLi$$
  
di-*tert*-butyl ketone  
$$\downarrow H_{3}O^{+} [(CH_{3})_{3}C]_{3}COH$$
  
(Highly hindered  
3° alcohol)

Now, let us take the case of  $\alpha$ , $\beta$ -unsaturated carbonyl compounds. We have seen that with  $\alpha$ ,  $\beta$ -unsaturated compounds, Grignard reagent can give either 1,2-addition or 1,4-addition product depending on the overall steric hindrance whereas organolithium compounds on the other hand, give exclusively 1,2-addition products.



[The reason for this is that organolithium compounds are aggressive nucleophiles.]



However, lithium dialkyl cuprates give exclusively 1,4-addition product.

• Reaction with carbon dioxide



So, here is an important difference with the Grignard reagent. In case of Grignard reagent, the reaction stops at stage 1 i.e., 1 mole addition of Grignard reagent but in case of organolithium compounds, the reaction goes upto stage 2 i.e., 2 moles addition of organolithium compound.

### Electrophilic displacement

R - X + R'Li  $\longrightarrow$  RLi + R' - X

A specific example of this kind can be given :

 $CH_2 = CHBr + nC_4H_9Li \longrightarrow$ Vinyl bromide *n*-Butyl lithium

 $CH_2 = CHLi + nC_4H_9Br$ 

### O Nucleophilic displacement

Here, a similar reaction as happens in Wurtz reaction takes place.

$$R \xrightarrow{f} X + \overbrace{R'} \stackrel{:}{\xrightarrow{k'}} Li^+ \xrightarrow{S_N^2} R - R' + LiX$$

It is to be remembered that in case of substitution in aromatic rings, elimination-addition reaction takes place. Take the following example,



1-Phenylnaphthalene

2-Phenylnaphthalene





ی چ

# AVAILABLE BOUND VOLUMES

PHYSICS for you 2016		Physics For You 2016 (January - December)	₹ 325 12 issues
2016	Mg	Mathematics Today 2016 (January - December)	₹ 325 12 issues
MATHEMATICS	BIOLOGY	Biology Today 2016 (January - June)	₹175 6 issues
	2016	Chemistry Today 2016 (January - June)	₹175 6 issues
2016 buy online at w	ww.mtg.in	Mathematics Today 2013 (January - December)	₹ 300 12 issues

### of your favourite magazines

How to order : Send money by demand draft/money order. Demand Draft should be drawn in favour of MTG Learning Media (P) Ltd. Mention the volume you require along with your name and address.

### Add ₹ 60 as postal charges

Older issues can be accessed on **digital.mtg.in** in digital form.

### Mail your order to :

Circulation Manager, MTG Learning Media (P) Ltd. Plot No. 99, Sector 44 Institutional Area, Gurgaon, (HR) Tel.: (0124) 6601200 E-mail : info@mtg.in Web : www.mtg.in





method)

• % of O =  $\frac{5 \times 16}{2 \times 127} \times \frac{\text{mass of I}_2 \text{ formed}}{\text{mass of compound taken}} \times 100$ 

• Stability of carbocations  $\propto +I$ -effect  $\propto \frac{1}{-I$ -effect  $\propto R$ -effect  $\propto \frac{1}{-R$ -effect  $\mathrm{Ph_3C}^+ > \mathrm{Ph_2CH}^+ > \mathrm{PhCH_2}^+ > \mathrm{Allyl} > 3^\circ > 2^\circ > 1^\circ > \overset{+}{\mathrm{CH}_3}$ 

### - p-type semiconductors : Group 14 elements doped with group 13 elements, holes increase conductivity.

 $p_A^{\circ} - p_A / p_A^{\circ} = ix_B; \Delta T_b = iK_b m; \Delta T_f = iK_f m; \pi = iCRT$ 

Degree of dissociation ( $\alpha$ ) = i - 1/n - 1; i > 1

• Modified colligative properties :



### HYDROLYSIS AND HYDRATION

**NOTE** : In this article equations are intensionally not balanced.

If you are ready to consider hydration along with hydrolysis, you will be amazed to see a large number of important chemical reactions of your syllabus comes under this single banner.

In hydrolysis, bonds break during the reaction while in hydration simply water is added. In hydrolysis, you watch more than one product while in hydration, there is only one product unless a disproportionation takes place.

 $CCl_4 + H_2O \longrightarrow$  No hydrolysis ordinarily

Carbon fails to expand its octet to receive the donation of electron pair from  $H_2O$  to form required intermediate. However, carbon uses the third energy shell to expand its octet.

$$CCl_4 + H_2O \xrightarrow{\Delta} COCl_2 + HCl$$
  
siceam SiCl\_4 + H\_2O  $\longrightarrow$  Si(OH)\_4 + HCl  
SiF\_4 + H\_2O  $\longrightarrow$  H\_2SiO\_3 + HF

But HF subsequently reacts with SiF<sub>4</sub>.

 $SiF_4 + 2HF \longrightarrow H_2SiF_6$ 

 $SiH_4$  is stable to water but rapidly hydrolysed by alkali.

 $SiH_4 + OH^- \longrightarrow SiO_4^{4-} + H_2$ 

In SiH<sub>4</sub>, the hydrogen carries a partial negative charge and in OH<sup>-</sup>, it carries a partial positive charge. This facilitates the release of H<sub>2</sub>.

 $GeH_4 + OH^- (30\%) \longrightarrow$  No hydrolysis Polarity of H in both  $GeH_4$  and  $OH^-$  is positive.

 $BF_3 + H_2O \longrightarrow H^+[BF_3(OH)]^-$ 

No hydrolysis, rather an acid-base reaction takes place. BCl<sub>3</sub> + H<sub>2</sub>O  $\longrightarrow$  B(OH)<sub>3</sub> + HCl

The general rule is the central atom produces the oxyacid maintaining its oxidation state while the peripheral atom produces the hydra-acid.

$$AlCl_3 + H_2O \longrightarrow Al(OH)_3 + HCl$$

Again, NF<sub>3</sub> + H<sub>2</sub>O  $\longrightarrow$  No hydrolysis

N – F bond is too strong to break.

 $NCl_3 + H_2O \longrightarrow NH_3 + HOCl$ 

(not HNO<sub>2</sub> and HCl)

Nitrogen has no room to accept electrons. Therefore, it is the Cl that accepts donated electrons.

 $PF_5 + H_2O \longrightarrow No hydrolysis$ 

P − F bond is again too strong to cleave.  
PCl<sub>3</sub> + H<sub>2</sub>O 
$$\longrightarrow$$
 H<sub>3</sub>PO<sub>3</sub> + HCl  
PCl<sub>5</sub> + H<sub>2</sub>O  $\longrightarrow$  H<sub>3</sub>PO<sub>4</sub> + HCl  
SF<sub>6</sub> + H<sub>2</sub>O  $\longrightarrow$  No hydrolysis

Steric factor is the primary reason of the failure as there is no room for  $H_2O$  to approach cental S atom, which is crowded by six fluorine atoms.

$$\begin{array}{ccc} \mathrm{SCl}_6 + \mathrm{H}_2\mathrm{O} & & \swarrow & \mathrm{S(OH)}_6 + \mathrm{HCl} \\ \mathrm{does \ not} & & \mathrm{does \ not} \\ \mathrm{exist} & & \mathrm{exist} & \mathrm{either} \end{array}$$

The lower electronegativity of six Cl<sup>-</sup> or six OH<sup>-</sup> is responsible for non-existence of these compounds.

$$BiCl_{3} + H_{2}O \longrightarrow BiOCl + HCl$$
White turbidity
$$SbCl_{3} + H_{2}O \longrightarrow SbOCl + HCl$$
White turbidity

There are  $BiO^+$  and  $SbO^+$  ions formed due to high positive charge density of  $Bi^{3+}$  and  $Sb^{3+}$  ions.

$$BrF_5 + H_2O \longrightarrow HBrO_3 + HF$$

\_\_\_\_

Similarly,

$$CH_{3} \xrightarrow{O} CH_{3} \xrightarrow{C} C=O + HOCH_{2}CH_{2}OH$$

$$CH_{3} \xrightarrow{O} CH_{3} \xrightarrow{C} OH^{-} No hydrolysis$$

$$R - C - NH_{2} \xrightarrow{H_{3}O^{+}} RCOOH + NH_{4}^{+}$$

$$R - C - NH_{2} \xrightarrow{OH^{-}} RCOO^{-} + NH_{3}$$
(Faster and easier)

Mukul C. Ray, Odisha

$$R - C = N \xrightarrow{H_{3}O^{+}} RCOOH + R'OH (reversible) (rev$$

OH<sup>-</sup> cannot attack either N or C.

$$R - O - R' \xrightarrow{H_3O^+} R - OH + R' - OH$$

 $RO^-$  or  $R'O^-$  both are bad leaving groups.

$$CH_{3} \xrightarrow[C]{H_{3}} CH_{3} \xrightarrow[C]{H_{2}O} (CH_{3})_{3}C \xrightarrow[C]{OH(S_{N}^{1})} CH_{3} \xrightarrow[C]{CH_{3}} CH_{2} \xrightarrow[C]{CH_{3}} CH_{2} \xrightarrow[C]{CH_{3}} CH_{3} \xrightarrow[C]{CH_{3}} No hydrolysis, rather elimination$$

It is not surprising that sometimes hydrolysis reactions give a twist.

 $XeF_4 + H_2O \longrightarrow Xe + HF + O_2 + XeO_3$ It is a clear cause of disproportionation during hydrolysis. Whereas, the other two fluorides of Xe behave in different ways :

$$XeF_{2} + H_{2}O \longrightarrow Xe + HF + O_{2}$$
$$XeF_{6} + H_{2}O \longrightarrow XeOF_{4} + HF$$
$$XeOF_{4} + H_{2}O \longrightarrow XeO_{3} + HF$$

Before we close, check the following two reactions :

$$\begin{array}{c} CH_{3} \\ H \\ C = NH \xrightarrow{H_{2}O} CH_{3} \\ H \\ C = O + NH_{3} \\ R - C - Cl \xrightarrow{H_{2}O} R - C - OH + HCl \end{array}$$

No catalyst even required for these two organic reactions. We shall be checking hydration in the next episode.

ی چ



### Who can participate

If you have taken any of the exams given below and possess plenty of grey cells, photographic memory then you are the right candidate for this contest. All you have to do is write down as many questions (with all choices) you can remember, neatly on a paper with name of the exam, your name, address, age, your photograph and mail them to us.



PMT: AIIMS, JIPMER, SRMJEEE .... etc.

(2)

Engineering: VITEEE, UPSEE, AMU, SRMJEEE, BITSAT, COMED-K.... etc.

### The Benefits

Plenty! Each complete question with answer will make you richer by Rs. 100\*. More the questions, the merrier it will be. We will make you famous by publishing your name (photo if possible). Also you can derive psychological satisfaction from the fact that your questions will benefit thousands of readers.

### And Lastly The Pitfalls

Don't send incomplete question. Our panel of experts will cross-check your questions. You have to send it within a month of giving the particular exam.

### Mail to : The Editor, MTG Learning Media Pvt. Ltd. Plot 99, Sector 44

Institutional Area, Gurgaon – 122003(HR) Tel. :0124-6601200

#### \*Conditions apply

- Payment will be made after the MCOs are published.
- Kindly note that each question should be complete.
- Payment will be made only for complete questions.
- Preference will be given to the reader sending the maximum complete and correct questions. Other conditions apply. The decision of the Editor, MTG shall be final and binding.



# CHEMISTRY MUSING

**SOLUTION SET 55** 

1. (d): 
$$cis X \xrightarrow{k_f} trans X$$
  
Initial  $a$   $0$   
at time  $t$   $a - x$   $x$   
at eqn  $a - x_e$   $x_e$   
 $K_{(eq)} = \frac{k_f}{k_b}; k_b = \frac{3 \times 10^{-4}}{0.1} = 3 \times 10^{-3}$   
As we know  $(k_f + k_b) = \frac{1}{t} \ln\left(\frac{x_e}{x_e - x}\right)$   
Given,  $x = \frac{x_e}{2}$   $\therefore$   $(k_f + k_b) = \frac{1}{t} \ln 2$   
or  $(3 \times 10^{-4} + 3 \times 10^{-3}) = \frac{0.693}{t}$   
 $\therefore$   $t = 210$  sec  
2. (d): Average  $C_{v,m} = \frac{n_1 C_{v,m_1} + n_2 C_{v,m_2}}{n_1 + n_2} = 2R$  ...(i)

For adiabatic process, dU = dw

$$n_1 C_{\nu,m_1} dT + n_2 C_{\nu,m_2} dT = -(n_1 RT + n_2 RT) \times \frac{d\nu}{V}$$
$$\frac{dT}{T} = -\frac{R}{\text{Average } C_{\nu,m}} \left(\frac{dV}{V}\right)$$

On substituting the value of average  $C_{v, m}$  from eqn (i), then on integration, we get

$$\ln \frac{T_2}{T_1} = -\frac{1}{2} \ln \left( \frac{V_2}{V_1} \right) \implies T_2 = 320 \times \left( \frac{1}{4} \right)^{1/2} = 160 \text{ K}$$
$$\Delta U = (n_1 C_{\nu, m_1} + n_2 C_{\nu, m_2}) \Delta T$$
$$= \left( 1 \times 3R + 2 \times \frac{3}{2} R \right) (160 - 320) = -960 R$$
$$3. \text{ (b):}$$

$$\text{Delocalised lone pair (5) NH}_2 = \frac{1}{2} \text{ Localised lone pair (4) } (3)$$

(2)

CH<sub>3</sub>



Delocalised

lone pair

Delocalised

lone pair

1 7 7



**4.** (b): With glycinato ligand,  $Pt^{2+}$  and  $Pt^{4+}$  both show isomerism but with ethylene diamine,  $Pt^{2+}$  does not show but  $Pt^{4+}$  show isomerism. In  $[Pt(en)_2]^{2+}$ ,  $Pt^{2+}$ is  $dsp^2$  hybrid in square planar splitting  $d_{x^2-y^2}$  has higher energy in comparison of  $d_{z^2}$  but in  $[Pt(en)_3]^{4+}$ ,  $d_{x^2-y^2}$  and  $d_{z^2}$  are at same energy level.

5. (a): 
$$\overline{O} - S - O^{-} + {}^{35}_{16}S \xrightarrow{\text{Boil}} \overline{O} - {}^{\text{II}}_{\text{II}} - \overline{O}$$
  
 ${}^{35}_{\text{S}} \xrightarrow{O} O^{-} - {}^{\text{S}}_{\text{HCl}} - O^{-} \xrightarrow{\Delta}_{\text{HCl}} SO_{2} + {}^{35}_{16}S + H_{2}O$   
 $O$ 

6. (b): For a constant volume process under adiabatic conditions,

$$\Delta U = \Delta U_{\text{heating}} + \Delta U_{298 \text{ K}} = 0$$
  
Hence,  $\Delta U_{\text{heating}} = -\Delta U_{298 \text{ K}}$ 
$$= -\int_{298 \text{ K}}^{T_f} \Sigma n C_v dT = -240.6 \text{ kJ} \qquad \dots(i)$$

Form IV				
1.	Place of Publication :	New Delhi		
2.	Periodicity of its Publication :	Monthly		
3.	Printer's Name :	HT Media Ltd.		
3a.	Publisher's Name :	MTG Learning Media Pvt. Ltd.		
	Nationality :	Indian		
	Address :	406, Taj Apartment,		
		New Delhi - 110029		
4.	Editor's Name :	Anil Ahlawat		
	Nationality :	Indian		
	Address :	19, National Media		
		Centre, Gurgaon,		
		Haryana - 122002		
5.	Name and address of individuals who :	Mahabir Singh Ahlawat		
	own the newspapers and partners or	64, National Media Centre,		
	shareholders holding more than one	Nathupur, Gurgaon		
	percent of the total capital :	Krishna Devi		
		64, National Media Centre,		
		Nathupur, Gurgaon		
	:	Anil Ahlawat & Sons		
		19, National Media Centre,		
		Nathupur, Gurgaon		
	:	Anil Ahlawat		
		19, National Media Centre,		
l		Nathupur, Gurgaon		
I, N	ianabir Singh, authorised signatory for M	IG Learning Media Pvt. Ltd. hereby		
deo bel	clare that particulars given above are true ief.	e to the best of my knowledge and		
		For MTG Learning Media Pvt. Ltd.		
		Mahabir Singh		
		Director		



H-l

Localised.

lone pair

Since, 2 moles unreacted N<sub>2</sub> are associated with 1/2 mole of O<sub>2</sub> (:: Ratio of N<sub>2</sub>: O<sub>2</sub> in air = 4:1), we have  $\Sigma nC_v = C_v(H_2O, g) + 2C_v(N_2, g)$ 

$$= (39.1 + 2 \times 26.4) \text{ J K}^{-1} = 91.9 \text{ J K}^{-1}$$

Hence, from eqn. (i), on integrating, we have 91.9 J K<sup>-1</sup> ( $T_f$  – 298) = 240, 600 J

 $T_f - 298 = 240, 600 \text{ J/91.9 J K}^{-1} = 2618 \text{ K}$ 

 $T_f = (2618 + 298) \text{ K} = 2916 \text{ K}$ 

7. (c) : In case of *G* as EWG or EWG attached to *ortho*or *para*-position with respect to group *G*,  $\sigma$ -complex (arenium ion) will be destabilised leading to decrease of rate of reaction for *ipso* substitution.



SO<sub>3</sub> and CO<sub>2</sub> are good leaving groups as formation of

C = C and S = O bonds provides thermodynamic stability.





CHEMISTRY TODAY | MARCH '18



Maximise your chance of success, and high rank in NEET, JEE (Main and Advanced) by reading this column. This specially designed column is updated year after year by a panel of highly qualified teaching experts well-tuned to the requirements of these Entrance Tests.

Unit 9

# **Principles Related to Practical Chemistry**

Practical work in labs improves the scientific knowledge and understanding of students, as well as providing opportunities for working scientifically and developing hands on skills.

Much of forensic science, medicine, pharmaceutical research and many chemical manufacturing processes, rely on accurate techniques and observations.

### QUANTITATIVE ESTIMATION

A quantitative analysis is one in which the amount or concentration of a particular species in a sample is determined accurately and precisely.

### Titration

The process of addition of the known solution from the burette to the measured volume of solution of the substance to be estimated until the reaction between the two is just complete.



These can be accurately weighed

and their solutions are not to be

standardised before use e.g., oxalic

### Unknown Solution

The solution consisting of substance to be estimated is termed as titrate or unknown solution.

### Acid-base titration

- When the strength of an acid is determined with the help of a standard solution of base, it is known as acidimetry.
- When the strength of a base is determined with the help of a standard solution of an acid, it is known as alkalimetry.

acid, potassium dichromate, etc. hydroxide, hydrochloric acid, etc. These titrations involve neutralisation of an acid

The solutions of these reagents are to be

standardised before use as these cannot

be weighed accurately e.g., sodium

with an alkali.  

$$HA + BOH \longrightarrow BA + H_2O$$
  
Acid Alkali Salt Water

A substance which helps in physical detection of completion of the titration is called indicator.



### Indicators

### • External Indicators

These indicators are not added in the reaction mixture. They are used outside the system *e.g.*, potassium ferricyanide is used in titration of Mohr's salt against potassium dichromate.

### Internal Indicators

These indicators are added in the reaction mixture *e.g.*, in acid-base titration, methyl orange, methyl red and phenolphthalein.

### Self Indicators

These indicators themselves show colour change at the end of titration *e.g.*, in titration of oxalic acid against  $KMnO_4$ ,  $KMnO_4$  acts as self indicator.

### How to choose a suitable indicator :

Indicator in a titration is selected on the basis of pH range at end point.

		Colour of indicator		
Indicator	pH-range	Acid	Alkaline	
Phenolphthalein	8.3 – 10	Colourless	Pink	
Methyl orange	3.2 - 4.4	Red	Yellow	
Methyl red	4.4 - 6.5	Red	Yellow	
Litmus	5.0 - 8.0	Red	Blue	
Phenol red	6.8 - 8.4	Yellow	Red	





### **Determination of Strength of NaOH**

**Principle :** Strength of NaOH solution can be determined by titrating it against standard solution of oxalic acid.

 $\begin{array}{c} \text{COOH} \\ | \\ \text{COOH} \end{array} + 2\text{NaOH} \rightarrow \begin{array}{c} \text{COONa} \\ | \\ \text{COONa} \end{array} + 2\text{H}_2\text{O} \\ \text{COONa} \end{array}$ 

Indicator : Phenolphthalein Preparation of standard solution :

Oxalic acid  $\begin{pmatrix} \text{COOH} \\ | \\ \text{COOH} \end{pmatrix}$ Equivalent weight =  $\frac{126}{2} = 63$ 

For preparing 250 mL solution of N/10 oxalic acid,

weight required = 
$$\frac{63 \times \frac{1}{10} \times 250}{1000} = 1.575 \text{ g}$$

Thus, for preparing N/10 oxalic acid, take 1.575 g of oxalic acid and make it upto 250 mL using distilled water.

### **Calculation :**

 $N_1V_1 = N_2V_2$ (Oxalic acid) (Caustic soda)  $N_2 = \frac{N_1V_1}{V_2}$ 

Strength of caustic soda = Normality × Equivalent weight =  $(N_2 \times 40)$  g/L

### **Determination of Strength of HCI**

**Principle :** When hydrochloric acid is titrated against Na<sub>2</sub>CO<sub>3</sub> solution, strength of HCl can be determined. Na<sub>2</sub>CO<sub>3</sub> + 2HCl  $\rightarrow$  2NaCl + H<sub>2</sub>O + CO<sub>2</sub> $\uparrow$ **Indicator :** Methyl orange

Preparation of standard solution :

Eq. weight  $=\frac{106}{2}=53$ 

For preparing 250 mL solution of N/10 Na<sub>2</sub>CO<sub>3</sub>,

weight required = 
$$\frac{53 \times \frac{1}{10} \times 250}{1000} = 1.325 \text{ g}$$

### **Calculation** :

 $\begin{array}{cc} N_1 V_1 = N_2 V_2 & ; & N_2 = \frac{N_1 V_1}{V_2} \\ (\mathrm{Na}_2 \mathrm{CO}_3) & (\mathrm{HCl}) & \end{array}$ 

Strength of given HCl = Normality × Eq. weight of HCl =  $(N_2 \times 36.5)$  g/L



CHEMISTRY TODAY | MARCH '18

### **QUALITATIVE ANALYSIS**

Qualitative analysis deals with the identification of various constituents present in a given material. For example, zinc blende contains zinc and sulphur in the form of  $Zn^{2+}$  and  $S^{2-}$ ions.

### **Preliminary Tests**

### Flame Test :

$$\begin{array}{c} & & \text{Greenish}\\ \text{Cu}^{2+} \xleftarrow{\text{blue}}\\ \text{K}^{+} \xleftarrow{\text{Lilac}}\\ \text{Golden}\\ \text{Na}^{+} \xleftarrow{\text{of the Bunsen burner}}\\ \text{Na}^{+} \xleftarrow{\text{of the Bunsen burner}} \\ & \text{of the Bunsen burner} \\ & \text{Sr}^{2+} \end{array}$$

### Borax Bead Test :

• Borax is heated on a loop of Pt wire, colourless, glassy bead of sodium metaborate and boric anhydride is formed.

$$Na_2B_4O_7 \cdot 10H_2O \xrightarrow{\Delta} Na_2B_4O_7 \xrightarrow{\Delta} 2Na$$

 $\underbrace{\text{MaBO}_2 + \text{B}_2\text{O}_3}_{\text{Glassy bead}}$ 

• Coloured salts are then heated on the glassy bead, coloured metaborate is formed in the oxidising flame.

Colour of bead in oxidising flame	Ion indicated
Green in hot, blue in cold	Copper
Pinkish violet in both hot and cold	Manganese
Yellowish brown in hot and pale yellow in cold	Iron
Brown in hot and pale brown in cold	Nickel

### **Acidic Radicals**

### Anions are acidic radicals

First group	Second	Third
$CO_2^{2-}, S^{2-}, SO_2^{2-}$	group	group
$CH_{2}COO^{-}$	Br <sup>−</sup> , Cl <sup>−</sup> , I <sup>−</sup>	$SO_4^{2-}, PO_4^{3-}$
$S_{2}O_{2}^{2-}$ , NO <sub>2</sub>	$NO_3^-, C_2O_4^{2-}$	Reagent used :
Reagent used :	Reagent used :	dil.H <sub>2</sub> SO <sub>4</sub> /
dil. HCl	conc. $H_2SO_4$	conc. $H_2SO_4$



# The most comprehensive **question bank books** that you cannot afford to ignore



30 Years' Physics, Chemistry & Biology contain not only chapterwise questions that have appeared over the last 30 years in NEET/AIPMT, but also full solutions, that too by experts. Needless to say, these question banks are essential for any student to compete successfully in NEET.

### HIGHLIGHTS:

- Chapterwise questions of last 30 years' (2017-1988) of NEET/AIPMT
- Chapterwise segregation of questions to help you assess the level of effort required to succeed
- An unmatched question bank series with close to 1,000 pages having detailed solutions by experts
- Fully solved questions of NEET 2017 included



Scan now with your smartphone or tablet\*



Available at all leading book shops throughout India. For more information or for help in placing your order: Call 0124-6601200 or email info@mtg.in \*Application to read QR codes required Visit www.mtg.in for latest offers and to buy online!

## iranchembook.ir/edu





### **Second Group :** Salt + conc. $H_2SO_4$

Effervescence or evolution of gases indicates the presence of group II acidic radicals.

Gas	Radical	<b>Observations and Reactions</b>
HCl (Colourless gas, pungent smell)	Chloride (Cl⁻)	$\begin{split} \text{NaCl} + \text{H}_2\text{SO}_4 & \rightarrow \text{NaHSO}_4 + \text{HCl}^{\uparrow} \\ \text{Salt} \\ \text{NH}_4\text{OH} + \text{HCl} & \rightarrow \text{NH}_4\text{Cl}^{\uparrow} + \text{H}_2\text{O} \\ & \text{White dense} \\ \text{fumes} \end{split}$
Br <sub>2</sub> (Brown fumes)	Bromide (Br <sup>-</sup> )	$NaBr + H_2SO_4 \rightarrow NaHSO_4 + HBr$ $2HBr + H_2SO_4 \rightarrow Br_2\uparrow + 2H_2O + SO_2$ Brown
I <sub>2</sub> (Deep violet gas)	Iodide (I <sup>-</sup> )	$2\text{KI} + 2\text{H}_2\text{SO}_4 \rightarrow 2\text{KHSO}_4 + 2\text{HI}$ $2\text{HI} + \text{H}_2\text{SO}_4 \rightarrow \text{I}_2^{\uparrow} + \text{SO}_2 + 2\text{H}_2\text{O}$ Violet
NO <sub>2</sub> (Light brown gas, pungent smell)	Nitrate (NO <sub>3</sub> <sup>-</sup> )	$NaNO_{3} + H_{2}SO_{4} \rightarrow NaHSO_{4} + HNO_{3}$ $4HNO_{3} \rightarrow 2H_{2}O + 4NO_{2}\uparrow + O_{2}\uparrow$ $Light brown$ fumes





Determination of sulphate by conductometric titration!

Titrimetric methods are much faster, but a good indicator has not been found for sulphate titrations. However, because the relative concentrations of ions in solution change during titration sequence, monitoring conductivity during the progress of a titration produces a signal which can be used to indicate equivalence point. A conductometric titration protocol has been developed which is relatively rapid, inexpensive and can produce accurate results.



### iranchembook.ir/edu

### **Confirmatory Tests for Group II**



Third Group : These anions are identified by their characteristic chemical reactions.



### **Basic Radicals**

Cations are the basic radicals.

Group	Group reagent	Cations	Form of ppt.
Ι	dil. HCl	$Pb^{2+}, Ag^+, Hg_2^{2+}$	Chlorides
II	dil. HCl + H <sub>2</sub> S gas	Pb <sup>2+</sup> , Hg <sup>2+</sup> , Cu <sup>2+</sup> , Cd <sup>2+</sup> , Bi <sup>3+</sup> ,	Sulphides
		Sb <sup>3+</sup> , As <sup>3+</sup> , Sn <sup>2+</sup> /Sn <sup>4+</sup>	

III	$NH_4Cl + NH_4OH$	Fe <sup>3+</sup> , Al <sup>3+</sup> , Cr <sup>3+</sup>	Hydroxides
IV	$NH_4Cl + NH_4OH + H_2S$ gas	Zn <sup>2+</sup> , Mn <sup>2+</sup> , Co <sup>2+</sup> , Ni <sup>2+</sup>	Sulphides
V	$(NH_4)_2CO_3 + NH_4OH + NH_4Cl$	Ca <sup>2+</sup> , Sr <sup>2+</sup> , Ba <sup>2+</sup>	Carbonates
VI	$Na_2HPO_4 + NH_4OH$	Mg <sup>2+</sup>	Phosphates

**Analysis of Basic Radicals :** Zero group  $[NH_4^+]$  : To the salt, add NaOH solution and heat. If ammonia gas evolves,  $NH_4^+$  is present.

Pass the gas through Nessler's reagent  $\rightarrow$  Brown ppt.  $\rightarrow$  NH<sub>4</sub><sup>+</sup> confirmed.







### Detection of N, S, Cl in organic compounds

### Lassaigne's extract :

A small pellet of metallic sodium together with a little amount of the substance is heated to red hot in an ignition tube. It is then suddenly plunged into about 10 mL of distilled water in a China dish. The mixture is boiled well and filtered. Filtrate is known as Lassaigne's extract (L.E.).

Element	Detection	Confirmatory test	Reactions
Nitrogen	Lassaigne's extract (L.E.)	L.E. + FeSO <sub>4</sub> + NaOH, boil	$FeSO_4 + 2NaOH \longrightarrow Fe(OH)_2$
	$Na + C + N \xrightarrow{\Delta} NaCN$	and cool + FeCl <sub>3</sub> + conc. HCl	$+ Na_2SO_4$
	(L.E.)	Gives blue or green colour.	$Fe(OH)_2 + 6NaCN \longrightarrow$
			$Na_4[Fe(CN)_6] + 2NaOH$
			$Na_4[Fe(CN)_6] + FeCl_3 \xrightarrow{HCl} \rightarrow$
			$NaFe[Fe(CN)_6] + 3NaCl$ Prussian blue
			or $3Na_4[Fe(CN)_6] + 4FeCl_3 \longrightarrow$
			$Fe_4[Fe(CN)_6]_3 + 12NaCl$ Prussian blue

CHEMISTRY TODAY | MARCH '18

Sulphur	$2Na + S \xrightarrow{\Delta} Na_2S$ (L.E.)	<ul> <li>(i) L.E. + sodium nitroprusside A deep violet colour.</li> <li>(ii) L.E. + CH<sub>3</sub>COOH + (CH<sub>3</sub>COO)<sub>2</sub>Pb Gives a black ppt.</li> </ul>	(i) $Na_2S + Na_2[Fe(CN)_5NO] \longrightarrow$ Sodium nitroprusside $Na_4[Fe(CN)_5NOS]$ Deep violet (ii) $Na_2S + (CH_3COO)_2Pb \xrightarrow{CH_3COOH} PbS \downarrow + 2CH_3COONa$ Black ppt.
Halogens	$Na + X \xrightarrow{\Delta} NaX$ (L.E.) $(X = Cl, Br, I)$	L.E. + $HNO_3 + AgNO_3$ (i) White ppt. soluble in aq. $NH_3$ (or $NH_4OH$ ) confirms Cl. (ii) Pale yellow ppt. partially soluble in aq. $NH_3$ (or $NH_4OH$ ) confirms Br. (iii) Yellow ppt. insoluble in aq. $NH_3$ (or $NH_4OH$ ) confirms I.	$NaX + AgNO_{3} \xrightarrow{HNO_{3}} AgX \downarrow + NaNO_{3}$ White ppt. $AgCl + 2NH_{4}OH_{(aq.)} \longrightarrow$ $[Ag(NH_{3})_{2}]Cl + 2H_{2}O$ Soluble
Nitrogen and sulphur together	$Na + C + N + S \xrightarrow{\Delta}$ NaSCN (L.E.) Sodium thiocyanate	As in test for nitrogen; instead of green or blue colour, blood red colour- ation confirms presence of N and S both.	$3$ NaSCN + FeCl <sub>3</sub> $\longrightarrow$ [Fe(SCN) <sub>3</sub> ] + 3NaCl Blood red colour



- 1. An aqueous solution of 6.3 g oxalic acid dihydrate is made upto 250 mL. The volume of 0.1 N NaOH required to completely neutralise 10 mL of this solution is
  - (a) 40 mL (b) 20 mL
  - (c) 10 mL (d) 4 mL
- 2. Which of the following compounds does not show Lassaigne's test for nitrogen?
  - (a) Urea (b) Hydrazine
  - (c) Phenylhydrazine (d) Azobenzene
- 3. The group reagent for the group V radicals is
  - (a)  $(NH_4)_2CO_3$ (b)  $(NH_4)_2SO_4$
  - (c) NH<sub>4</sub>Cl (d)  $(NH_4)_2C_2O_4$
- **4.** In qualitative analysis,  $Cd^{2+}$  is under
  - (a) group IV (b) group III
  - (d) group I. (c) group II

- A colourless salt gives violet colour in Bunsen 5. flame, it may be
  - (a)  $Na_2CO_3$ (b) Na<sub>2</sub>CrO<sub>4</sub> (c)  $K_2CO_3$ (d) BaCO<sub>3</sub>.
- 6. A doctor by mistake administered  $Ba(NO_3)_2$ solution to a patient for radiography investigations. Which of the following should be given as the best to prevent the absorption of soluble barium?
  - (a)  $Na_2CO_3$ (b) NH<sub>4</sub>Cl (c) NaCl (d)  $Na_2SO_4$
- 7. Which of the following compounds are partially soluble or insoluble in NH<sub>4</sub>OH solution?
  - (2)  $Ag_2CrO_4$ (1) Fe(OH)<sub>3</sub> (4)  $Ag_2CO_3$
  - (3)  $Al(OH)_3$
  - (5) Ni(OH)<sub>2</sub>
  - (a) 1, 3 (b) 2, 3, 5 (c) 1, 3, 5 (d) 2, 3, 4

### iranchembook.ir/edu

8. Unknown salt 'A' +  $K_2Cr_2O_7$  + conc.  $H_2SO_4$  $\rightarrow$  Reddish brown fumes

Which is the correct statement regarding the above observation?

- (a) It confirms the presence of  $Br^{-}$  ion.
- (b) It neither confirms Cl<sup>-</sup> nor Br<sup>-</sup> ion unless it is passed through NaOH solution.
- (c) It confirms the presence of  $Cl^-$  ion.
- (d) It confirms the presence of both Cl<sup>-</sup> and Br<sup>-</sup> ions.
- **9.** A pink coloured salt turns blue on heating. The presence of which cation is most likely?
  - (a)  $Cu^{2+}$  (b)  $Fe^{2+}$  (c)  $Zn^{2+}$  (d)  $Co^{2+}$ (JEE Main Online 2015)

10. The difference between titrant and titrate is that

- (a) titrant is a solution of an accurately known concentration whereas a titrate is the substance whose concentration is to be determined by titration
- (b) titrant is a solution of unknown concentration whereas titrate is a solution of known concentration
- (c) one is titrated against the other
- (d) one is the reactant and the other is the product.
- **11.** In the titration of HCl against  $NH_4OH$ , the pH at the equivalence point will be
  - (a) less than 7 (b) greater than 7
  - (c) equal to 7 (d) none of these.
- **12.** What is the pH of the resulting solution when equal volumes of 0.1 M NaOH and 0.01 M HCl are mixed?

(a) 2.0 (b) 7.0 (c) 1.04 (d) 12.65 (NEET 2015)

- 13. A solution containing  $Na_2CO_3$  and  $NaHCO_3$ is titrated against HCl. The volume of the acid used when methyl orange is added as indicator is 'x' mL and the volume of the acid used when phenolphthalein is added as indicator is 'y' mL. The volume of HCl used for the neutralization of NaHCO<sub>3</sub> will be
  - (a) (x y) mL (b) (x 2y) mL
  - (c) (x + y) mL (d) (x + 2y) mL
- A solution containing Na<sub>2</sub>CO<sub>3</sub> and NaOH requires 300 mL of 0.1 N HCl using phenolphthalein as an

indicator. Methyl orange is then added to above titrated solution when a further 25 mL of 0.2 N HCl is required. The amount of NaOH present in solution is

- (a) 0.8 g (b) 1.0 g
- (c) 1.5 g (d) 2 g
- **15.** In Lassaigne's test, the organic compound is fused with a piece of sodium metal in order to
  - (a) increase the ionisation of the compound
  - (b) decrease the melting point of the compound
  - (c) increase the reactivity of the compound
  - (d) convert the covalent compound into a mixture of ionic compounds.
- 16. The only cations present in a slightly acidic solution are  $Fe^{3+}$ ,  $Zn^{2+}$  and  $Cu^{2+}$ . The reagent that when added in excess to this solution would identify and separate  $Fe^{3+}$  in one step is
  - (a) 2 M HCl (b) 6 M NH<sub>3</sub>
  - (c) 6 M NaOH (d)  $\text{H}_2\text{S}$  gas.
- 17. An aqueous solution of a salt *X* turns blood red on treatment with KSCN and blue on treatment with  $K_4[Fe(CN)_6]$ . *X* also gives positive chromyl chloride test. The salt *X* is
  - (a) CuCl<sub>2</sub> (b) FeCl<sub>3</sub> (c) Cu(NO<sub>3</sub>)<sub>2</sub> (d) Fe(NO<sub>3</sub>)<sub>3</sub> (JEE Main Online 2015)
- **18.** Before testing for halogens, the Lassaigne's extract is boiled with
  - (a) conc. HCl (b) NaOH
  - (c) conc.  $HNO_3$  (d) any of these.
- 19. A solid compound X on heating gives CO<sub>2</sub> gas and a residue. The residue when mixed with water forms Y. On passing an excess of CO<sub>2</sub> through Y in water, a clear solution Z is obtained. On boiling Z, compound X is reformed. The compound X is
  (a) Ca(HCO<sub>3</sub>)<sub>2</sub>
  (b) CaCO<sub>3</sub>
  - (c)  $Na_2CO_3$  (d)  $K_2CO_3$
- **20.** A salt is heated first with dil.  $H_2SO_4$  and then with conc.  $H_2SO_4$ . No reaction takes place. It may be
  - (a) nitrate (b) sulphide
  - (c) oxalate (d) sulphate.
- **21.** The phenomenon in which white transparent crystal changes into white powder is called
  - (a) efflorescence (b) allotropy
  - (c) sublimation (d) none of these.



- 22. Which of the following salts will give highest pH in water?
  - (a) KCl (b) NaCl
  - (c)  $Na_2CO_3$  (d)  $CuSO_4$  (NEET 2014)
- **23.** Give the correct order of initials T or F for the following statements. Use T if statement is true and F if it is false :
  - (i) Cu<sup>+</sup> undergoes disproportionation to Cu and Cu<sup>2+</sup> in aqueous solution.
  - (ii) HgCl<sub>2</sub> does not give chromyl chloride test.
  - (iii) Sulphide ions react with sodium nitroprusside to form a green coloured complex. In this reaction, oxidation state of iron changes.
  - (a) TFT (b) TTF (c) TFF (d) FTT
- **24.** The presence of magnesium is confirmed in the qualitative analysis by the formation of a white crystalline precipitate of
  - (a)  $Mg(HCO_3)_2$  (b)  $MgNH_4PO_4$
  - (c)  $MgNH_4(HCO_3)_3$  (d)  $MgCO_3$

**25.** Match the following :

	Colu	ımn-]	[			Column-II
P.	Presence of halogen			ogen	1.	HNO <sub>3</sub> /AgNO <sub>3</sub>
Q.	Presence of sulphur			ohur	2.	FeCl <sub>3</sub>
R.	Presence of nitrogen			ogen	3.	$Na_2[Fe(CN)_5 NO]$
S.	Presence of nitrogen			ogen	4.	FeSO <sub>4</sub> /dil. H <sub>2</sub> SO <sub>4</sub>
	and sulphur					
	Р	Q	R	<b>S</b>		
(a)	1	3	4	2		
(b)	2	4	1	2		
(c)	1	2	3	4		
(d)	4	3	2	1		

26.  $Cl_2$  gas is continuously passed with constant shaking through Lassaigne's extract containing  $CS_2$ . If the extract contains both NaBr and NaI, first a violet colour is produced, then the organic layer turns colourless and finally orange colour is seen in the  $CS_2$  layer. The  $CS_2$  turns colourless between violet and orange colour due to the formation of

(a) IBr (b) ICl (c) BrCl<sub>3</sub> (d) NaI<sub>3</sub>

- **27.** Positive Beilstein test for halogens shows that
  - (a) a halogen is absent

62

- (b) a halogen is definitely present
- (c) a halogen may be present
- (d) all the statements are correct.

- The weight of oxalic acid required to neutralise 100 mL of 1 N NaOH solution, is
  - (a) 6.3 g (b) 126 g (c) 530 g (d) 63 g
- **29.** In the titration of HCl against NaOH the indicator(s) used and the end point(s) obtained are
  - (a) phenolphthalein, pink to colourless
  - (b) methyl orange, yellow to red
  - (c) both (a) and (b) separately
  - (d) none of these.
- 30. Reddish-brown (chocolate) ppt. is formed with
  - (a)  $Cu^{2+}$  and  $Fe(CN)_{6}^{4-}$  (b)  $Ba^{2+}$  and  $SO_{4}^{2-}$
  - (c)  $Pb^{2+}$  and  $I^{-}$  (d) none of these.

#### SOLUTIONS

1. (a): Normality of oxalic acid =  $\frac{6.3 \times 1000}{63 \times 250} = 0.4$  $N_1 V_{1(\text{NaOH})} = N_2 V_{2 \text{ (oxalic acid)}}$  $0.1 \times V_1 = 0.4 \times 10 \implies V_1 = 40 \text{ mL}$ 

2. (b): Hydrazine  $(NH_2NH_2)$  does not contain carbon and hence on fusion with Na metal, it cannot form NaCN, consequently hydrazine does not show Lassaigne's test for nitrogen.

- 3. (a) 4. (c) 5. (c)
- 6. (d):  $Ba(NO_3)_2$  gives precipitate of  $BaSO_4$  with  $Na_2SO_4$ .
- 7. (a): Due to insoluble nature of  $Fe(OH)_3$  and  $Al(OH)_3$  in  $NH_4OH$  such metals are detected in group III analysis.
- **8.** (b): When brown vapours of chromyl chloride  $(CrO_2Cl_2)$  are passed through NaOH solution it gives yellow solution of Na<sub>2</sub>CrO<sub>4</sub> which confirms the presence of Cl<sup>-</sup> ions.
- **9.** (d) : In aqueous solution,  $Co^{2+}$  ion exists as  $[Co(H_2O)_6]^{2+}$  ion and has pink colour. When heated, it turns blue due to dehydration.

**10.** (a) : Titrant is a solution of known concentration generally taken in the burette and titrate is the solution of the substance whose concentration is to be determined by titration.

11. (a) : HCl + NH<sub>4</sub>OH 
$$\xrightarrow{\text{Neutralization}}$$
 NH<sub>4</sub>Cl + H<sub>2</sub>O  
NH<sub>4</sub>Cl + H<sub>2</sub>O  $\xrightarrow{\text{Hydrolysis}}$  NH<sub>4</sub>OH + HCl

Since on hydrolysis, strong acid, HCl is produced, the pH of the solution will be less than 7.

12. (d): One mole of NaOH is completely neutralised by one mole of HCl.

Hence, 0.01 mole of NaOH will be completely neutralised by 0.01 mole of HCl.

 $\Rightarrow$  NaOH left unneutralised = 0.1 – 0.01 = 0.09 mol As equal volumes of two solutions are mixed,

$$[OH]^{-} = \frac{0.09}{2} = 0.045 \text{ M}$$

 $\Rightarrow$  pOH = -log(0.045) = 1.35

∴ pH = 14 – 1.35 = 12.65

**13.** (b): Methyl orange (*M*) indicates the neutralization of both Na<sub>2</sub>CO<sub>3</sub> and NaHCO<sub>3</sub> in a mixture with HCl solution and the volume used is 'x' mL.

$$Na_2CO_3 + 2HCl \xrightarrow{M} 2NaCl + H_2O + CO_2$$
  
NaHCO<sub>2</sub> + HCl  $\xrightarrow{M}$  NaCl + H<sub>2</sub>O + CO<sub>2</sub>

Phenolphthalein (P) indicates the neutralization of Na<sub>2</sub>CO<sub>3</sub> only upto single stage, *i.e.*, upto NaHCO<sub>3</sub> stage with HCl and the volume used is '*y*' mL.

 $Na_2CO_3 + HCl \xrightarrow{P} NaHCO_3 + NaCl$ 

 $\therefore$  For complete neutralization of Na<sub>2</sub>CO<sub>3</sub>, the volume of HCl used will be 2y mL. Hence, the volume of HCl used for netralization of NaHCO<sub>3</sub> only = (x - 2y) mL.

14. (b): Let x eq. of  $Na_2CO_3$  and y eq. of NaOH be present in the solution. In presence of phenolphthalein the reactions are

 $Na_2CO_3 + HCl \longrightarrow NaHCO_3 + NaCl$  $NaOH + HCl \longrightarrow NaCl + H_2O$ 

$$\therefore \quad \frac{x}{2} + y = \frac{300 \times 0.1}{1000} = 0.03$$

In presence of methyl orange rest of Na<sub>2</sub>CO<sub>3</sub> reacts

 $\frac{x}{2} = \frac{25 \times 0.2}{1000} = 0.005$ 

 $\therefore x = 0.01$ 

Then weight of  $Na_2CO_3 = 0.01 \times 53 = 0.53$  g and y = 0.03 - 0.005 = 0.025 thus, weight of NaOH  $= 0.025 \times 40 = 1$  g

15. (d): To convert covalent compounds into ionic compounds such as NaCN, Na<sub>2</sub>S, NaX, etc.

16. (b): When  $NH_3$  is added to  $Fe^{3+}$ ,  $Zn^{2+}$  and  $Cu^{2+}$  ions we get brown ppt. of  $Fe(OH)_3$ , colourless  $[Zn(NH_3)_4]^{2+}$ soluble and blue  $[Cu(NH_3)_4]^{2^+}$  soluble complex. Fe<sup>3+</sup> + 3NH<sub>3</sub> + 3H<sub>2</sub>O  $\longrightarrow$  Fe(OH)<sub>3</sub> $\downarrow$  + 3NH<sub>4</sub><sup>+</sup>

**17.** (b):  $Fe^{3+}$  radical gives blood red colour with KSCN and blue colour with K<sub>4</sub>[Fe(CN)<sub>6</sub>]. Cl<sup>-</sup> radical gives chromyl chloride test. Thus, the salt X is FeCl<sub>3</sub>.

**18.** (c) : Lassaigne's extract is boiled with  $HNO_3$  to decompose Na<sub>2</sub>S and NaCN which would otherwise interfere with the tests of halogens. HCl cannot be used because we cannot add Cl<sup>-</sup> ions when we have to test for Cl in the organic compound. NaOH also cannot be used because it would precipitate AgNO<sub>3</sub> as AgOH.

**19.** (b): Solid compound X is  $CaCO_3$  which on heating gives CaO (residue) and CO<sub>2</sub> gas. CaO gives Ca(OH)<sub>2</sub> (Y) on reacting with  $H_2O$ . Ca(OH)<sub>2</sub> reacts with excess  $CO_2$  to give a clear solution of  $Ca(HCO_3)_2(Z)$  which on boiling gives  $CaCO_3(X)$  again.

**20.** (d):  $SO_4^{2-}$  does not react with dil. and conc.  $H_2SO_4$ .

21. (a): Efflorescence is the loss of water of crystallisation of hydrated crystals to change into lower hydrate or anhydrous form.

**22.** (c) :  $Na_2CO_3$  which is a salt of NaOH (strong base) and H<sub>2</sub>CO<sub>3</sub> (weak acid) will produce a basic solution with pH greater than 7.

23. (b)

**24.** (b): 
$$MgCl_2 + Na_2HPO_4 + NH_4OH \rightarrow Mg(NH_4)PO_4 + 2NaCl + H_2O$$
  
White ppt.

25. (a)

**26.** (b):  $I_2$  first liberated combines with  $Cl_2$  to form colourless iodine monochloride.

$$I_2 + Cl_2 \longrightarrow 2ICl$$

27. (c): Positive Beilstein test does not necessarily mean that halogen is definitely present because certain compounds such as urea, thiourea, pyridine, etc. which do not contain halogen, also give this test. Therefore, positive Beilstein test shows that halogen may be present.

28. (a): Number of eq. of NaOH = Number of eq. of oxalic acid

$$\therefore \quad \frac{100 \times 1}{1000} = \frac{\text{Wt.of oxalic acid}}{63}$$
  
Weight of oxalic acid = 6.3 g

29. (c): In the titration of strong acid vs strong base, both the indicators phenolphthalein (pink to colourless) and methyl orange (yellow to red) can be used separately.

**30.** (a):  $Cu_2[Fe(CN)_6]$  is chocolate brown colour precipitate.

۲ 🔇



Class XII

Maximise your chance of success, and high rank in NEET, JEE (Main and Advanced) by reading this column. This specially designed column is updated year after year by a panel of highly qualified teaching experts well-tuned to the requirements of these Entrance Tests.



# **Principles Related to Practical Chemistry**

### SURFACE CHEMISTRY

### **Preparation of Lyophilic Sol**

• **Starch sol :** Starch is an intrinsic colloid, forms a hydrophilic sol directly when mixed with hot water. It is quite stable as it is electrically neutral.

**NEET** JEE

- **Gum-arabic :** Gum-arabic is hydrophilic colloid and therefore, readily forms sol on shaking with water. The sol is stable as it is not affected by ionic impurities.
- **Egg albumin :** Egg albumin forms hydrophilic sol with water at room temperature. It is quite stable and not affected by the presence of any impurity.

### **Preparation of Lyophobic Sol**

• Hydrated ferric oxide sol : It is prepared indirectly by the hydrolysis of ferric chloride with boiling water. Ferric ions produced from FeCl<sub>3</sub> solution are adsorbed on the surface of the particles of Fe(OH)<sub>3</sub>.

$$\operatorname{FeCl}_{3(aq)} + \operatorname{3H}_2O_{(l)} \xrightarrow{\Delta} \operatorname{Fe}(OH)_{3(s)} + \operatorname{3HCl}_{(aq)}$$
  
Red sol

• **Hydrated aluminium oxide sol :** It is prepared by hydrolysis of aluminium chloride.

$$AlCl_{3(aq)} + 3H_2O_{(l)} \xrightarrow{\Delta} Al(OH)_{3(s)} + 3HCl_{(aq)}$$
  
White sol

• Arsenious sulphide sol : It is prepared by passing hydrogen sulphide gas through a dilute aqueous solution of arsenious oxide.

$$As_2O_{3(aq)} + 3H_2S_{(g)} \longrightarrow As_2S_{3(s)} + 3H_2O_{(l)}$$
  
Yellow sol

### **CHEMICAL KINETICS**

### Rate of Decomposition of $H_2O_2$ :

 $2H_2O_2 \longrightarrow 2H_2O + O_2$ 

The reaction is catalysed by iodide ions.

$$H_2O_2 + I^- \xrightarrow{Slow} H_2O + IO^-$$
  
Hypoiodite ion

$$IO^- + IO^- \xrightarrow{I \text{ ust}} 2I^- + O_2$$

or  $H_2O_2 + IO^- \xrightarrow{Fast} H_2O + I^- + O_2$ Thus, rate law equation is

Rate = 
$$-\frac{1}{2} \frac{d[H_2O_2]}{dt} = k' [H_2O_2] [I^-]$$

If concentration of  $I^-$  ions remains constant, then the rate law equation is

Rate = 
$$-\frac{1}{2} \frac{d[H_2O_2]}{dt} = k [H_2O_2]$$
 (:: k' [I<sup>-</sup>] = k)

*i.e.*, it is a first order reaction.

# Reaction of lodide lon with $H_2O_2$ at Room Temperature :

Here, reactions involved are

• Main reaction :

$$H_2O_2 + 2I^- + 2H^+ \xrightarrow{Slow} I_2 + 2H_2O$$





# Mad about rehearsing?



Tune. Fine tune. Reach the peak of your readiness for JEE with MTG's 40+16 Years Chapterwise Solutions. It is undoubtedly the most comprehensive 'real' question bank, complete with detailed solutions by experts.

Studies have shown that successful JEE aspirants begin by familiarising themselves with the problems that have appeared in past JEEs as early as 2 years in advance. Making it one of the key ingredients for their success. How about you then? Get 40+16 Years Chapterwise Solutions to start your rehearsals early. Visit www.mtg.in to order online.



Available at all leading book shops throughout the country. For more information or for help in placing your order: Call 0124-6601200 or email:info@mtg.in Visit www.mtg.in for latest offers and to buy online!



**Monitor reaction :** 

$$I_2 + 2S_2O_3^{2-} \xrightarrow{\text{Fast}} S_4O_6^{2-} + 2I^{-}$$

**Indicator reaction :** 

- $I_2$  + Starch  $\longrightarrow$  Blue colour complex
- As the concentration of thiosulphate ion is kept constant, the different time taken  $(t_c)$  for the appearance of blue colour with change in concentration of either reactant indicates the relative rate of reaction.

Initial rate ∝-

The rate of reaction decreases with decrease in the concentration of KI.

The graph of  $1/t_c$  versus volume of KI solution is a straight line.



Rate of reaction ∝ Concentration of KI

Similarly, by keeping I<sup>-</sup> ion concentration constant and taking different concentrations of H<sub>2</sub>O<sub>2</sub>, the rate w.r.t. H<sub>2</sub>O<sub>2</sub> can be found out.

### THERMOCHEMISTRY

### **Enthalpy of Dissolution of Copper Sulphate**

It is the amount of heat absorbed or evolved on dissolving one mole of substance in excess of solvent at a given temperature and pressure. It is +ve when heat is absorbed and -ve when heat is given out during dissolution.

For copper sulphate :

 $CuSO_4.7H_2O_{(s)} + H_2O_{(l)} \longrightarrow CuSO_{4(aq)} + \Delta H; \Delta H = ?$ Heat gained or lost  $(Q) = Mass(m) \times specific heat of the$ solution (*s*) × change in temperature ( $\Delta t$ )

Heat of the dissolution

 $= \frac{Q \times \text{Molecular weight of compound}}{Q \times Molecular \text{ weight of compound}}$ Weight of compound dissolved

66 CHEMISTRY TODAY | MARCH '18

### Enthalpy of Neutralisation for a Strong Acid and a Strong Base

Heat of neutralisation is the heat evolved, when one mole of H<sup>+</sup> ions is completely neutralised by a base. The neutralisation reaction in case of HCl and NaOH is in accordance with Arrhenius theory of ionisation, can be written as

$$\begin{split} \text{NaOH}_{(aq)} + \text{HCl}_{(aq)} &\longrightarrow \text{NaCl}_{(aq)} + \text{H}_2\text{O}_{(l)} \\ \text{H}^+_{(aq)} + \text{OH}^-_{(aq)} &\longrightarrow \text{H}_2\text{O}_{(l)} + 57.3 \text{ kJ} \end{split}$$

Thus, the reaction between HCl and NaOH solution is, in fact, the reaction between  $H^+_{(aq)}$  and  $OH^-_{(aq)}$ , therefore, the heat of neutralisation will be same for neutralisation of all strong acids with strong bases.

### ELECTROCHEMISTRY

### Variation of Cell Potential in Zn|Zn<sup>2+</sup>|| Cu<sup>2+</sup>|Cu with Change in Concentration of Electrolytes (CuSO<sub>4</sub> and ZnSO<sub>4</sub>) at Room Temperature

Reduction potential of an electrode increases with increase in concentration of the electrolyte.

$$M^{n+}_{(aq)} + ne^- \longrightarrow M_{(s)}$$

In the zinc-copper electrochemical cell, zinc electrode acts as anode while copper electrode acts as cathode.

$$E_{\text{cell}}^{\circ} = E_{\text{cathode}}^{\circ} - E_{\text{anode}}^{\circ}$$

 $E_{\text{cell}}^{\circ}$  increases if  $E_{\text{cathode}}^{\circ}$  increases and  $E_{\text{anode}}^{\circ}$  decreases. Thus, higher conc. of Cu<sup>2+</sup> and lower conc. of Zn<sup>2+</sup> ions increase the  $E_{cell}^{\circ}$  for Zn | Zn<sup>2+</sup> || Cu<sup>2+</sup> | Cu.

The relation between conc. of the eletrolyte and the standard electrode potential is given in the form of Nernst equation :

$$E_{\text{cell}} = E_{\text{cell}}^{\text{o}} - \frac{0.059}{n} \log \frac{[\text{Zn}^{2+}]}{[\text{Cu}^{2+}]}$$

Every therapeutic drug has a narrow range of optimal water content, which needs to be controlled to avoid potential adverse effect on patients. Recently, chemists have invented a method to quantify water content in solid pharmaceutical drugs that is faster, cheaper, more accurate and more precise than Karl Fischer titration method. In this method, water determination is done using ionic liquids and headspace gas chromatography.



### CHROMATOGRAPHY

Chromatography is a modern and sensitive technique used for rapid and efficient analysis and separation of components of a mixture and purification of compounds.

**Principle :** This technique is based on the differential migration of individual components of a mixture through a stationary phase under the influence of moving phase.

The components of the mixture rise up at different rates and thus, get separated from one another and then the  $R_f$  value is calculated using the given formula :





### **PREPARATION OF ORGANIC COMPOUNDS**



CHEMISTRY TODAY | MARCH '18

### **DETECTION OF FUNCTIONAL GROUPS**

### **Test for Unsaturation**

Bromine test :  $C = C + Br_2 - Alkene (Orange)$ CBr — CBr Dibromoalkane red) (Colourless)  $-C \equiv C - + 2Br_2$  $-CBr_2 - CBr_2 - CBr_2$ (Orange Tetrabromoalkane Alkyne red) (Colourless) Baeyer's test or Permanganate test :

 $3CH_2 \equiv CH_2 + 2KMnO_4 + 4H_2O \longrightarrow$ Ethene (Pink)  $2KOH + 2MnO_2 + 3CH_2OH.CH_2OH$ (Brown) Glycol (Colourless)  $CH \equiv CH + 2KMnO_4 + 2H_2O \longrightarrow$ Ethyne (Pink)  $2KOH + 2MnO_2 + (COOH)_2$ 

(Brown) Oxalic acid (Colourless)

### Test for Alcoholic (-OH) Group

•

•

.

- Sodium metal test :  $\rightarrow RONa + 1/2H_{2(g)}$  $ROH_{(l)} + Na_{(s)}$ -Ethanol Sodium Sodium Hydrogen (sample) ethoxide (effervescence)
- Ceric ammonium nitrate test :  $2ROH + (NH_4)_2[Ce(NO_3)_6] -$ Colourless Ceric ammonium (sample) nitrate (yellow)  $(ROH)_2Ce(NO_3)_4 + 2NH_4NO_3$
- (Pink or purple) (Colourless) Ester test : Conc. H<sub>2</sub>SO<sub>4</sub> ROH + CH<sub>3</sub>COOH Ethanol Acetic acid

 $CH_3COOR + H_2O$ (sample) Ethyl acetate Water (fruity smell) Acetyl chloride test :

 $R - OH + CH_3COCl \longrightarrow CH_3COOR + HCl^{\uparrow}$ Alcohol Acetvl Ester (sample) chloride  $HCl\uparrow + NH_4OH \rightarrow \mathrm{NH_4Cl}^\uparrow + \mathrm{H_2O}$ White dense fumes

Xanthate test : (i)  $\Delta$ ROH + KOH  $\rightarrow ROK + H_2O$ (ii) Cool Alcohol (iii) Ether (Sample) ROK +

$$CS_2 \longrightarrow RO - C \swarrow S^-K^+$$

Yellow ppt.

Distinction between 1°, 2° and 3° alcohols : Organic compound + Lucas reagent (Anhy. ZnCl<sub>2</sub> + conc. HCl) and shake :

- Separation of an insoluble layer at once confirms 3° alcohol.
- Appearance of cloudiness within 4-5 minutes confirms 2° alcohol.
- Clear solution confirms 1° alcohol.

### **Test for Phenolic Group**

- Ferric chloride test :  $6C_6H_5OH + FeCl_3 \longrightarrow [Fe(C_6H_5O)_6]^{3-} + 3HCl$ Phenol (Neutral) (Green/Red/Violet colour) (sample)
- **Litmus test :** Sample solution + Blue litmus → Red colour
- Br<sub>2</sub>-water test : • Gives white ppt. of 2,4,6-Tribromophenol.
- Liebermann's test :



Phthalein test :



## iranchembook.ir/edu



CHEMISTRY TODAY | MARCH '18



### Distinction between 1°, 2° and 3° Amine (Hinsberg's Test)

### Test for Carboxylic (—COOH) Group

- Sodium bicarbonate test :  $RCOOH + NaHCO_3 \rightarrow RCOONa + H_2O + CO_2^{\uparrow}$ (Sample) (Effervescence)
- Litmus test :  $RCOOH + Blue litmus \rightarrow Red$ (Sample)

Note : Phenols also give this test.

Ester test : •  $RCOOH + C_2H_5OH \xrightarrow{\text{conc. } H_2SO_4} RCOOC_2H_5 + H_2O$ Ester (Fruity smell)

### **Characteristic Tests for Carbohydrates, Fats** and Proteins

Tests for carbohydrates :



- **Tests for lipids :** 
  - Grease spot test : Lipid leaves translucent spot  $\geq$ on unglazed brown paper bags.
  - Sudan red test : Sudan red is a fat soluble dye that stains lipids red.

## Test for proteins :

**Biuret test :** 

For peptides with Biuret solution Purple or chain length of at (Blue) + Sample pink solution least 3-amino acids

### Volumetric Analysis

- In a titration, a volume of a standardized solution containing a known concentration of reactant 'A' is added incrementally to a sample containing an unknown concentration of reactant 'B' till reactant 'B' is just consumed (stoichiometric completion). This is known as the equivalence point. At this point we have,  $N_1V_1 = N_2V_2$ .
- Strength = Normality × Equivalent weight Relation between normality and molarity : Normality =  $n \times$  Molarity where, n = number of equivalents in 1 mole.

### Titration of Oxalic Acid with KMnO<sub>4</sub>

 $2KMnO_4 + 3H_2SO_4 \rightarrow K_2SO_4 + 2MnSO_4 + 3H_2O + 5[O]$ 

$$\begin{array}{c} \text{COOH} \\ 5 \mid & + 5[\text{O}] \rightarrow 5\text{H}_2\text{O} + 10\text{CO}_2^{\uparrow} \\ \text{COOH} \end{array}$$

- Indicator : KMnO<sub>4</sub> is self indicator.
- End point : Appearance of light pink colour.
- **Observations :**

S.No.	Initial burette reading (R <sub>1</sub> )	Final burette reading (R <sub>2</sub> )	Volume of KMnO <sub>4</sub> used $V_1 = (R_2 - R_1)$
1			
2			

**Calculation :**  $n_1 \times M_1 \times V_1 = n_2 \times M_2 \times V_2$ (KMnO<sub>4</sub>) (Oxalic acid)





Molarity of KMnO<sub>4</sub> ( $M_1$ ) =  $\frac{n_2 \times M_2 \times V_2}{n_1 \times V_1}$ where,  $n_1 = 5$  and  $n_2 = 2$ 

### Titration of Mohr's Salt vs KMnO<sub>4</sub>

- $2KMnO_4 + 8H_2SO_4 + 10FeSO_4 \cdot (NH_4)_2SO_4 \cdot 6H_2O$   $\rightarrow K_2SO_4 + 2MnSO_4 + 5Fe_2(SO_4)_3 + 10(NH_4)_2SO_4$  $+ 68H_2O$
- **Indicator** : KMnO<sub>4</sub> is self indicator.
- End Point : Appearance of light pink colour.

### **P**REPARATION OF INORGANIC COMPOUNDS

• Observation :

S.No.	Initial	Final	Volume of
	burette	burette	KMnO <sub>4</sub> used
	reading $(R_1)$	reading (R <sub>2</sub> )	$V_1 = (R_2 - R_1)$
1			
2			

**Calculation :** 
$$n_1 \times M_1 \times V_1 = n_2 \times M_2 \times V_2$$
  
(KMnO<sub>4</sub>) (Mohr's salt)  
Molarity of KMnO<sub>4</sub>( $M_1$ ) =  $\frac{n_2 \times M_2 \times V_2}{n_1 \times V_1}$   
where,  $n_1 = 5$  and  $n_2 = 1$ 

Ferrous Ammonium Sulphate (Mohr's S<u>alt)</u>\_\_\_\_\_

It is a double salt, prepared by dissolving equimolar quantities of iron(II) sulphate and ammonium sulphate in water containing a little amount of sulphuric acid, and then subjecting the resulting solution to crystallization.

 $\begin{array}{ll} \text{FeSO}_4.7\text{H}_2\text{O} + (\text{NH}_4)_2\text{SO}_4 \longrightarrow \text{FeSO}_4.(\text{NH}_4)_2\text{SO}_4.6\text{H}_2\text{O} + \text{H}_2\text{O} \\ (1 \text{ mole}, 278 \text{ g}) & (1 \text{ mole}, 132 \text{ g}) & (1 \text{ mole}, 392 \text{ g}) \end{array}$ 

### Potash alum (*Phitkari*)

It is a double salt of potassium sulphate and aluminium sulphate prepared by concentrating a solution containing equimolar quantities of  $K_2SO_4$  and  $Al_2(SO_4)_3.18H_2O$  to crystallization point followed by cooling.

 $\begin{array}{rl} K_2SO_4 &+& Al_2(SO_4)_3.18H_2O + 6H_2O \longrightarrow K_2SO_4.Al_2(SO_4)_3.24H_2O \\ (1 \text{ mole}, 174 \text{ g}) && (1 \text{ mole}, 666 \text{ g}) && (1 \text{ mole}, 948 \text{ g}) \end{array}$ 



- 1. Liebermann's test is used for identification of which functional group in organic compounds?
  - (a) Alcohol (b) 1° Amine
  - (c) Phenolic (d) Aldehyde
- 2. Which of the following statements is not correct regarding preparation of *p*-nitroacetanilide?
  - (a) Nitration of acetanilide gives a mixture of *o* and *p*-nitroacetanilide in which *p*-isomer predominates.
  - (b) Nitration is carried out with a mixture of conc.  $HNO_3$  + conc.  $H_2SO_4$
- (c) During crystallisation of the crude product,
   *p*-nitroacetanilide crystallises out leaving
   *o*-nitroacetanilide in the mother liquor.
- (d) *o*-Nitroacetanilide is colourless but *p*-nitroacetanilide is yellow coloured
- **3.** The purpose of adding dilute sulphuric acid in the preparation of Mohr's salt is to
  - (a) prevent the hydrolysis of ferrous sulphate
  - (b) increase the solubility of the salts used
  - (c) prevent the precipitation of carbonates of metals
  - (d) neutralize ammonium salts.



- **4.** Some pale green crystals are strongly heated. The gases given off are passed into a container surrounded by ice and then through a solution of acidified KMnO<sub>4</sub>. KMnO<sub>4</sub> is decolourised and a waxy white solid is formed in the ice container, this is dissolved in water. The solution will
  - (a) give a precipitate with silver nitrate solution
  - (b) give a precipitate with barium chloride solution
  - (c) turn red litmus blue
  - (d) give blue colour with starch solution.
- Experiments on the oxidation of I<sup>-</sup> ions by H<sub>2</sub>O<sub>2</sub> in presence of dilute H<sub>2</sub>SO<sub>4</sub>, S<sub>2</sub>O<sub>3</sub><sup>2-</sup> ions and starch were repeated, taking the same amount of 3% H<sub>2</sub>O<sub>2</sub> but varying amounts of 0.5 M KI solution (*e.g.*, 20 mL, 16 mL, 12 mL etc. and making the volume to 20 mL in each case by adding water). The time for the appearance of blue colour was noted. Then
  - (a) a plot of volume of KI solution *vs* time will be linear
  - (b) a plot of volume of KI solution *vs* 1/time will be linear
  - (c) a plot of 1/volume of KI solution *vs* 1/time will be linear
  - (d) there is no relationship between volume of KI solution and time.
- 6. A solution containing a group IV cation gives a precipitate on passing  $H_2S$  gas. A solution of this precipitate in dil. HCl produces a white precipitate with NaOH solution and bluish white precipitate with basic potassium ferrocyanide. The cation is (a)  $Co^{2+}$  (b)  $Ni^{2+}$  (c)  $Zn^{2+}$  (d)  $Mn^{2+}$ .

### (JEE Main Online 2017)

- 7. 3.92 g of ferrous ammonium sulphate reacts completely with 50 mL of  $\frac{N}{10}$  KMnO<sub>4</sub> solution. The percentage purity of the sample is (a) 50 (b) 78.4 (c) 80 (d) 39.2.
- 8. A metal chloride dissolves appreciably in cold water and when placed on a platinum wire in Bunsen flame, no distinctive colour is noticed, the cation would be

(a) 
$$Mg^{2+}$$
 (b)  $Ba^{2+}$  (c)  $Pb^{2+}$  (d)  $Ca^{2+}$ .

**9.** A white crystalline solid *A*, on boiling with caustic soda solution gave a gas *B* which when passed through an alkaline solution of potassium mercuric iodide gave a brown ppt. The substance *A* on heating gave a gas *C* which rekindled a glowing splinter but



CHEMISTRY TODAY | MARCH '18

did not give brown fumes with nitric oxide. The gases *B*, *C* and the substance *A* respectively are

- (a)  $H_2S$ ,  $NO_2$  and NaCl
- (b)  $NH_3$ ,  $N_2O$  and  $NH_4NO_3$
- (c) HCl, NO and  $NH_4Cl$
- (d)  $CO_2$ ,  $SO_2$  and  $Na_2SO_3$ .
- **10.** Match the list-I with list-II and choose the correct option using the codes given below :

List - I				List - II		
(Functional group)				(Reagent)		
P.	-OF	ł		(i)	Ceric ammonium	
					nitrate	
Q.	-CC	OOH		(ii)	2, 4-DNP	
R.	}c=	0		(iii)	NaHCO <sub>3</sub>	
S.	-NF	$H_2$		(iv)	C <sub>6</sub> H <sub>5</sub> SO <sub>2</sub> Cl	
	Р	Q	R		S	
(a)	(i)	(ii)	(iii)	(i	iv)	
(b)	(iv)	(iii)	(ii)	(	(i)	
(c)	(i)	(ii)	(iv)	(	ii)	
(d)	(i)	(iii)	(ii)	(i	iv)	

**11.** Dimethylglyoxime (DMG) gives a red precipitate with Ni<sup>2+</sup>, which is used for its detection. To get this precipitate, the best pH range is

(a) < 1 (b) 2 - 3 (c) 3 - 4 (d) 9 - 11

12. The pair(s) of ions where both the ions are precipitated upon passing H<sub>2</sub>S gas in presence of dilute HCl, is(are)

(a) 
$$Ba^{2+}, Zn^{2+}$$
 (b)  $Bi^{3+}, Fe^{3+}$   
(c)  $Cu^{2+}, Pb^{2+}$  (d)  $Hg^{2+}, Bi^{3+}$ 

(JEE Advanced 2015)

- 13. The brown ring test for NO<sub>2</sub><sup>-</sup> and NO<sub>3</sub><sup>-</sup> ions is due to the formation of complex ion with formula,
  (a) [Fe(H<sub>2</sub>O)<sub>6</sub>]<sup>2+</sup>
  (b) [Fe(NO)(CN)<sub>5</sub>]<sup>2-</sup>
  - (c)  $[Fe(H_2O_5(NO)]^{2+}$  (d)  $[Fe(H_2O)(NO)_5]^{2+}$
- 14. (Clear solution)  $B \xleftarrow{\text{dil. HCl}} A \xrightarrow{\text{K}_2\text{CrO}_4} D$  (Yellow ppt.)  $\downarrow^{\text{dil.H}_2\text{SO}_4} C$  (White ppt.)

Compound *A* is

- (a) barium carbonate (b) calcium carbonate
- (c) lead carbonate (d) red lead.
- **15.** KMnO<sub>4</sub> reacts with oxalic acid according to the equation,
  - $2MnO_4^- + 5C_2O_4^{2-} + 16H^+ \rightarrow 2Mn^{2+} + 10CO_2 + 8H_2O$ Here 20 mL of 0.1 M KMnO<sub>4</sub> is equivalent to

- (a) 50 mL of 0.5 M  $C_2H_2O_4$
- (b) 20 mL of 0.1 M  $C_2H_2O_4$
- (c) 20 mL of 0.5 M  $C_2H_2O_4$
- (d) 50 mL of 0.1 M  $C_2H_2O_4$ .
- 16. An organic compound produces effervescence with NaHCO<sub>3</sub> solution. It dissolves in NaOH solution and gets precipitated when dil. HCl is added to it. The compound is
  - (a) acetic acid (b) benzoic acid
  - (c) oxalic acid (d) formic acid.
- **17.** Leveling bulb is used during experiment to study kinetics of dissociation of hydrogen peroxide to ensure
  - (a) pressure within the reaction vessel is same as that in the room
  - (b) uniform pressure difference between the room and the gases in the system
  - (c) same temperature as that of room
  - (d) none of these.
- **18.** Gold sol has been prepared by shaking gold chloride (AuCl<sub>3</sub>) solution with formaldehyde. Which one of the following is true?
  - (a)  $AuCl_3$  has been oxidised by formaldehyde.
  - (b) AuCl<sub>3</sub> has been reduced by formaldehyde.
  - (c) AuCl<sub>3</sub> has been reduced as well as oxidised, *i.e.*, it is a disproportionation reaction.
  - (d) Reaction between AuCl<sub>3</sub> and HCHO is a displacement reaction.
- 19. Which of the following statements is incorrect?
  - (I) In  $S_2O_3^{2-}$ , both sulphur are different in nature.
  - (II) Sodium acetate and lead acetate on heating give same type of products, whereas Mn, Sn, Fe oxalate salts give different types of products.
  - (III) Aqueous solution of OCl<sup>-</sup>,  $S^{2-}$  and  $CO_3^{2-}$  ions is basic in nature.
  - (IV)  $NO_2^-$  oxidises  $I^-$  ion whereas  $Br_2$  and  $Cl_2$  oxidises  $NO_2^-$  ions.
  - (a) II, IV only (b) I, II, IV only
  - (c) II only (d) II, III, IV only
- **20.** Which of the following compounds is responsible for yellow colour in the xanthate test for alcoholic group?

(a) 
$$RO - \overset{\text{o}}{C} - O^{-}K^{+}$$
 (b)  $RO - \overset{\text{o}}{C} - S^{-}K^{+}$   
(c)  $RO - \overset{\text{o}}{C} - SR$  (d)  $RO - \overset{\text{o}}{C} - OR$ 

- **21.** The cation that will not be precipitated by H<sub>2</sub>S in the presence of dil. HCl is
  - (a)  $Cu^{2+}$  (b)  $Pb^{2+}$  (c)  $As^{3+}$  (d)  $Co^{2+}$ (JEE Main Online 2015)
- **22.** To determine the enthalpy of neutralization of HCl with NaOH, we use
  - (a) 0.2 N HCl and 0.2 N NaOH
  - (b) 0.5 N HCl and 0.5 N NaOH
  - (c) 1 N HCl and 1 N NaOH
  - (d) any of these.
- 23. Sometimes a brown precipitate of hydrated MnO<sub>2</sub> (MnO<sub>2</sub>.H<sub>2</sub>O) is observed in KMnO<sub>4</sub> titrations. This is due to
  - (a) insufficient quantity of dil. H<sub>2</sub>SO<sub>4</sub> which results in incomplete reduction of KMnO<sub>4</sub>
  - (b) over heating of the solution
  - (c) dropwise addition of KMnO<sub>4</sub> with constant shaking
  - (d) addition of hydrochloric acid.
- 24. In the preparation of *p*-nitroacetanilide from aniline, nitration is not done by using nitrating mixture (a mixture of conc.  $H_2SO_4$  and conc.  $HNO_3$ ) because




- (a) it gives *o*-nitroacetanilide
- (b) it gives a mixture of *o* and *p*-nitroaniline
- (c)  $-NH_2$  group gets oxidised
- (d) it forms a mixture of *o*-and *p*-nitroacetanilide.
- **25.** Few drops of HNO<sub>3</sub> are added before proceeding for group III in order to
  - (a) convert  $Fe^{2+}$  to  $Fe^{3+}$
  - (b) convert  $Fe^{3+}$  to  $Fe^{2+}$
  - (c) precipitate group III radicals
  - (d) none of these.
- 26. A known coloured solution containing two metal ions, was treated with excess cold sodium hydroxide solution. When filtered a whitish solid, slowly changing to brown, was retained on the filter paper and a colourless solution collected as the filtrate. Dropwise addition of hydrochloric acid to the filtrate produced a white ppt. which dissolved in excess acid. Treatment of the residue on filter paper with a solution of strong oxidiser produced a reddish-violet solution. Indicate the pairs of ions.
  - (a)  $Mn^{2+}$  and  $Mg^{2+}$  ions
  - (b)  $Fe^{2+}$  and  $Zn^{2+}$  ions
  - (c)  $Zn^{2+}$  and  $Mn^{2+}$  ions
  - (d)  $Mg^{2+}$  and  $Zn^{2+}$  ions
- 27. A solution of metal hydroxide (MOH) with copper sulphate and mixed tartarate of metal M with another metal  $M_1$  of the same group, is used in the detection of —CHO group. Metal M and  $M_1$  are respectively

**28.** In the electrochemical cell :

Zn|ZnSO<sub>4</sub>(0.01 M)||CuSO<sub>4</sub>(1.0 M)|Cu, the emf of this Daniell cell is  $E_1$ . When the concentration of ZnSO<sub>4</sub> is changed to 1.0 M and that of CuSO<sub>4</sub> changed to 0.01 M, the emf changes to  $E_2$ . Which one is the correct relationship between  $E_1$  and  $E_2$ ? (Given, RT/F = 0.059)

(a) 
$$E_1 < E_2$$
 (b)  $E_1 > E_2$ 

(c) 
$$E_2 = 0 E_1$$
 (d)  $E_1 = E_2$  (NEET 2017)

29. Ferric hydroxide sol can be prepared

- (a) by shaking FeCl<sub>3</sub> solution with dilute NaOH solution
- (b) by shaking freshly precipitated  $\mbox{FeCl}_3$  with cold water
- (c) by shaking freshly precipitated Fe(OH)<sub>3</sub> with water
- (d) by shaking FeCl<sub>3</sub> with boiling water.



- **30.** The sol of egg albumin is prepared by shaking
  - (a) the white of the egg with cold water
  - (b) the yellow of the egg with cold water
  - (c) the white of the egg with hot water
  - (d) yellow of the egg with hot water.

### SOLUTIONS

#### 1. (c)

**2.** (d): *o*-Nitroacetanilide is yellow coloured while *p*-nitroacetanilide is colourless.

3. (a) :  $FeSO_4$  undergoes hydrolysis to produce  $Fe(OH)_2$ and  $H_2SO_4$ .

 $FeSO_4 + 2H_2O \rightarrow Fe(OH)_2 + H_2SO_4$ 

To prevent hydrolysis, dil.  $H_2SO_4$  is added in the preparation of Mohr's salt.

4. (b): The salt can be  $FeSO_4.7H_2O$  which produces  $SO_2$ .  $SO_2$  decolourises  $KMnO_4$  and gets oxidised to  $SO_3$  which dissolves in water to give  $H_2SO_4$  which gives white ppt. of  $BaSO_4$  with  $BaCl_2$  solution.

5. (b): Volume of KI solution represents concentration and 1/time represents rate of reaction.

As rate of reaction  $\infty$  concentration of KI solution, hence a plot of volume of KI solution versus 1/time will be linear.

$$M \times V = 392 \times 1$$
  
Let % purity be x.

then,  $N_1V_1 = N_2V_2$  $\frac{x}{100}(0.01 \times 1000) = \left(\frac{1}{10} \times 50\right)$   $\left(\frac{x}{100} \text{ is the factor of purity}\right)$   $x = \frac{100 \times 50 \times 0.1}{0.01 \times 1000} = 50\%$ 

8. (a):  $MgCl_2$  is soluble in cold water and does not give flame test.

9. (b): Substance A is  $NH_4NO_3$  which on boiling with NaOH gives  $NH_3$  (gas B)

$$NH_4NO_3 + NaOH \rightarrow NaNO_3 + NH_3 + H_2O$$
(A)
(B)

 $NH_3$  gives brown ppt. with Nessler's reagent ( $K_2HgI_4$ )  $2K_2[HgI_4] + NH_3 + 3KOH \rightarrow$ 

$$H_2NHgO.HgI + 7KI + 2H_2C$$
  
Brown ppt.

10. (d)

11. (d): At pH 9-11, dimethylglyoxime forms a red precipitate with Ni<sup>2+</sup> as Ni-DMG complex.

12. (c, d) : Group II radicals will get precipitated out upon passing H<sub>2</sub>S gas in presence of dilute HCl. CuS, PbS, HgS and  $Bi_2S_3$  give black precipitates with dil. HCl. 13. (c)

14. (a): A is  $BaCO_3$  which gives clear solution by reacting with dil. HCl but gives white ppt. of BaSO4 with dil. H<sub>2</sub>SO<sub>4</sub> while yellow ppt. of BaCrO<sub>4</sub> with K<sub>2</sub>CrO<sub>4</sub> solution.

**15.** (d): 
$$2MnO_4^- + 5C_2O_4^{2-} + 16H^+ \rightarrow 2Mn^{2+} + 10CO_2 + 8H_2O$$

 $\therefore$  2 moles of MnO<sub>4</sub><sup>-</sup> = 5 moles of C<sub>2</sub>O<sub>4</sub><sup>2-</sup> 20 mL of 0.1 M KMnO<sub>4</sub> = 2 mmol of KMnO<sub>4</sub>

Also, 50 mL of 0.1 M  $C_2H_2O_4 = 5 \text{ mmol of } C_2O_4^2$ Therefore, these are equivalent.

16. (b): Since the compound dissolves in NaOH and gets precipitated on addition of HCl, therefore, it must be benzoic acid.

17. (a)

**18.** (b):  $AuCl_3$  is reduced by formaldehyde to form colloidal gold sol.

19. (c) 20. (b)

**21.** (d):  $Co^{2+}$  ion present in group IV is precipitated by H<sub>2</sub>S in presence of NH<sub>4</sub>OH. Other ions are precipitated as sulphides in presence of dil. HCl in group II.

22. (d): As in every case, normalities of HCl and NaOH are equal, any of the given pair of acid and base can be used.

23. (a): A brown ppt. of  $MnO_2$  will be formed if insufficient quantity of dil. H<sub>2</sub>SO<sub>4</sub> is added or H<sub>2</sub>SO<sub>4</sub> is not added at all in the titration process. This is due to incomplete reduction of KMnO<sub>4</sub>.

24. (c) : Nitration of aniline is difficult to carry out with nitrating mixture since -NH<sub>2</sub> group gets oxidised which is not required. So, the amino group is first protected by acylation to form acetanilide which is then nitrated to give *p*-nitroacetanilide as a major product.

25. (a) **26.** (c) :  $Zn^{2+} + Mn^{2+}$ NaOH salt  $ZnO_2^{2-}$  $MnO_4^{2-}$ Manganate solution White ppt. soluble (slowly changes in excess NaOH into brown ppt.) (filtrate) oxidation HCl  $MnO_4^ ZnCl_2$ (Reddish violet solution) White ppt

(Dissolve in excess of HCl)

**27.** (c) : NaOH + CuSO<sub>4</sub> + sodium (*M*), potassium ( $M_1$ ) tartarate solution is known as Fehling solution and is used for the detection of -CHO group.

28. (b): 
$$E_{\text{cell}} = E_{\text{cell}}^{\circ} - \frac{0.059}{n} \log \frac{[\text{Zn}^{2+}]}{[\text{Cu}^{2+}]}$$
  
 $E_1 = E^{\circ} - \frac{0.059}{2} \log \frac{0.01}{1}$   
 $E_1 = E^{\circ} - \frac{0.059}{2} (-2) = E^{\circ} + 0.059$   
 $E_2 = E^{\circ} - \frac{0.059}{2} \log \frac{1}{0.01} = E^{\circ} - 0.059$ 

Hence,  $E_1 > E_2$ .

**29.** (d):  $Fe(OH)_3$  sol is prepared by shaking  $FeCl_3$  with boiling water.

**30.** (a) : The sol of egg albumin is prepared by shaking white of the egg with cold water.

Your favourite MTG Books/Magazines available in WEST BENGAL at

- Progressive Book Centre Kharagpur Ph: 03222-279956; Mob: 9932619526, 9434192998
- International Book Trust Kolkata
- Ph: 033-22414947. 24792343: Mob: 9830360012 Rukmani Agencies - Kolkata Ph: 033-24666173. 224839473: Mob: 9830037811
- Every Books Kolkata
- Ph: 033-22418590, 22194699; Mob: 9830162977, 8599985926
- Katha Kolkata Ph: 033-22196313; Mob: 9830257999
- Saraswati Book Store Kolkata Ph: 22198108, 22190784; Mob: 9831349473
- Chhatra Pustak Bhawan Medinipur Mob: 9609996099,9332341750
- Novelty Books Siliguri Ph: 0353-2525445; Mob: 7797938077
- Om Traders Siliguri Ph: 0353-6450299; Mob: 9434310035, 9749048104

Visit "MTG IN YOUR CITY" on www.mtg.in to locate nearest book seller OR write to info@mtg.in OR call 0124-6601200 for further assistance.





Time Allowed : 3 hours Maximum Marks: 70

#### **GENERAL INSTRUCTIONS**

- All questions are compulsory. (i)
- (ii) Q. no. 1 to 5 are very short answer questions and carry 1 mark each.
- (iii) Q. no. 6 to 10 are short answer questions and carry 2 marks each.
- (iv) Q. no. 11 to 22 are also short answer questions and carry 3 marks each.
- (v) Q. no. 23 is a value based question and carry 4 marks.
- (vi) Q. no. 24 to 26 are long answer questions and carry 5 marks each.
- (vii) Use log tables if necessary, use of calculators is not allowed.
- 1. Which forces are involved in holding the drugs to the active site of enzymes?
- 2. Write the name of monomer of PMMA and state one use of it.
- 3. Does a tetrahedral complex show geometrical isomerism? Give reason.
- 4. Why La(OH)<sub>3</sub> is more basic than  $Lu(OH)_3$ ?
- 5. Why does compound (A) given below not form an oxime?



6. The value of  $\Delta_f G^\circ$  for the formation of  $Cr_2O_3$  is  $-540 \text{ kJ mol}^{-1}$  and that of Al<sub>2</sub>O<sub>3</sub> is  $-827 \text{ kJ mol}^{-1}$ . Is the reduction of Cr<sub>2</sub>O<sub>3</sub> possible with Al?



- (iii)
  - NO<sub>2</sub>

7. In each of the following pairs of compounds, identify the compound which will undergo  $S_{\!\scriptscriptstyle\rm N} 1$ reaction.





- Write down the IUPAC names for the following 8. compounds :
  - (i) PhCH<sub>2</sub>CH<sub>2</sub>COOH





#### OR

Cyclohexanone forms cyanohydrin in good yield but 2, 2, 6-trimethylcyclohexanone does not. Explain.

- 9. Physical adsorption decreases with increase in temperature while chemical adsorption first increases and then decreases with increase in temperature. Explain.
- 10. An element has a body-centred cubic structure with edge length of 288 pm. The density of the element is 7.2 g cm<sup>-3</sup>. Calculate the number of atoms in 208 g of the element.
- 11. Non-ideal solutions exhibit either positive or negative deviations from Raoult's law. What are these deviations and why are they caused? Explain with one example for each type.
- **12.** Identify *A* and *B* in the following reactions :

(i) 
$$\underset{\text{NH}_2}{\overset{\text{NaNO}_2, \text{HCl}}{273 \text{ K}}} A \xrightarrow{\text{Cu}_2\text{Cl}_2/\text{HCl}} B$$
  
(ii)  $\text{CH}_3\text{CH}_2\text{CH}_2\text{Br} \xrightarrow{\text{NaCN}} A \xrightarrow{\text{LiAlH}_4} B$   
 $\underset{\text{NO}_2}{\overset{\text{NO}_2}{}}$ 

(iii) 
$$\xrightarrow{\text{H}_2\text{SO}_4/\text{SO}_3} \xrightarrow{\text{Fe/HCl}} B$$

13. 1 mL of methyl acetate was added to 25 mL of N/2 HCl. 2 mL of reaction mixture were withdrawn from time to time during the progress of the reaction and titrated against standard NaOH solution. Final reading was obtained by completing hydrolysis by boiling the reaction mixture. The following results were obtained :

Time (min) : 0 25 40 61  $\infty$ NaOH used : 19.24 24.20 26.60 29.50 42.10 (mL)

Show that reaction is of first order.

- **14.** Answer the following :
  - (i) Silver crystallises with face-centred cubic unit cell. Each side of this unit cell has a length of 409 pm. What is the radius of silver atom? Assume the atoms just touch each other on the diagonal across the face of the unit cell.
  - (ii) Explain how we can determine the atomic mass of an unknown metal if density and the dimension of unit cell are known.

#### OR

In a face-centred lattice of X and Y, X atoms are present at the corners while Y atoms are at face centres.

- What is the formula of the compound? (a)
- What would be the formula of the compound if **(b)** 
  - (i) one of the X atom is missing from a corner in each unit cell.
  - (ii) one atom of X from a corner is replaced by Z atom (also monovalent)?
- 15. The vapour pressure of a dilute aqueous solution of glucose is 750 mm Hg at 373 K. Calculate the molality and mole fraction of the solute.
- **16.** Answer the following questions :
  - (a) Write down the electronic configuration of the *d*-orbitals of Ti in  $[Ti(H_2O)_6]^{3+}$  ion in an octahedral crystal field.
  - (b) Why is  $[Ti(H_2O)_6]^{3+}$  ion coloured? Explain on the basis of distribution of electrons in the *d*-orbitals.
  - (c) How does the colour change on heating  $[Ti(H_2O)_6]^{3+}$  ion?
- **17.** How will you synthesise :
  - (i) 1-Phenylethanol from a suitable alkene
  - (ii) Cyclohexylmethanol using an alkyl halide by  $S_N 2$  reaction
  - (iii) Pentan-1-ol using a suitable alkyl halide?
- **18.** For the reaction,  $2NO + Cl_2 \rightarrow 2NOCl$  at 300 K following data is obtained :

Expt.	Initial concentration (mol L <sup>-1</sup> )		Initial rate
NO.	[NO]	[Cl <sub>2</sub> ]	(moll s)
1.	0.010	0.010	$1.2  imes 10^{-4}$
2.	0.010	0.020	$2.4  imes 10^{-4}$
3.	0.020	0.020	$9.6 \times 10^{-4}$

Write rate law of the reaction. What is the order of the reaction? Also calculate the specific rate constant.

- **19.** A white crystalline compound X is heated with  $K_2Cr_2O_7$  and conc.  $H_2SO_4$ , a reddish-brown gas A is evolved. On passing A into caustic soda solution, a yellow-coloured solution of B is obtained. Neutralising the solution B with acetic acid on subsequent addition of lead acetate, a yellow precipitate *C* is obtained. Identify *A*, *B*, *C* and *X*.
- **20.** Give reason for the following :
  - (a) Carboxylic acids do not give characteristic reactions of carboxyl group.



- (b) Treatment of benzaldehyde with HCN gives a mixture of two isomers which cannot be separated even by careful fractional distillation.
- (c) Sodium bisulphite is used for the purification of aldehydes and ketones.
- 21. Define the following with one example each
  - (i) addition polymers,
  - (ii) condensation polymers, (iii) copolymers.
- 22. Define the following as related to proteins (i) Peptide linkage (ii) Primary structure (iii) Denaturation.
- 23. A seminar about harmful effects of chlorofluorocarbons (CFCs) was conducted in a school in New Delhi by professor K.G. Sharma who is an ecologist. Professor explained that CFCs were used as refrigerants, propellants for aerosols, as solvents for dry cleaning etc. He explained the destruction of ozone layer by CFCs. He also explained the industrial use of freon 12.
  - (i) How will freon 12 be manufactured?
  - (ii) How does CFCs destroy ozone layer?
  - (iii) What should be done to minimise the harmful effects of CFCs?
  - (iv) Write two values that are learnt by the students?
- 24. (a) Consider the figure and answer the following questions.



- Cell 'A' has  $E_{cell} = 2$  V and cell 'B' has (i)  $E_{\text{cell}} = 1.1 \text{ V}$  which of the two cells 'A' or 'B' will act as an electrolytic cell? Which electrode reactions will occur in this cell?
- (ii) If cell 'A' has  $E_{cell} = 0.5$  V and cell 'B' has  $E_{\text{cell}} = 1.1 \text{ V}$  then what will be the reactions at anode and cathode?
- (b) Conductivity of 0.00241 M acetic acid is 7.896  $\times$  10<sup>-5</sup> S cm<sup>-1</sup>. Calculate its molar conductivity. If  $\Lambda_m^{\circ}$  for acetic acid is 390.5 S cm<sup>2</sup> mol<sup>-1</sup>, what is its dissociation constant?

OR

- (a) An acidic solution of  $Cu^{2+}$  salt containing  $0.4 \text{ g of } \text{Cu}^{2+}$  is electrolysed until all the copper is deposited. The electrolysis is continued for seven more minutes with volume of solution kept at 100 mL and the current at 1.2 amp. Calculate the gases evolved at NTP during the entire electrolysis.
- **(b)** Write down the electrode process when aqueous solutions of following halides are electrolysed:

(i) NaCl<sub>(aq)</sub> (ii)  $NaBr_{(aq)}$  (iii)  $NaF_{(aq)}$ 

- 25. (a) Write the formula of reagents used in the following reactions :
  - (i) Bromination of phenol to
  - 2,4,6- tribromophenol (ii) Hydroboration of propene and then
  - oxidation to propanol.
  - (b) Arrange the following sets of compounds in the increasing order of their property indicated:
    - p-nitrophenol, ethanol, phenol (acidic (i) character)
    - (ii) propanol, propane, propanal (boiling point)
  - Write the mechanism (using curved arrow (c) notation) of the following reaction :

$$CH_3 - CH_2 - OH \xrightarrow{(i) H^+} (ii) CH_3 CH_2 OH$$

#### OR

- (a) How would you obtain the following :
  - (i) benzoquinone from phenol (ii) 2-methylpropan-2-ol from
    - methyl magnesium bromide
  - (iii) propan-2-ol from propene?
- Name the different reagents needed to perform **(b)** the following reactions :
  - Reduction of phenol to benzene (i)
  - (ii) Dehydration of propan-2-ol to propene
  - (iii) Friedel-Crafts alkylation of anisole to *p*-methoxytoluene
  - (iv) Dehydrogenation of ethanol to ethanal
- Deduce the shapes of the following on the 26. (a) basis of VSEPR theory.
  - (i) BrF<sub>3</sub> (ii)  $I_3^-$ (iii) IF<sub>7</sub> (b) Explain
    - (i)  $BiCl_3$  is less covalent than  $PCl_3$ .
    - (ii) In HNO<sub>3</sub>, N-O bond (121 pm) is shorter than the N—OH bond (140 pm). OR
  - (a) Why does the reactivity of nitrogen differ from that of phosphorus?



- (b) Give at least one example to explain the following properties :
  - (i) Sulphuric acid is a dibasic acid.
  - (ii) Sulphuric acid is a dehydrating agent.
  - (iii) Sulphuric acid is an oxidising agent.

### SOLUTIONS

1. Substrates bind to the active site of the enzyme through a variety of interactions such as ionic bonding, hydrogen bonding, van der Waals' interaction or dipole-dipole interaction.

2. PMMA is polymethylmethacrylate, its monomer is

$$CH_2 = C - COCH_2$$

Methylmethacrylate

Use : PMMA is used in making glassy decorative materials.

**3.** No, because the relative positions of the unidentate ligands coordinated to the central metal atom are the same w.r.t. each other.

**4.** Due to lanthanoid contraction, the covalent character of hydroxides increases from  $La^{3+}$  to  $Lu^{3+}$  (Fajan's rule) and hence the basic strength decreases.

5. Glucose pentaacetate (*A*) does not have free —OH at C1, thus cannot be converted to the open chain form to give —CHO group hence, it does not form the oxime.

6. Given that,

$$2AI_{(s)} + \frac{3}{2}O_{2(g)} \rightarrow AI_2O_{3(s)}; \Delta_f G^\circ = -827 \text{ kJ mol}^{-1}...(i)$$

$$2Cr_{(s)} + \frac{3}{2}O_{2(g)} \rightarrow Cr_2O_{3(s)}; \Delta_f G^\circ = -540 \text{ kJ mol}^{-1}...(ii)$$

On substracting equation (ii) from equation (i), we get

$$2\mathrm{Al}_{(s)} + \mathrm{Cr}_2\mathrm{O}_{3(s)} \rightarrow \mathrm{Al}_2\mathrm{O}_{3(s)} + 2\mathrm{Cr}_{(s)};$$
  
$$\Delta_r G^\circ = -287 \text{ kJ mol}^{-1}$$

Since  $\Delta_r G^\circ$  of this reaction is –ve, therefore, reduction of  $\operatorname{Cr}_2O_3$  by Al is possible.

7. (i) Since 3° carbocations are more stable than 1° carbocations, therefore, *tert*-butyl chloride undergoes  $S_N1$  reaction but 1-chlorobutane does not.

(ii) Benzyl chloride readily forms benzyl carbocation which is stabilized by resonance. Thus, benzyl chloride undergoes  $S_N1$  reaction but chlorobenzene does not undergo  $S_N1$  reaction.

- 8. (i) 3-Phenylpropanoic acid
- (ii) 3-Methylbut-2-enoic acid
- (iii) 2-Methylcyclopentanecarboxylic acid
- (iv) 2, 4, 6-Trinitrobenzoic acid

Consider the following reactions :



OR

In reaction (a), cyanohydrin is formed by the nucleophilic attack of cyanide ion on the carbonyl carbon. But in reaction (b), substrate is sterically hindered due to the presence of the three methyl groups. Hence, the nucleophilic attack of cyanide ion on the carbonyl carbons does not take place. As a result, the reaction does not occur.

**9.** Physisorption is a type of adsorption in which adsorbate molecules are held to the solid surface (adsorbent) by weak van der Waals' forces. When temperature increases, these forces are weakened



resulting in decrease in extent of physisorption. In case of chemisorption, adsorbate molecules are held with adsorbent by chemical bonds with high activation energy. Initial increase in chemisoption is due to heat supplied which acts as activation energy required in chemisorption. But later it decreases due to the exothermic nature of adsorption at equilibrium.

10. Density 
$$d = \frac{MZ}{a^3 N_0}$$
  
where,  $M = \text{molar mass}$   
 $d = \text{density} = 7.2 \text{ g cm}^{-3}$   
 $N_0 = \text{Avogadro's number} = 6.02 \times 10^{23} \text{ mol}^{-1}$   
 $a = \text{edge length} = 288 \text{ pm} = 288 \times 10^{-10} \text{ cm}$   
 $Z = 2 \text{ (for bcc)}$   
 $M = \frac{dN_0 a^3}{Z} = \frac{7.2 \times 6.02 \times 10^{23} \times (288 \times 10^{-10})^3}{2}$   
 $M = 51.77 \text{ g mol}^{-1}$ 

CHEMISTRY TODAY | MARCH '18

79

Thus, moles of the element in 208 g =  $\frac{208}{51.77}$  = 4.02 Hence, number of atoms =  $4.02 \times N_0$ 

 $= 4.02 \times 6.02 \times 10^{23} = 2.42 \times 10^{24}$  atoms

11. Positive deviation : For non-ideal solution, if the vapour pressure is higher, then it is said to exhibit positive deviation. A-B interactions are weaker than A - A or B - B interactions. Due to this, vapour pressure increases which results in positive deviation. In positive deviation, intermolecular force decreases, volume increases, vapour pressures increases, enthalpy of mixing increases. Therefore,  $\Delta H_{mix} = +ve$ ,  $\Delta V_{mix} = +ve$ . e.g., ethanol + acetone and carbon disulphide + acetone show positive deviation.

Negative deviation : For non-ideal solution, if the vapour pressure is lower, then it is said to exhibit negative deviation. A-B interactions are stronger than A - A and B - B interactions. Due to this, vapour pressure decreases which results in negative deviation. In negative deviation, intermolecular force increases, volume decreases, vapour pressure decreases and heat is released. Therefore,  $\Delta H_{mix} = -ve$ ,  $\Delta V_{mix} = -ve$ e.g., phenol + aniline and chloroform + acetone show negative deviation.



- **13.** Let amount of NaOH used at time t = 0 be  $V_0$ .
- *:*.  $V_0 \propto$  HCl present in the mixture

Let amount of NaOH used at any time t be  $V_t$ .

*:*..  $V_t \propto$  HCl present + acetic acid formed.

*:*.. Concentration of acetic acid formed after time t, *i.e.*,  $x \propto (V_t - V_0)$ 

Let amount of NaOH used at  $t = \infty$ , when reaction is complete, be  $V_{\infty}$ .

 $\therefore$   $V_{\infty} \propto$  (HCl present + maximum amount of CH<sub>3</sub>COOH formed)

But maximum  $CH_3COOH$  formed  $\propto$  initial concentration of CH<sub>3</sub>COOCH<sub>3</sub>

$$\therefore \quad a \propto V_{\infty} - V_0$$
  

$$\therefore \quad (a - x) \propto V_{\infty} - V_0 - (V_t - V_0) \propto V_{\infty} - V_t$$
  

$$\therefore \quad \text{Rate law, } k = \frac{2.303}{t} \log \frac{a}{(a - x)} \text{ can be written as :}$$
  

$$k = \frac{2.303}{t} \log \frac{V_{\infty} - V_0}{V_{\infty} - V_t}$$

Here,  $V_{\infty} - V_0 = 42.10 - 19.24 = 22.86 \text{ mL}$ 

t (minute)	V <sub>t</sub>	$V_{\infty}$ – $V_t$	$k = \frac{2.303}{t} \log \frac{V_{\infty} - V_0}{V_{\infty} - V_t}$
25	24.20	42.10 - 24.20 = 17.90	$k = \frac{2.303}{25} \log \frac{22.86}{17.90}$ $= 9.785 \times 10^{-3} \text{ min}^{-1}$
40	26.60	42.10 - 26.60 = 15.50	$k = \frac{2.303}{40} \log \frac{22.86}{15.50}$ $= 9.715 \times 10^{-3} \min^{-1}$
61	29.50	42.10 - 29.50 = 12.60	$k = \frac{2.303}{61} \log \frac{22.86}{12.60}$ $= 9.767 \times 10^{-3} \min^{-1}$

Nearly constant values of rate constant prove that reaction is of first order.

14. (i) As the atoms just touch each other on the diagonal across the face of unit cell, therefore

$$b^{2} = a^{2} + a^{2} = 2a^{2}$$
  
 $b = \sqrt{2}a$  ...(i)  
Also  $b = r + 2r + r = 4r$  ...(ii)

From (i) and (ii), we get

$$4r = \sqrt{2}a \implies r = \frac{\sqrt{2}}{4}a;$$
$$r = \frac{1.414 \times 409 \text{ pm}}{4}$$
$$= 144.58 \text{ pm}$$



# Concerned about your performance in Class XII Boards?



#### Well, fear no more, help is at hand .....

To excel, studying in right direction is more important than studying hard. Which is why we created the Excel Series. These books – for Physics, Chemistry, Biology & Mathematics – have been put together totally keeping in mind the prescribed syllabus and the pattern of CBSE's Board examinations, so that students prepare and practice with just the right study material to excel in board exams.

Did you know nearly all questions in CBSE's 2017 Board Examination were a part of our Excel books? That too fully solved !

#### **HIGHLIGHTS:**

- Comprehensive theory strictly based on NCERT, complemented with illustrations, activities and solutions of NCERT questions
- Practice questions & Model Test Papers for Board Exams
- Value based questions
- Previous years' CBSE Board Examination Papers (Solved)
- CBSE Board Papers 2017 Included





Available at all leading book shops throughout the country. For more information or for help in placing your order: Call 0124-6601200 or email: info@mtg.in \*Application to read QR codes required

CHEMISTRY TODAY | MARCH '18 81

Visit www.mtg.in for latest offers

and to buy

online

(ii) We can determine the atomic mass of an unknown metal by using the formula of density of its unit cell.  $Z(No. of atoms per unit cell) \times M(Atomic mass)$ 

$$a = \frac{1}{a^3(\text{Cell edge}) \times N_4 \text{ (Avogadro's number)}}$$

By knowing density and dimension of unit cell we can calculate M, the atomic mass of metal for a given unit cell as Z is fixed and  $N_A$  is a universal constant.

#### OR

(a) No. of *X* atoms in the unit cell 
$$= 8 \times \frac{1}{8} = 1$$

No. of Y atoms in the unit cell =  $6 \times \frac{1}{2} = 3$ Formula of the compound =  $XY_3$ 

(b) (i) If one *X* atom is missing, no. of *X* atoms in the

unit cell  $=\frac{7}{8}$ 

Formula =  $X_{7/8}Y_3$  or  $X_7Y_{24}$ 

(ii) No. of X atoms = 
$$\frac{7}{8}$$
, No. of Z atoms =  $\frac{1}{8}$ 

No. of Y atoms = 3.

Formula =  $X_{7/8}Y_3Z_{1/8}$  o  $X_7Y_{24}Z$ 

15. Vapour pressure of solution  $(p_s)$  at 373 K = 750 mm Hg

Vapour pressure of pure solvent, *i.e.*, water  $(p^{\circ})$  at 373 K = 760 mm Hg

: Lowering in vapour pressure,

 $\Delta p = 760 - 750 = 10 \text{ mm Hg}$ 

Also, molar mass of glucose  $(C_6H_{12}O_6) = 180 \text{ g mol}^{-1}$ According to Raoult's law,

$$\frac{\Delta p}{p^{\circ}} = \frac{n_2}{n_1 + n_2}$$
 (Mole fraction of solute)  

$$\therefore \text{ Mole fraction of glucose} = \frac{10 \text{ mm}}{760 \text{ mm}} = 0.013$$

Molality is the number of moles of solute in 1000 g ... of water.

$$\therefore \quad \text{Number of moles of water } (n_1) = \frac{1000}{18} = 55.55 \text{ mol}$$

$$\therefore \frac{n_2}{55.55 + n_2} = 0.013$$

On calculating, we get  $n_2 = 0.73 \text{ mol/kg or } 0.73 \text{ m}$ 

**16.** (a) In 
$$[Ti(H_2O)_6]^{3+}$$
 ion,

Oxidation state of Ti = +3

There is only one electron in the *d*-orbital and its electronic configuration in an octahedral crystal field is  $t_{2g}^1 e_g^0$ .

(b) Due to d-d transition configuration changes from  $t_{2g}^1 e_g^0$  to  $t_{2g}^0 e_g^1$  and hence coloured.



(c) On heating  $[Ti(H_2O)_6]^{3+}$  ion becomes colourless as there is no ligand (H<sub>2</sub>O) left on heating. In the absence of ligand, crystal field splitting does not occur.

HO  

$$CH=CH_2$$
  
 $HO$   
 $CH=CH_2$   
 $HO$   
 $CH=CH_3$   
 $HO$   
 $HO$   
 $CH=CH_3$   
 $HO$   
 $HO$   

$$CH_3 - (CH_2)_3 - CH_2 - OH$$
  
Pentan-1-ol

**18.** Let the rate law for the reaction be  
Rate = 
$$k[NO]^{x}[Cl_{2}]^{y}$$
  
From Expt. (1), 1.2 × 10<sup>-4</sup> =  $k[0.010]^{x}[0.010]^{y}$  ...(i)  
From Expt. (2), 2.4 × 10<sup>-4</sup> =  $k[0.010]^{x}[0.020]^{y}$  ...(ii)  
Dividing Eq. (ii) by Eq. (i),  
 $\frac{2.4 \times 10^{-4}}{1.2 \times 10^{-4}} = \frac{[0.020]^{y}}{[0.010]^{y}} \implies 2 = (2)^{y} \implies y = 1$   
From Expt. (2), 2.4 × 10<sup>-4</sup> =  $k[0.010]^{x}[0.020]^{y}$  ...(iii)  
From Expt. (2), 2.4 × 10<sup>-4</sup> =  $k[0.020]^{x}[0.020]^{y}$  ...(iv)  
Dividing Eq. (iv) by Eq. (iii),  
 $\frac{9.6 \times 10^{-4}}{2.4 \times 10^{-4}} = \frac{[0.020]^{x}}{[0.010]^{x}} \implies 4 = 2^{x} \implies x = 2$   
Order of reaction =  $x + y = 2 + 1 = 3$   
Rate law for the reaction is  
Rate =  $k[NO]^{2}[Cl_{2}]$   
Considering Eq. (i) again,  
 $1.2 \times 10^{-4} = k[0.010]^{2}[0.010]$   
 $k = \frac{1.2 \times 10^{-4}}{100} = 1.2 \times 10^{2} \text{ mol}^{-2}[\text{litre}^{2} \text{ sec}^{-1}]$ 

**19.** When sodium chloride is heated with  $K_2Cr_2O_7$  and conc. H<sub>2</sub>SO<sub>4</sub>, a reddish-brown gas CrO<sub>2</sub>Cl<sub>2</sub> (chromyl chloride) is evolved.

 $[0.010]^3$ 



$$4\text{NaCl} + \text{K}_2\text{Cr}_2\text{O}_7 + 6\text{H}_2\text{SO}_4 \longrightarrow (\text{conc.})$$
  
2KHSO<sub>4</sub> + 4NaHSO<sub>4</sub> + 2CrO<sub>2</sub>Cl<sub>2</sub>↑ + 3H<sub>2</sub>O

When vapours of chromyl chloride (*A*) are passed into caustic soda (NaOH), a yellow-coloured solution is obtained due to formation of  $Na_2CrO_4(B)$ .

$$\begin{array}{c} \operatorname{CrO}_2\operatorname{Cl}_2 + 4\operatorname{NaOH} \longrightarrow \operatorname{Na}_2\operatorname{CrO}_4 + 2\operatorname{NaCl} + 2\operatorname{H}_2\operatorname{O}_{(A)} \\ (B) \end{array}$$

On subsequent addition of lead acetate solution, a yellow precipitate of  $PbCrO_4(C)$  is obtained.

 $Na_{2}CrO_{4} + (CH_{3}COO)_{2}Pb \rightarrow PbCrO_{4}\downarrow + 2CH_{3}COONa$ (B)
(C)

Thus, *A* is chromyl chloride, *B* is  $Na_2CrO_4$ , *C* is PbCrO<sub>4</sub> and *X* is a chloride (say, NaCl).

**20.** (a) Due to resonance, CO group of carboxylic acid is no more electrophilic.

(b) It is because we get two optical isomers which have same physical properties, therefore, cannot be separated by fractional distillation.

Benzaldehyde cyanohydrin

(c) Aldehydes and ketones form addition products with NaHSO<sub>3</sub> whereas impurities do not. On hydrolysis, we get pure aldehydes and ketones back.

$$CH_{3} - C - H + NaHSO_{3} \Longrightarrow CH_{3} - CH - SO_{3}Na$$

$$(H_{2}O/H^{+}) = CH_{3} - CH + NaHSO_{3}$$

**21.** (i) Addition polymers : The polymers formed by the addition reaction of a large number of unsaturated monomers are called addition polymers. *e.g.*, Polyethene.

(ii) Condensation polymers : The polymers formed by the condensation of two or more bifunctional monomers are called condensation polymers. *e.g.*, Nylon 6, 6.

(iii) **Copolymers :** The polymers made by addition polymerisation from two different compounds are known as copolymers. *e.g.*, Buna-S.

**22.** (i) **Peptide Linkage** : Proteins are the polymers of  $\alpha$ -amino acids which are connected to each other by peptide bond or peptide linkage. Chemically, peptide linkage is an amide formed between —COOH group and —NH<sub>2</sub> group. The reaction between two molecules of similar or different amino acids, proceeds through the combination of the amino group of one molecule with the carboxyl group of the other. This results in the elimination of a water molecule and formation of a peptide bond —CO—NH—.

(ii) **Primary Structure :** Proteins may have one or more polypeptide chains. Each polypeptide in a protein has amino acids linked with each other in a specific sequence and it is this sequence of amino acids that is said to be the primary structure of that protein. Any change in this primary structure *i.e.*, the sequence of amino acids creates a different protein.

(iii) **Denaturation :** Protein found in a biological system with a unique three-dimensional structure and biological activity is called a native protein. When a protein in its native form, is subjected to physical change like change in temperature or chemical change in pH, the hydrogen bonds are disturbed. Due to this, globules unfold and helix get uncoiled and protein loses its biological activity. This is called denaturation of protein.

During denaturation 2° and 3° structures are destroyed but 1° structure remains intact. The coagulation of egg white on boiling is a common example of denaturation of protein.

**23.** (i) Freon 12 is manufactured from carbon tetrachloride by Swarts reaction.

$$3CCl_4 + 2SbF_3 \xrightarrow{SDCl_5} 3CCl_2F_2 + 2SbCl_3$$
  
Freon 12

(ii) NO + O<sub>3</sub>  $\rightarrow$  NO<sub>2</sub> + O<sub>2</sub> NO<sub>2</sub> + O  $\rightarrow$  NO + O<sub>2</sub> CF<sub>2</sub>Cl<sub>2</sub>  $\rightarrow$  CF<sub>2</sub>Cl<sup>•</sup> + Cl<sup>•</sup> (In presence of light) CFCl<sub>3</sub>  $\rightarrow$  CFCl<sub>2</sub><sup>•</sup> + Cl<sup>•</sup> Cl<sup>•</sup> + O<sub>3</sub>  $\rightarrow$  ClO<sup>•</sup> + O<sub>2</sub> ClO<sup>•</sup> + O  $\rightarrow$  Cl<sup>•</sup> + O<sub>2</sub>

#### Solution Senders of Chemistry Musing

Set - 55

Ashwini Mukherjee, West Bengal

Set - 54

- Varun Gowda, Karnataka
- Anushka Singh, Rajasthan



(iii) CFCs must be replaced by other compounds which are not harmful like natural refrigerant, isobutane can be used in domestic refrigerator.

(iv) Concern and environment conservation are values learnt by the students through this seminar.

24. (a) (i) Cell B will act as electrolytic cell as it has lower emf.

The electrode reaction will be

 $Zn^{2+} + 2e^- \rightarrow Zn$  (Cathode)

 $Cu \rightarrow Cu^{2+} + 2e^{-}$  (Anode)

(ii) With the values of emf, cell *B* will act as galvanic cell due to higher emf and will push electrons into cell A. The electrode reactions will be

 $Zn \rightarrow Zn^{2+} + 2e^{-}$  (Anode)  $Cu^{2+} + 2e^{-} \rightarrow Cu$  (Cathode)  $\kappa \times 1000$  (7.896  $\times 10^{-5}$ )  $\times 1000$ 

(b) 
$$\Lambda_m^c = \frac{\kappa \times 1000}{C} = \frac{(7.000 \times 10^{-7}) \times 1000}{0.00241}$$
  
= 32.76 S cm<sup>2</sup> mol<sup>-1</sup>

$$\alpha = \frac{\Lambda_m^c}{\Lambda_m^o} = \frac{32.76}{390.5} = 8.4 \times 10^{-2}$$

$$K_a = \frac{C\alpha^2}{1-\alpha} = \frac{0.00241 \times (8.4 \times 10^{-2})^2}{1-(8.4 \times 10^{-2})} = 1.86 \times 10^{-5}$$
OR

(a) 0.4 g of  $Cu^{2+} = \frac{0.4}{31.75} = 0.0126$  g equivalent

At the same time, the oxygen deposited at anode

= 0.0126 g equivalent =  $\frac{8}{32} \times 0.0126 = 0.00315$  g mol

After the complete deposition of copper, the electrolysis will discharge hydrogen at cathode and oxygen at anode. The amount of charge passed =  $1.2 \times 7 \times 60$ 

= 504 coulomb

So, oxygen liberated =  $\frac{1}{96500} \times 504 = 0.00523$  g equivalent  $=\frac{8}{32}$  × 0.00523 = 0.001307 g mol

Hydrogen liberated = 0.00523 g equivalent

$$=\frac{1}{2} \times 0.00523 = 0.00261$$
 g mol

Total gases evolved = (0.00315 + 0.001307 + 0.00261) g mol = 0.007067 g mol

Volume of gases evolved at NTP

 $= 22400 \times 0.007067 \text{ mL} = 158.3 \text{ mL}$ 

(b) (i)  $2H^+ + 2e^- \rightarrow H_2^\uparrow$ (Cathode)  $2\text{Cl}^- \rightarrow \text{Cl}_2 + 2e^-$ (Anode)



84 CHEMISTRY TODAY | MARCH '18

(ii) 
$$2H^+ + 2e^- \rightarrow H_2^\uparrow$$
 (Cathode)

$$2Br^{-} \rightarrow Br_{2} + 2e^{-} \qquad (Anode)$$

$$(Cathode)$$

$$2OH^{-} \rightarrow H_2O + \frac{1}{2}O_2 + 2e^{-}$$
 (Anode)

25. (a) (i)  $Br_2/H_2O$ 

(ii)  $BH_3$  in THF/H<sub>2</sub>O<sub>2</sub>

(b) (i) *p*-Nitrophenol > phenol > ethanol (ii) Propanol > propanal > propane

(c) 
$$CH_3 - CH_2 - \ddot{O}H \xrightarrow{H^+} CH_3 - CH_2 - \ddot{O}H_2 \xrightarrow{CH_3 - CH_2 - \ddot{O}H_2} \xrightarrow{CH_3 - CH_2 - \ddot{O}H_2} \xrightarrow{CH_3 - CH_2 - CH_3} CH_3 - CH_2 - CH_2 - CH_3$$

$$CH_3 - CH_2 - O - CH_2 - CH_3$$

(a) (i) 
$$\underset{\text{Phenol}}{\overset{\text{OH}}{\longrightarrow}} \xrightarrow{\text{Na}_2\text{Cr}_2\text{O}_7/\text{H}_2\text{SO}_4} \xrightarrow{\text{O}}$$

(ii) 
$$CH_3 - C - CH_3 + CH_3MgBr \rightarrow OH CH_3 - C - CH_3 \leftarrow CH_3 - C - CH_3 \leftarrow CH_3 - C - CH_3 \leftarrow CH_3 - C - CH_3 -$$

Benzoquinone

(iii) 
$$CH_3CH = CH_2 \xrightarrow{H_2SO_4} CH_3 \xrightarrow{I} CH_2 \xrightarrow{I} CH_2$$

(b) (i) Zinc dust

(ii) Concentrated  $H_2SO_4$ 

2-Methylpropan-2-ol

26. (a) (i)  $BrF_3$ : This molecule has two lone pairs and three bond pairs.

To minimise the lone pair-lone pair and lone pairbond pair repulsions, these two lone pairs occupy the equatorial positions. Also, axial fluorine atoms will be bent towards the equatorial fluorine atom in order to minimise lone pair-lone pair repulsions.

**Shape :** slight bent T-shaped



(ii) Triiodide ion  $(I_3^-)$ :  $I_3^-$  ion is made up of an  $I_2$  molecule with an  $I^-$  bonded to it by means of a coordinate bond in which  $I_2$  is lone pair acceptor (Lewis acid) and  $I^-$  is lone pair donor (Lewis base). There are two bond pairs and three lone pairs in the



outer shell of central atom. To minimise the repulsive forces the three lone pairs occupy the equatorial position. The ion is therefore linear in shape with a bond angle of exactly 180°.

(iii) Iodine heptafluoride  $(IF_7)$ : This is the only common example of a non-transition element using seven orbitals for bonding giving a pentagonal bipyramidal shape.



(b) (i) By Fajan's rule,

- smaller the size of cation
- larger the size of anion
- larger the charge

then larger the polarising power and hence, greater the covalent character.

 $Bi^{3+} > P^{3+}$  (size)

Charges are identical, anions are same. Thus, polarising power is decided by size of cation. Thus,  $BiCl_3$  is less covalent than  $PCl_3$ .

(ii) HNO<sub>3</sub> exists as two resonating structures.

$$HO - N \sim O \leftrightarrow HO - N \sim O$$

(N-O) bond is resonance stabilised and it has average bond length between = and — bond, while (N-OH)bond is purely single. Thus, (N-O) bond length is shorter than the (N-OH) bond.

OR

(a) Molecular nitrogen exists as a diatomic molecule having a triple bond between the two nitrogen atoms  $(N \equiv N)$ . The bond dissociation energy is very high (945 kJ mol<sup>-1</sup>). Thus, under ordinary conditions, nitrogen behaves as inert gas. On the other hand, white

or yellow phosphorus exists as a tetratomic molecule ( $P_4$ ) having single bonds. The bond dissociation energy of P - P bond is low (213 kJ mol<sup>-1</sup>). Thus, phosphorus is much more reactive than nitrogen.

(b) (i)  $H_2SO_4$  forms two series of salts, *i.e.*, both the hydrogen atoms are replaceable.

$$\begin{array}{rcl} H_2SO_4 \rightleftharpoons H^+ + HSO_4^- \rightleftharpoons 2H^+ + SO_4^{2-} \\ H_2SO_4 + NaOH & \longrightarrow & NaHSO_4 & + H_2O \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ H_2SO_4 + 2NaOH & \longrightarrow & Na_2SO_4 & + 2H_2C \end{array}$$

$$H_2SO_4 + 2NaOH \longrightarrow Na_2SO_4 + 2H_2O$$
  
Sodium sulphate  
(Normal salt)

(ii)  $H_2SO_4$  has great affinity for water molecules and hence, acts as a dehydrating agent.

HCOOH  $\xrightarrow{\text{Conc. H}_2\text{SO}_4}$  H<sub>2</sub>O + CO Formic acid

(iii)  $H_2SO_4$  oxidises metals, non-metals and other compounds.

 $\overset{0}{\text{Cu}} + 2\text{H}_2\text{SO}_4(\text{conc.}) \longrightarrow \overset{+2}{\text{Cu}} \overset{2}{\text{Cu}} \overset{-2}{\text{Cu}} \overset{+2}{\text{Cu}} \overset{-2}{\text{Cu}} \overset{-2}{\text{Cu}} \overset{+2}{\text{Cu}} \overset{-2}{\text{Cu}} \overset$ 

### EXAM CORNER 2018

Exam	Date		
VITEEE	4 <sup>th</sup> to 15 <sup>th</sup> April		
JEE Main	8 <sup>th</sup> April (Offline), 15 <sup>th</sup> & 16 <sup>th</sup> April (Online)		
SRMJEEE	16 <sup>th</sup> to 30 <sup>th</sup> April		
Karnataka CET	18 <sup>th</sup> & 19 <sup>th</sup> April		
WBJEE	22 <sup>nd</sup> April		
Kerala PET	23 <sup>rd</sup> & 24 <sup>th</sup> April		
NEET	6 <sup>th</sup> May		
MHT CET	10 <sup>th</sup> May		
COMEDK (Engg.)	13 <sup>th</sup> May		
AMU (Engg.)	13 <sup>th</sup> May (Revised)		
BITSAT	16 <sup>th</sup> to 31 <sup>st</sup> May		
JEE Advanced	20 <sup>th</sup> May		
AIIMS	27 <sup>th</sup> May		
JIPMER	3 <sup>rd</sup> June		



ی چ

# Now, save up to Rs 2,020\*





Get digital editions of MTG Magazines on http://digital.mtg.in/











### **CREATING LEADERS OF TOMORROW**

International and National Rankings \_\_\_\_

WORLD UNIVERSITY RANKINGS 2018	18 <sup>th</sup> joint rank among 42 institutions within India (801-1000 <sup>th</sup> among 1102 institutions worldwide)	Z018	29 <sup>th</sup> rank among 42 Universities in India (251-300 <sup>th</sup> rank among Asian Universities)		
2018	210 <sup>th</sup> rank among 400 institutions from 17 countries	2017	26 <sup>th</sup> rank in 'Indian Engineering Institutions' category and 46 <sup>th</sup> rank in 'University' category		
INDIA TODAY BEST COLLEGES 2017 2017	10 <sup>th</sup> among Best Engineering Colleges	CSR-GHRDC Engineering College Survey 2017	2 <sup>nd</sup> rank among Top Engineering Colleges of Super Excellence		
CAREERS 360	• 8 <sup>th</sup> rank in Technology & Science • 52 <sup>nd</sup> rank in Research • 2 <sup>nd</sup> rank among (Private) Engineering Institutions	BRICS Rankings	137 <sup>th</sup> rank		
<ul> <li>Technology) of three UG programs in Mechanical, Civil and Electronics Engineering <ul> <li>NBA accredited UG programs</li> </ul> </li> <li>Credit transfer International Engineering Program (2+2) - Trinity College Dublin, University of New South Wales, Australia</li> <li>2018-19 Admission Open <ul> <li>a) BE, B.Tech (Bio-Technology), BE-MBA, BE (LEET), MBA, ME, M.Tech, MCA, MSc, MA, PG Diploma (Professional)- Clinical &amp; Counselling Psychology, PhD</li> </ul> </li> <li>b) UG Admission criteria: <ul> <li>BE, BE-MBA on JEE Main-2018 score.</li> <li>B.Tech (Bio-Technology) on NEET-2018 score.</li> <li>Direct admission &amp; tuition fee waiver in BE, B.Tech for 5 toppers from recognized Secondary Boards</li> <li>Scholarships worth ₹10 Crores</li> <li>I seat per discipline for J&amp;K &amp; NE students</li> </ul> </li> </ul>					
ME, MEA – April 29   B. Tech (Bio-Technology) – July 13   BE (LEE 1) – June 15   MBA – June 15   ME, M.Tech – June 30   MCA – June 30   MSc – June 30   MA – June 30   PG Diploma (Professional) – June 30   PhD – June 30					
Thapar Institute of Engineering & Technology   www.thapar.edu P.O. Box 32, Bhadson Road, Patiala, Pin -147004   Email ID : admissions@thapar.edu   Tel.: 8288008120, 8288008121.					







NAAC ACCREDITED WITH **A<sup>+</sup> GRADE** MHRD CLASSIFIED AS CATEGORY - I

### GAT-2018 GITAM ADMISSION TEST

ONLINE TEST FOR ADMISSION INTO Engineering, Architecture and Pharmacy Programs at Bengaluru, Hyderabad & Visakhapatnam campuses

**B.Tech. Programs** 

Aerospace Biotech. Civil CSE IT EEE ECE EIE Industrial Mechanical

Dual Degree (B.Tech. + M.Tech.) Programs - 6 years

ECE Mechanical

### **M.Tech. Programs**

 CST
 SE
 Cyber Forensics
 IT
 Data Sciences
 RF & ME
 VLSI
 Embedded Systems

 DSSP
 EI
 PSA
 MD
 CAD / CAM
 Industrial
 SE & NDM
 Biotech.
 FPT

B.Arch. & M.Arch. Programs

B.Pharm. & M.Pharm. Programs

#### **Record Placements**

Over 200 corporate majors visited GITAM campuses and offered **2250** jobs with lucrative packages ranging from 3.5 to 12.5 lakhs per annum in 2017-18.

Our recruiters: TCS, Accenture, Deloitte, Microsoft, Amazon, ServiceNow, Futures First, FactSet, Maruti Suzuki, Fiat to name a few.

### Entrepreneurship Development

Technology incubation and Entrepreneurship development centres provide opportunities in nurturing the young minds of GITAM for innovation and startups.

### Extra Edge

- Internships in multi national corporations.
- Science activity centre provides an exclusive platform to nourish creativity and unleash the inherent talent.
- Live Classes for GATE / GRE / CAT / Civil Services & Foreign Languages.

How to Apply : (i) Apply online at www.gitam.edu

(ii) Applications also available at selected branches of Union Bank of India, Indian Bank and Karur Vysya Bank.
 (iii) Filled-in applications along with a DD for ₹ 1,000/- (₹ 600/- for female candidates), drawn in favour of GITAM (Deemed to be University) payable at Visakhapatnam on any scheduled bank, be sent to the Director-Admissions, GITAM, Gandhi Nagar Campus, Rushikonda, Visakhapatnam-530045, Andhra Pradesh.

Online Admission Test at 48 Centres: Visit www.gitam.edu for the list of test centres

Scholarships : 
100% Fee waiver to top 10 rankers of GAT-2018 
50% Fee waiver to top 11-100 rankers of GAT-2018 
Merit - cum - Means Scholarships 
Scholarships to every M.Tech. & M.Arch. students 
Teaching Assistantships to M.Tech. students 
Scholarships to every Aerospace Engineering students 
Scholarships to every B.Pharm. & M.Pharm. students

Important Dates		
Last date for receipt of Application	s : 26 <sup>th</sup> March 2018	SCAN EST
Online Slot Booking	: 5 <sup>th</sup> to 8 <sup>th</sup> April 2018	QR 😥 🚱
Download of E-Hall Tickets	: 5 <sup>th</sup> April 2018 onward	S CODE CODE
Online (Computer based) Tests	: 11 <sup>th</sup> to 26 <sup>th</sup> April 2018	
💮 www.gitam.edu	IVRS: 0891-2866444	GITAMAdmissionTest
BENGALURU CAMPUS 080-28098008, 28098000	HYDERABAD CAMPUS 08455-221266 / 200 / 204	VISAKHAPATNAM CAMPUS 0891-2866555 / 2730177

89



### SANDEEPANA ACADEMY

A Gateway to explore more (A Unit of Sandeepana Foundation Trust) An ISO 9001: 2015 Certified Organization)

### **One Stop Solution for CBSE OUR JOURNEY OF EXCELLENCE CONTINUES......** OUR TOPPERS OF AISSCE









### **ADMISSION YEAR**

2014-15 ABOVE 90%=41 STUDENTS





### For Classes (VI, VII, VIII, IX, X, XI & XII) SANDEEPANA CLASSES **Learn from Experts**

### MEDICAL COACHING **NEET & AIIMS**

### Crash course for NEET 2018 starts from 29th March

### WHAT MAKES US NO-1 CBSE BRAND IN THE CITY

- Coaching + Study Tour guided by Teachers. 1.
- Well equipped and digitized A/C Class Rooms. 2.
- 3. Periodic Assessment of students through Well designed tests (PETs, CSTs, VSTs & Revision Test Series)
- Online Test Series for CBSE as well as all other competitive Exams. 4.
- Intimation of Test Results through SMS & Website. 5.
- 6. Hostel Facility (Separate hostels for Boys & Girls) & Transportation Facility Available.

**Centre – I : SCR – 93, KHARVEL NAGAR, BBSR, OPPOSITE TO BUDHA SOCIETY TEMPLE.** Centre – II: 191/A, KHARVEL NAGAR, BBSR, BACKSIDE TO EXHIBITION GROUND MOB.: 9438677417. 7064441570/71/72/73, PH. 0674-2534417/18 E-Mail: mail@sandeepana.co.in. www.sandeepana.co.in

90





### **VITEEE 2018** 4<sup>th</sup> April to 15<sup>th</sup> April (Computer Based Test)

Applications are invited for B.Tech Programmes

Biotech., Bioengg., Chemical, Civil, CSE, CSE (Specialisation in Digital Forensics & Cyber security), CSE (Bioinfo.), CSE (Data Analytics), CSE (Gaming Tech.), CSE (Information Security), CSE (Networks & Security), ECE, ECE (Biomedical), ECE (Embedded Systems), ECE (IOT), ECE (Sensors & Wearable Tech.), ECE (VLSI) , EEE, EIE, Electronics & Computer Engg., IT, Mech., Mech (Auto), Mech. (Energy), Production & Industrial Engg., B.Des.Ind.Design

To apply online and to get details like eligibility and fee, please visit www.vit.ac.in ONLINE APPLICATION FEE: ₹1150

### READY. STEADY. VITEEE EXPERIENCE EDUCATION THAT'S AT PAR WITH TOP UNIVERSITIES ABROAD

### **ADVANTAGE VIT**

- Curriculum for Applied Learning (CAL)<sup>™</sup>
- Project Based Learning (PBL)
- Opportunity for Semester Abroad Programme
- One module of every subject handled by Industry experts
- 35,000 students from
   60 countries
- 110 student clubs and chapters
- More than 170 Adjunct Professors from reputed universities abroad

• International Transfer Programme (ITP)

# LAST DATE FOR SUBMISSION EXTENDED TO : 15th MARCH 2018

### For enquiries contact us on our helpline number +91-416-220 2125 | +91-416-330 5555

VELLORE CHENNAI

NATIONAL INSTITUTIONAL PANKING ITAMEANEK



BHOPA

Registered R.N.I. No. 53544/1991 *Published on 1st of every month* Postal Regd. No. DL-SW-01/4044/18-20 Lic. No. U(SW)-28/2018-20 to post without prepayment of postage N.D.P.S.O.N.D-1 on 2-3rd same month

## MAKE THE RIGHT MOVE IN YOUR CAREER





ACST is the right step in the right direction that shapes your future and gives you a chance to build your career dreams under the expert guidance of Aakash. It offers scholarships up to 60% and with that a chance to seek admission in Aakash for preparation of Medical / Engineering Entrance Exams, Olympiads & Other Competitive Exams. With ACST, you can determine a bright future ahead as it brings the best in you.

### **ADMISSION OPEN**

### ONE YEAR / TWO YEAR INTEGRATED CLASSROOM COURSES 2019, 2020



For Class XI students moving to Class XII and for Class X students moving to Class XI respectively



For Class XI students moving to Class XII and for Class X students moving to Class XI respectively



For Class VIII students moving to Class IX and for Class IX students moving to Class X

### **DIRECT ADMISSION**

### **CRASH COURSE / TEST SERIES COURSE 2018**

#### for CBSE NEET-UG & AIIMS / Other Medical Ent. Exams | JEE (Main & Adv.) / Other Engg. Entrance Exams

 Dates of Commencement Crash Courses:
 15th March, 29th March, & 07th April, 2018 (Medical)
 24th March, 2018 (Engineering)

 Dates of Commencement Test Series Courses:
 29th March, 2018 (Medical)
 26th March, 2018 (Engineering)

www.aakash.ac.in — 💿 instagram.com/aakasheducation 👎 facebook.com/aakasheducation 😸 youtube.com/AakashEducation 🈏 twitter.com/aakash\_twitted 🗈 aakashinstitute.blogspot.com



CIVE A MISSED CALL: 9599280605

TOLL FREE: 1800-212-1238

SMS Aakash to **53030** 

Registered Office : Aakash Tower, 8, Pusa Road, New Delhi-110005. Ph.: (011) 47623456 E-mail: medical@aesl.in | iitjee@aesl.in | info.afs@aesl.in